

Original Research Article

Carob (*Ceratonia siliqua* L.) fruit hydro-alcoholic extract alleviates reproductive toxicity of lead in male mice: Evidence on sperm parameters, sex hormones, oxidative stress biomarkers and expression of *Nrf2* and *iNOS*

Ali Soleimanzadeh^{1,*}, Mehdi Kian¹, Sajjad Moradi¹, Soraya Mahmoudi²

¹Department of Theriogenology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

²Department of Pathobiology, Faculty of Veterinary medicine, Urmia University, Urmia, Iran

Article history:

Received: Feb 27, 2019

Received in revised form:

Apr 16, 2019

Accepted: May 26, 2019

Vol. 10, No. 1, Jan-Feb 2020,
35-49.

*** Corresponding Author:**

Tel: +98-44-32774737

+98-44-32770508

Fax: +98-44-32777099

+98-44-32271926

a.soleimanzadeh@urmia.ac.ir

Keywords:

Ceratonia siliqua

Pb poisoning

Oxidative stress

Lipid peroxidation

Nrf2

Inducible nitric oxide synthase

Male reproductive system

Abstract

Objective: Carob (*Ceratonia siliqua* L.) is an evergreen tree with fruits that have potent antioxidant activity. The aim of this study was to investigate alleviative effects of carob fruit hydro-alcoholic extract (CFHAE) against reproductive toxicity induced by lead (Pb) in male mice.

Materials and Methods: Forty-two NMRI adult male mice were randomly categorized into 7 groups (N=6). Group I was the control group and received no treatment. Group II was the sham group and received 0.2 ml distilled water per day. Group III (Pb group) received Pb acetate 1000 ppm/kg/day. Groups IV and V received CFHAE 500 and 1000 mg/kg/day, respectively. Groups VI and VII received both Pb 1000 ppm/kg/day and CFHAE at doses of 500 and 1000 mg/kg/day, respectively at the same time. The groups were treated by gavage. After 35 days, sperm parameters (count, motility, morphology, viability, DNA damage, and teratozoospermia index), total antioxidant capacity (TAC), reduced glutathione content (GSH), antioxidant enzymes (SOD, CAT, and GPx) activity, MDA levels, and sex hormones (FSH, LH, and testosterone) concentrations in serum, testicular expression of *Nrf2* and *iNOS* genes and histopathological alterations were evaluated.

Results: Our findings revealed that co-administration of CFHAE with Pb significantly ($p < 0.05$ to $p < 0.01$) improved sperm parameters, elevated sex hormones, TAC, GSH content, and antioxidant enzymes activity of serum, decreased serum MDA levels, and down-regulated testicular expression of *Nrf2* and *iNOS* genes compared with Pb group. Also, CFHAE ameliorated histopathological alterations in testis tissue caused by Pb.

Conclusion: CFHAE can alleviate reproductive toxicity following Pb exposure in male mice.

Please cite this paper as:

Soleimanzadeh A, Kian M, Moradi S, Mahmoudi S. Carob (*Ceratonia siliqua* L.) fruit hydro-alcoholic extract alleviates reproductive toxicity of lead in male mice: evidence on sperm parameters, sex hormones, oxidative stress biomarkers and expression of *Nrf2* and *iNOS*. Avicenna J Phytomed, 2020; 10(1): 35-49.

Introduction

Infertility is considered one of the most important issues in married couples. It has

been estimated, a male factor infertility plays a role in about 50% of all infertility cases (Sigman, 2007). Environmental

exposures to toxicants are directly associated with male infertility (Wong and Cheng, 2011). Lead (Pb) is an abundant toxic heavy metal in the environment (Bae et al., 2001; Shotyk and Le Roux, 2005). This heavy metal is widely used in industries and present in many products and some cosmetics (Ayinde et al., 2012; Chowdhury, 2009). Human exposure to Pb could occur via air, water, soil, food, and consumer products (Hammond, 1977).

Evidence elucidated that Pb is toxic for male reproductive system. Exposure to Pb causes generation of reactive oxygen species (ROS), lipid peroxidation and depletion of antioxidants which lead to induction of oxidative stress in testis tissue (Moniem et al., 2010; Reshma Anjum et al., 2011; Ayinde et al., 2012; Reshma Anjum et al., 2017; Elgawish and Abdelrazek, 2014). Also, Pb attenuates sperm parameters and diminishes the activity of spermatozoa (Chowdhury, 2009). Pb exposure impairs sperm normal function by decreasing the levels of sperm intracellular cyclic adenosine monophosphate and calcium and reducing tyrosine phosphorylation of sperm proteins (He et al., 2016). Many studies indicated that administration of antioxidant herbal plants can alleviate oxidative stress induced by Pb in experimental models (El-Nekeety et al., 2009; Dorostghoal et al., 2014; Dkhil et al., 2016; Soleimanzadeh et al., 2018).

Carob (*Ceratonia siliqua L.*) which belongs to the *Fabaceae* family is an evergreen tree cultivated in the Mediterranean countries (Dakia et al., 2007; Özcan et al., 2007). This plant grows widely in Fars province, Iran (Mokhtari et al., 2012; Vafaei et al., 2018). The carob fruit is a brown pod with length of 10–25 cm that consists of 80–90% pod and 10–20% seed by weight (Naghmouchi et al., 2009; Tetik et al., 2011). Carob fruit possesses high amounts of antioxidant compounds such as flavonoids and polyphenols and has strong radical scavenging activity (Custódio et al., 2009; Kumazawa et al., 2002; Owen et al., 2003; Papagiannopoulos et al., 2004). Previous

studies indicated protective effects of carob against oxidative stress in different organs (Abdel-Rahman et al., 2018; Rahman et al., 2016; Rtibi et al., 2016a, 2016b, 2015; Suzek et al., 2017; Vafaei et al., 2018).

In Persian traditional medicine, carob fruit is used as an aphrodisiac and to increase human semen volume to treat male infertility. Recently, it has been demonstrated that aqueous extract of carob pod has a beneficial effect on male reproductive parameters in infertile mice (Vafaei et al., 2018). In another study done by Mokhtari et al., administration of carob seeds hydro-alcoholic extract increased levels of sex hormones and sperm density in seminiferous tubules (Mokhtari et al., 2012). Also, it has been reported that carob pods extract has a possible positive influence against Pb toxicity and may interfere with the absorption of Pb (Said et al., 2017). Hence, in the present study, possible alleviative effects of carob fruit hydro-alcoholic extract (CFHAE) on Pb - induced oxidative damages on male reproductive parameters, were evaluated.

Materials and Methods

Animals

Forty-two, 8-10-week-old NMRI male mice (30±6 g) were purchased from the animal house of the Urmia Medical University, Iran. The animals were maintained under standard laboratory conditions (with 12:12 hr light/dark cycle at 25±2°C). They were acclimatized for one week before the study and were provided with tap water and standard laboratory diet *ad libitum*. The experimental protocols were approved by the Animals Ethics Committee at Urmia University (AECVU-170-2018).

Plant extract preparation

Carob fruits were purchased from a Persian herbal market and their identity was confirmed in Natural Resource Center (Herbarium No. 7123). The fruits (seeds+pods) were ground to a fine powder

and soaked in 96% ethanol (Merck, Darmstadt, Germany) and water (50:50 ratio) for 72 hr. Whatman filter No. 40 was utilized to filter the extract; after that, the extract was concentrated at 50°C using a rotary evaporator (Heidolph Laborota 4000 efficient, Schwabach, Germany). The remaining brown pasty extract was dried in an oven at 40°C (Mokhtari *et al.*, 2012). Eventually, the CFHEA was maintained at -20°C in order to use in subsequent *in-vitro* and *in-vivo* experiments.

HPLC analysis to identify and quantify phenolic and flavonoids compounds

Based on a modified version of Singleton's method by Dewanto *et al.* (Dewanto *et al.*, 2002), the Folin-Ciocalteu reagent was used to analyze the phenolic contents. One aliquot (0.125 ml) of a desirable methanolic extract which was diluted, was poured into 0.5 ml of deionized water which contained 0.125 ml of Folin-Ciocalteu reagent. Then, 1.25 ml of 7% Na₂CO₃ was added to the mixture, and the solution was shaken and left for 6 min before adding Na₂CO₃. In order to increase the final volume of the solution to 3 ml, deionized water was added. Afterwards, the mixture was totally stirred. The absorbance was read at 760 nm versus prepared blank following incubation for 90 min at 23°C.

Agilent Technologies 1100 series liquid chromatograph (Reversed phase HPLC) coupled with an UV-vis multi-wavelength detector, was adopted to analyze the phenolic compound. With a 250×4.6 mm, 4 µm Hypersil ODS C18 reversed-phase column at ambient temperature, the separation was performed. The injected volume was 20 µl and prior to injection, the samples were filtered through a 0.45 µm membrane filter and peaks were monitored at 280 nm. The phenolic compounds were distinguished from other components based on their retention duration, spectral features of the peaks versus the standards and transfixing the sample with standards. The Analyses were carried out with three replications.

Experimental design

The animals were categorized into 7 groups and each group included 6 animals. Group I was the control group and received no treatment. Group II was the sham group and received distilled water per day by gavage. Group III (Pb group) received Pb acetate (Sigma-Aldrich, USA; CAS Number 6080-56-4) 1000 ppm/kg/day by gavage. Groups IV and V received CFHAE, 500 and 1000 mg/kg/day, respectively by gavage. Groups VI and VII received Pb 1000 ppm/kg/day and CFHAE at doses of 500 and 1000 mg/kg/day, respectively by gavage at the same time. The animals received treatments for 35 consecutive days. The dose of Pb was chosen based on Soleimanzadeh *et al.* (2018).

Plasma sampling

After 35 days, animals were anesthetized using intraperitoneal administration of a combination of 2% xylazine (10 mg/kg, Alfasan, Woerden, Holland) and 5% ketamine hydrochloride (90 mg/kg, Rotexmedica, Trittau, Germany) according to Kanter (2010) method. Then, animals were sacrificed by cutting their necks, blood samples were collected in laboratory tubes and centrifuged at 3000 g for 15 min. The separated serum samples were stored in microtubes at -20°C until used for biochemical and hormonal evaluations (Soleimanzadeh *et al.*, 2018).

Collection of epididymal sperms

Sperm samples were obtained from the cauda epididymis of the testes of each mice. The cauda epididymis was excised into small pieces and placed in Petri dishes containing 1 ml of Human Tubal Fluid (HTF) medium at 37°C for 30 min.

Sperm motility

Assessment of the sperm motility was performed according to the WHO laboratory manual protocol for the examination of human semen (2010). In brief, 10 µl of the sperm suspension was placed on a semen analysis chamber. A

minimum of five microscopic fields were assessed in terms of sperm motility on at least 200 sperm for each animal.

Sperm viability

Eosin-Nigrosine staining was used to assess sperm viability according to the WHO protocol (World Health Organization, 2010). Briefly, Eosin (Merck, Darmstadt, Germany) and Nigrosine (Merck, Darmstadt, Germany) were prepared in distilled water. One volume of sperm suspension was mixed with two volumes of 1% eosin. After 30 sec, an equal volume of nigrosine was added to this mixture. Thin smears were then prepared and observed under a light microscope (Model CHT, Olympus optical Co. Ltd., Tokyo, Japan) at 1000X magnification. Viable sperm remained colorless while nonviable sperm was stained red.

Teratozoospermia index (TZI)

The TZI or multiple anomalies index is the number of defects per abnormal spermatozoon. Each abnormal spermatozoon may have one to four abnormalities including head, neck/mid piece and tail defects or presence of cytoplasmic residues. TZI values have been read between 1.00 (each abnormal spermatozoon has only one defect) and 3.00 (each abnormal spermatozoon has a head, midpiece, and tail defects) as described by Krassas et al. (Krassas et al., 2008).

Assessment of DNA damage

Sperm suspensions were washed 3 times with 5 ml of phosphate buffered solution (PBS) and centrifuged at 3000 rpm for 5 min. Thin smears were prepared from the sperm solution and allowed to air-dry. To test sperm DNA integrity, the smears were stained with Acridine-Orange (AO). The AO staining was performed according to a protocol described by Tejada et al. (Tejada et al., 1984). In brief, the smears were fixed for 24 hr in methanol/acetic acid (3:1) at 4°C and stained with AO solution (0.19%

in phosphate-citrate buffer, pH 2.5) for 10 min. The slides were gently washed by distilled water for 5 min and air dried. The stained smears were then observed under a fluorescence microscope (Model GS7, Nikon Co., Tokyo, Japan) at 1000X magnification. Three types of staining patterns were considered in sperm head; green spermatozoa (double-stranded DNA), and yellow and red spermatozoa (single-stranded DNA). At least 100 spermatozoa per slide were counted to evaluate the percentage of double-stranded DNA in the spermatozoa (Tejada et al., 1984).

Estimation of total antioxidant capacity (TAC)

The TAC of the serum was estimated by ferric reduction antioxidant power (FRAP) assay (Benzie and Strain, 1999). Here, 100 µl of cellular supernatant was added to 1 ml of fresh ferric reducing antioxidant power reagent (FRAP; Tripiridyltriazine; Merck) and incubated at 37°C for 10 min in the dark. Reading of the blue-colored reagent was done at 595 nm every 20 sec for 10 min. An aqueous solution of FeII (FeSO₄·7H₂O) and appropriate concentrations of freshly prepared ascorbic acid were used as blank and standard solutions, respectively. The results were expressed as µmol of TAC per L serum.

Estimation of reduced glutathione (GSH)

Glutathione (GSH) was assayed using a modified method described by Beutler et al. (1963). In brief, 0.9 ml distilled water and 1.5 ml of precipitating reagent were added (3.34 g metaphosphoric acid, 0.4 g EDTA, and 60.0 g sodium chloride) to 0.1 ml of sample. The tubes were vortexed and left for 5 min at room temperature. The solution was centrifuged for 15 min at 4000 rpm at 4°C. Next, 4.0 ml of phosphate solution (0.3 M disodium hydrogen phosphate) and 0.5 ml 5-50-dithiobis-(2-nitrobenzoic acid) (DTNB) (80 mg in 1% sodium citrate) were mixed with 1.0 ml supernatant. The absorbance of the mixture containing a

yellow color complex was read immediately at 412 nm spectrophotometrically. A standard curve for GSH was prepared and GSH concentration in the experimental samples was extrapolated from the standard curve. The results were expressed as U/L serum.

Measurement of enzymatic antioxidants activity

The activity of superoxide dismutase (SOD) was measured in terms of inhibition of the nicotinamide adenine dinucleotide (reduced)-phenazinemetosulphate-nitrobluetetrazolium reaction system according to the Nishikimi *et al.* (1972) method. The catalase (CAT) activity was determined based on the reduction of dichromate in acetic acid to chromic acetate when heated in the presence of H₂O₂ according to Sinha (1972). The activity of glutathione peroxidase (GPx) was estimated according to the method described by Paglia and Valentine (1967) based on the fact that GPx catalyzed the oxidation of glutathione by cumene hydroperoxide. The results were presented as U/L serum.

Lipids peroxidation measurement

The rate of lipid peroxidation in the serum samples was estimated by determination of malondialdehyde (MDA) using the thiobarbituric acid reactive substances (TBARS) test (Placer *et al.*, 1966). The results were presented as $\mu\text{mol/L}$.

Hormonal assessments

Serum concentration of testosterone was measured by enzyme-linked immunosorbent assay (ELISA) as described in the instructions of the manufacturer's kit (Demeditec Diagnostics GmbH, Germany). The results were presented as $\mu\text{mol/L}$ serum.

Serum levels of LH and FSH were determined by ELISA using specific commercial kits (Amersham, Buckinghamshire, UK) according to

Loraine and Bell, 1971 methods. The results were presented as mIU/ml serum.

Histopathological evaluations

Right testes of animals were fixed in formalin 10% solution. Then, the samples were dehydrated by a graded series of ethanol and embedded in paraffin. Thin sections were cut by microtome into 7 μm thickness. Next, prepared sectioned were stained by hematoxylin and eosin (H&E) staining according to the method described by Suvarna *et al.* (2018). The stained sections were evaluated for histopathological alterations by using a light microscope (Olympus Model BH-2, Tokyo, Japan).

RNA extraction

Using Cinna Pure-RNA kit (CinnaGen, Cat. No. PR891620, Iran), the total RNA of the left testis of mice was extracted. ND-2000 spectrophotometer (Nanodrop Technologies, Thermo Fisher Scientific, USA) was used to determine concentration and purity of the total RNA and then the total RNA was stored at -70°C until cDNA synthesis.

cDNA synthesis

Using random hexamers primers (Primer+dNTP Mix+Nuclease-free water), first strand cDNA was synthesized from 10 μg of total RNA and cDNA Synthesis Mix (M-MLV Reverse Transcriptase+10X Buffer M-MuLV+Nuclease-free Water) (Vivantis Technologies, Cat. No. RTPL12 100 app, Malaysia) was incubated at 42°C for 60 min and the reaction was terminated by incubating at 85°C for 5 min followed by cooling at 4°C . Synthesized cDNA was stored at -20°C until processed. The electrophoresis was performed with the PCR products to verify the primers specificity (Figure 1).

Quantitative real-time PCR assay

Using SinaSyber Blue HF- qPCR Mix (CinnaGen, Cat. No. MM2171, Iran) on a StepOne™ real-time PCR system (Applied

Carob fruit ameliorates Pb toxicity in reproductive system of male mice

Biosystems, USA) using quantitative real-time polymerase chain reaction (qRT-PCR), the mRNA expression levels of *Nrf2*, *iNOS* and *18S rRNA* were quantified. The characteristics of the primer pairs used in the present study, are presented in Table 1. Reactions included 12.5 μ l of SinaSyber Blue HF- qPCR Mix (CinnaGen, Cat. No. MM2171, Iran), 0.25 μ l of each primer (2 μ M), 2 μ l of cDNA, 9.5 μ l of nuclease-free water at 95°C for 10 s, followed by 40 cycles of 95°C for 10 s, and 60°C for 30 s.

A standard curve from the cDNA reaction mixture was used to detect PCR efficiencies. The efficiencies of PCR were between 90 and 110 percent. The amounts for R2 for all curves were observed from 0.997 to 0.999. The amounts of CT and expression of the relative expression level of the target gene as 2-CT were used for quantification of the product of PCR. Following, the CT was measured Δ CT after

subtracting CT of the housekeeping gene (*18S rRNA*), from CT of the target gene.

Statistical analysis

Statistical analysis was performed using SPSS version 21 (IBM Co., Chicago, USA). The data were analyzed by one-way ANOVA and the Tukey test was used as *post hoc*. A p value of less than 0.05 was considered significant.

Results

Phytochemical analysis

The phytochemical investigation using RP-HPLC method revealed various phenolic and flavonoid compounds in CFHAE, as shown in Table 2. The compounds are including gallic acid, caffeic acid, chlorogenic acid, rutin, coumaric acid quercetin, cinnamic acid, and apigenin.

Table 1. The characteristics of primer pairs used in the present study

Name/gene	Primer sequence	Ref. sequence (accession number acc. to GeneBank)	Product length (base pairs, bp)
<i>iNOS</i>	F: 5' CACCTTGGAGTTCACCCAGT 3' R: 5' ACCACTCGTACTTGGGATGC 3'	NM_010927	170 bp
<i>Nrf2</i>	F: 5' CACCTTGGAGTTCACCCAGT 3' R: 5' ACCACTCGTACTTGGGATGC 3'	NM172086	263 bp
<i>18S rRNA</i>	F: 5' GCAATTATCCCATGAACG 3' R: 5' GGCCTCACTAAACCATCCAA 3'	NR_003278	123 bp

iNOS: Inducible Nitric Oxide Synthase; *Nrf2*: nuclear factor erythroid 2-related factor 2




Figure 1. Primer specificity and cDNA synthesis products. A) *Nrf2* (263 bp), B) and *iNOS* (170 bp) gene primer; Lane M: 100 bp DNA ladder, Lane 1, Control group; Lane 2, Pb group; Lane 3, Sham group; Lane 4, CFHAE 500 group; Lane 5, CFHAE 1000 group; Lane 6, Pb+CFHAE 500 group; Lane 7, Pb+CFHAE 1000 group.

Effects on sperm parameters

Data summarized in Table 3 show that exposure to Pb significantly ($p < 0.01$ for all cases) attenuated sperm parameters compared to the control group but co-administration of CFHAE and Pb significantly ($p < 0.05$ to $p < 0.01$) enhanced these parameters compared with Pb group (Figures 2-4).

Effects on TAC, GSH content and antioxidant enzymes activity in serum

Exposure to Pb significantly ($p < 0.01$ for all cases) decreased TAC, GSH content and activities of enzymatic antioxidants (SOD, CAT, and GPx) in serum compared to

control and sham groups while co-administration of CFHAE and Pb significantly ($p < 0.05$ to $p < 0.01$) and dose-dependently increased them in comparison with Pb group (Table 4).



Figure 2. Sperm viability in groups; A) Viable sperm (colorless), B) Dead sperm (red); (eosin/nigrosine, 1000X).




Figure 3. Morphologically normal and abnormal sperms in groups; normal (black arrows), and abnormal (white arrows); (eosin/nigrosine, 1000X).




Figure 4. DNA damage in groups; A) Normal sperm (green), B) Damaged DNA (yellow); (AO, 400X).

Table 2. The amounts of total phenolic and flavonoid compounds in CFHAE as determined by HPLC analysis.

Compounds	mg/kg
Gallic acid	408.37
Caffeic acid	100.58
Chlorogenic acid	123.58
Rutin	89.25
Coumaric acid	121.81
Quercetin	88.68
Cinnamic acid	20.80
Apigenin	62.38

Effects on serum lipid peroxidation

The animals that were only treated with Pb had a significant ($p < 0.01$) reduction in serum MDA level compared to the control group. However, co-administration of CFHAE and Pb, significantly ($p < 0.05$) increased MDA levels (Figure 5).

Effects on sex hormones

Hormonal evaluations showed a significant ($p < 0.01$ for all cases) decrease in concentrations of sex hormones in Pb group compared to control and sham groups while co-administration of CFHAE with Pb significantly increased sex hormones levels in comparison to Pb group ($p < 0.05$ to $p < 0.01$) (Table 5).




Figure 5. The serum mean levels MDA in different groups. Values represent means \pm SEM. ** $p < 0.01$ vs. control; # $p < 0.05$ vs. Pb.

Carob fruit ameliorates Pb toxicity in reproductive system of male mice

Table 3. Comparison of sperm parameters in different groups.

Groups	Control	Sham	Pb	CFHAE 500	CFHAE 1000	Pb+CFHAE 500	Pb+CFHAE 1000
Sperm count (10 ⁶)	27.11±1.36	26.35±1.20	11.77±0.81**	28.13±1.63	27.15±1.09	19.17±1.44##	20.42±1.02##
Sperm viability (%)	81.08±0.95	81.10±1.82	57.84±1.16**	82.28±0.77	83.71±1.50	70.03±1.27##	75.94±1.49##
Sperm motility (%)	76.24±1.33	75.28±1.21	51.49±0.78**	80.34±0.90*	78.65±0.61	63.50±1.38##	60.71±1.05##
Sperm morphology (%)	90.37±1.71	88.23±0.58	76.48±1.29**	89.47±1.40	91.20±1.01	86.34±0.68##	85.97±1.22##
DNA damage (%)	3.08±0.61	3.24±0.41	18.79±0.48**	3.08±0.71	3.04±1.15	13.59±0.96#	9.14±1.23##
TZI (%)	1.12±0.32	1.14±0.37	1.89±0.59**	1.11±0.30	1.13±0.74	1.60±0.41#	1.35±0.35##

TZI: Teratozoospermia index. Values represent means±SEM. *p<0.05, **p<0.01 vs. control; #p<0.05, ##p<0.01 vs. Pb.

Table 4. The mean amounts of antioxidant activity and lipid peroxidation in different groups

Groups	Control	Sham	Pb	CFHAE 500	CFHAE 1000	Pb+CFHAE 500	Pb+CFHAE 1000
TAC (µmol/L)	3.21±0.81	3.18±0.70	0.41±0.29**	3.32±0.43	3.45±0.38*	0.97±0.41##	1.61±0.37##
GSH (U/L)	81.19±1.48	80.37±0.56	44.25±1.17**	81.08±1.79	81.75±1.22	54.69±0.44##	62.95±1.62##
SOD (U/L)	1637±1.39	1620±1.81	1124±1.72**	1652±1.05	1681±1.19	1318±1.30##	1394±0.93##
CAT (U/L)	1.29±0.39	1.28±0.33	0.64±0.41**	1.30±0.61	1.37±0.56	0.85±0.42#	1.04±0.33##
GPx (U/L)	7.45±1.12	7.41±1.75	2.37±1.02**	7.42±1.34	7.63±1.60	4.85±0.37##	5.30±0.51##

TAC: Total antioxidant capacity, GSH: Glutathione SOD: Superoxide dismutase, CAT: Catalase, GPx: Glutathione peroxidase. Values represent means±SEM. *p<0.05, **p<0.01 vs. control; #p<0.05, ##p<0.01 vs. Pb.

Table 5. The serum mean concentrations of sexual hormones in different groups

Groups	Control	Sham	Pb	CFHAE 500	CFHAE 1000	Pb+CFHAE 500	Pb+CFHAE 1000
FSH (mIU/ml)	4.40±0.18	4.38±0.18	1.52±0.18**	4.49±0.18	4.45±0.18	2.95±0.18##	3.12±0.18##
LH (mIU/ml)	3.20±0.66	3.21±0.66	1.64±0.66**	3.29±0.66	3.41±0.66	2.35±0.66#	2.70±0.66##
Testosterone (µmol/L)	6.07±1.49	5.89±1.49	3.19±1.49**	6.11±1.49	6.28±1.49	4.79±1.49##	5.10±1.49##

Values represent means±SEM. **p<0.01 vs. control; #p<0.05, ##p<0.01 vs. Pb.

Histopathological findings

In the control group, seminiferous tubules, interstitium, Leydig cells and stromal elements have normal structures. Seminiferous tubules lined by multilayered epithelium contained Sertoli cells, spermatogonia (type A and B), primary spermatocytes, secondary spermatocytes, spermatids and spermatozoa (Figure 6A).

The sham group was similar to the control group and had a normal structure (Figure 6B). In the Pb group, seminiferous tubules were thin with the dilated lumen and specific degenerative changes. The basal membrane was irregular and disrupted. The Leydig cells were reduced in interstitial tissue. Spermatogenic cells which included spermatogonia, spermatocyte and

spermatids, were decreased and remaining cells had pyknotic nuclei and no Sertoli cells were observed. Spermatozoa were not present in the lumen which indicated suppression of spermatogenesis. Exfoliated cells were obvious in the lumen (Figure 6C). In groups treated with CFHAE 500 and 1000 mg/kg, seminiferous tubules with a normal different type of germ cells were seen (Figures 6D and E). In the group treated with Pb and CFHAE 500

mg/kg/day, degenerative changes caused by Pb were reduced after treatment with CFHAE. Also, increase of spermatogonia cells in seminiferous tubules is obvious (Figure 6F). In the group treated with Pb and CFHAE 1000 mg/kg, germinal cells in all the tubules were increased. Basement membrane structure was normalized and marked increases in spermatozoa in the lumen can be seen (Figure 6G).




Figure 6. Light micrographs of testis tissue exhibiting protective effects of co-administration of CFHAE against Pb toxicity in testis. (A): Control, (B): Sham. (C): Pb, (D): CFHAE 500, (E): CFHAE 1000, (F): Pb+CFHAE 500, and (G): Pb+CFHAE 1000. H&E staining ($\times 400$).

Effects on expression of *Nrf2* and *iNOS* genes in testis

The results in Figure 7 show that Pb significantly ($p < 0.01$) increased *Nrf2* expression compared with control group. Also, co-administration of CFHAE with Pb significantly ($p < 0.05$ at CFHAE 500 mg/kg and $p < 0.01$ at CFHAE 1000 mg/kg) decreased its expression compared with Pb group.

Data in Figure 8 show that Pb group exhibited a significant ($p < 0.01$) increase in *iNOS* expression in comparison with the control and sham groups. However, co-administration of CFHAE with Pb significantly ($p < 0.05$ at CFHAE 500 mg/kg and $p < 0.01$ at CFHAE 1000 mg/kg) decreased the expression of *iNOS* compared to Pb group.




Figure 7. The densitometry analyses of *Nrf2*, which were normalized, based on the expression level of 18srRNA. ** $p < 0.01$ vs. control; # $p < 0.05$, ## $p < 0.01$ vs. Pb.

Carob fruit ameliorates Pb toxicity in reproductive system of male mice




Figure 8. The densitometry analyses of iNOS, which were normalized, based on the expression level of 18srRNA. ** $p < 0.01$ vs. control; # $p < 0.05$, ## $p < 0.01$ vs. Pb.

Discussion

In the present study, our findings indicated that administration of CFHAE improved sperm parameters, elevated sex hormones, TAC, GSH content and antioxidant enzymes activity of serum, decreased serum MDA levels, down-regulated testicular expression of *Nrf2* and *iNOS* genes, and also ameliorated histopathological alterations in testis tissue caused by Pb.

Similar to previous studies (Dorostghoal et al., 2014; Mabrouk and Ben Cheikh, 2014; Reshma Anjum et al., 2017, 2011), our results showed that Pb attenuates sperm parameters. These findings may be caused as Pb can pass through the blood-testis barrier (Creasy, 2001), induces oxidative stress and causes lipid peroxidation and DNA damages (Zhang et al., 2014) which results in impairment of spermatogenesis. Since testis tissue possesses high content of polyunsaturated lipids in its cells' membrane, it is susceptible to oxidative stress (Mishra and Acharya, 2004). Products of lipid peroxidation detrimentally influence membrane function and motility of sperms which leads to diminishing their fertility potential (Choudhary et al., 2010).

Administration of Pb decreased serum TAC, GSH content and activity of antioxidant enzymes. The mechanism

through which Pb altered antioxidant enzymes activity, is mediated by direct inhibition of functional SH groups in antioxidant enzymes and/or by binding to metal cofactors of these enzymes such as Cu, Zn and Mn (Patra et al., 2011). Also, MDA level increased in Pb group. These results are in line with prior reports (Dkhil et al., 2016; Dorostghoal et al., 2014; Patra et al., 2011; Reshma Anjum et al., 2017).

Effects of Pb on LH and FSH are contradictory. As some studies, in line with the present research, reported a reduction of these hormones following exposure to Pb (Ayinde et al., 2012; Dorostghoal et al., 2014), other studies reported increases in them (Anjum and Reddy, 2015). The difference in various reports is attributed to the factors such as Pb concentration, exposure interval, and physiological status of the testis and reproductive axis (Gandhi et al., 2017).

The reduction in the concentration of serum testosterone in the present research, is in accordance with prior studies (Anjum and Reddy, 2015; Dkhil et al., 2016; Mabrouk and Ben Cheikh, 2014; Reshma Anjum et al., 2017). Possibly, Pb exposure adversely affected hypothalamic–pituitary–gonadal axis (Gandhi et al., 2017). Researchers revealed that exposure to Pb degenerates gonadotrophic cells of the pituitary gland (Hamadouche et al., 2013). Moreover, Pb induces extrinsic apoptosis signal pathway in Leydig cells (He et al., 2017). Oxidative damages make Leydig cells less sensitive to LH which can affect testosterone synthesis (Glade and Smith, 2015). Pb suppresses the activity of steroidogenic enzymes in Leydig cells. Thereby, another reason of decrement of testosterone might be due to inhibition of androgen synthesis by Pb (Huang and Liu, 2004; Liu et al., 2001).

Exposure to Pb induced expression of *iNOS* gene in testis tissue. *iNOS* catalyzed the synthesis of nitric oxide (NO) from L-arginine (Aktan, 2004). NO incorporates with superoxide anion in the formation of peroxynitrite anion (ONOO^-), a strong

ROS which is able to cause membrane lipid peroxidation (Moneim, 2015). Dkhil *et al.* (2016) reported that Pb exposure increases generation of NO in rats' which is in line with our results.

Nrf2 is the primary regulator of antioxidant response and participates in protection against oxidative stress (Lindl and Jordan-Sciutto, 2008). Nrf2 induces the expression of phase II antioxidant enzymes in the nucleus (Jung and Kwak, 2010). It has been found that defective Nrf2 signaling causes spermatogenesis defects (Yu *et al.*, 2012). Exposure to heavy metals results in activation of the Nrf2 pathway (Simmons *et al.*, 2011). Our study indicated that Pb up regulates *Nrf2* gene expression. Wang *et al.* (2012) showed that exposure to Pb up-regulates expression of Nrf2 in testis which is in agreement with our results.

To the best of the authors knowledge, the present research is the first study on ameliorative effects of CFHAE against Pb-induced reproductive toxicity. The present research showed that co-administration of CFHAE improved sperm parameters in animals treated with Pb. Similar to our findings, prior studies showed that carob aqueous extract has beneficial effects on sperm parameters in experimental animals (Ata *et al.*, 2018; Vafaei *et al.*, 2018). These effects may be due to antioxidant properties of carob fruit. CFHAE enhanced antioxidant activity of serum which is similar to data reported by Rtibi *et al.* (Pedersen, 2001). Also, many studies indicated that administration of carob increases antioxidant enzymes in different organs (Abdel-Rahman *et al.*, 2018; Rtibi *et al.*, 2016a, 2016b, 2015; Suzek *et al.*, 2017).

The antioxidant ability of CFHAE might be related to its high phenolic and flavonoid contents. Phytochemical screening in this study showed that CFHAE possesses gallic acid, caffeic acid, chlorogenic acid, rutin, coumaric acid quercetin, cinnamic acid and apigenin which have antioxidant activity.

Besides, carob fruit contains trace elements such as Fe, Cu, Mn and Zn (Özcan *et al.*, 2007; Oziyci *et al.*, 2014) which are

co-factors of antioxidant enzymes (Harris, 1992) which boost antioxidant system in Pb exposure. The MDA levels in serum decreased in Pb + CFHAE groups compared to Pb group which indicates that administration of CFHAE reduces lipid peroxidation. It could be a consequence of the reduction of cell damages due to neutralization of free radicals by CFHAE's antioxidant compounds. Similar to this research, several studies showed that administration of carob aqueous extract decreased MDA levels (Rtibi *et al.*, 2016a; Vafaei *et al.*, 2018).

Data from real time PCR revealed that *Nrf2* expression was down-regulated in Pb + CFHAE groups which demonstrated that co-administration of CFHAE reduced oxidative stress in animals exposed to Pb. This is the first study that demonstrated CFHAE decreased expression of *Nrf2*. Also, co-administration of CFHAE with Pb down-regulated *iNOS* which is not reported so far and indicates that CFHAE decreases the production of NO. In line with the present results, Al-Olayan *et al.* (2016) showed that carob pod aqueous extract reduced NO level in the liver of mice experimentally infected with *Schistosoma mansoni*. Abdel-Rahman *et al.* (2018) showed that treatment of rats exposed to water pipe smoke, with carob aqueous extract decreased NO levels in lung tissue.

Other researchers reported protective potential of carob pods against oxidative stress which is similar to our findings (Abdel-Rahman *et al.*, 2018; Rtibi *et al.*, 2016a, 2016b, 2015; Suzek *et al.*, 2017; Vafaei *et al.*, 2018).

Co-administration of CFHAE elevated sex hormones in groups that were simultaneously exposed to Pb. Increases in LH and FSH levels might be due to protecting effects of CFHAE on endocrine cells of the pituitary gland since Pb attenuates antioxidant enzymes and causes degenerative changes in these cells (Hamadouche *et al.*, 2013). In line with our results, previous researches elucidated that administration of carob extract increases

Carob fruit ameliorates Pb toxicity in reproductive system of male mice

serum concentrations of LH and testosterone (Mokhtari et al., 2012; Vafaei et al., 2018). Oxidative stress reduces antioxidant activity and increases lipid peroxidation in Leydig cells which results in impairment of testosterone synthesis (Glade and Smith, 2015). Furthermore, increases in testosterone levels in Pb + CFHAE groups occurred because CFHAE decreases oxidative stress caused by Pb in Leydig cells.

This study demonstrated that CFHAE alleviates Pb toxicity in the reproductive system by diminishing the development of oxidative stress. Hence, it can be utilized as an antioxidant agent in the diet of infertile male patients and the people who may have chances of occupational and environmental exposure to heavy metals such as Pb and other agents that induce oxidative stress.

Acknowledgment

The authors extend their appreciation to the Faculty of Veterinary Medicine and Urmia University Research Council for funding this study.

Conflict of interest

The authors declare that they have no conflicts of interest.

References

- Abdel-Rahman M, Bauomy AA, Salem FEH, Ahmed Khalifa M. 2018. Carob extract attenuates brain and lung injury in rats exposed to waterpipe smoke. *Egypt J Basic Appl Sci*, 5:31–40.
- Aktan F. 2004. iNOS-mediated nitric oxide production and its regulation. *Life Sci*, 75:639–653.
- Al-Olayan EM, El-Khadragy MF, Alajmi RA, Othman MS, Bauomy AA, Ibrahim SR, Moneim AE. 2016. *Ceratonia siliqua* pod extract ameliorates *Schistosoma mansoni*-induced liver fibrosis and oxidative stress. *BMC Complement Altern Med*, 16: no pagination.
- Anjum MR, Reddy PS. 2015. Recovery of lead-induced suppressed reproduction in male rats by testosterone. *Andrologia*, 47:560–567.
- Ata A, Yildiz-Gulay O, Güngör S, Balic A, Gulay MS. 2018. The effect of carob (*Ceratonia siliqua*) bean extract on Male New Zealand white rabbit semen. *World Rabbit Sci*, 26:209–215.
- Ayinde OC, Ogunnowo S, Ogedegbe RA. 2012. Influence of Vitamin C and Vitamin E on testicular zinc content and testicular toxicity in lead exposed albino rats. *BMC Pharmacol Toxicol*, 13:1–8.
- Bae D-S, Gennings C, Carter Jr WH, Yang RSH, Campaign JA. 2001. Toxicological interactions among arsenic, cadmium, chromium, and lead in human keratinocytes. *Toxicol Sci*, 63:132–142.
- Benzie I, Strain J. 1999. Direct measure of total antioxidant activity of biological fluids and modified version for simultaneous measurement of total antioxidant power and ascorbic. *Methods Enzymol*, 299:15–27.
- Beutler E, Duron O, Kelly BM. 1963. Improved method for the determination of blood glutathione. *J Lab Clin Med*, 61:882–888.
- Choudhary R, Chawala VK, Soni ND, Kumar J, Vyas RK. 2010. Oxidative stress and role of antioxidants in male infertility. *Pakistan J Physiol*, 6:54–59.
- Chowdhury AR. 2009. Recent advances in heavy metals induced effect on male reproductive function-A retrospective. *Al Ameen J Med Sci*, 2:37–42.
- Creasy DM. 2001. Pathogenesis of Male Reproductive Toxicity. *Toxicol Pathol*, 29:64–76.
- Custódio L, Fernandes E, Escapa AL, López-Avilés S, Fajardo A, Aligué R, Alberício F, Romano A. 2009. Antioxidant activity and in vitro inhibition of tumor cell growth by leaf extracts from the carob tree (*Ceratonia siliqua*). *Pharm Biol*, 47:721–728.
- Dakia PA, Wathelet B, Paquot M. 2007. Isolation and chemical evaluation of carob (*Ceratonia siliqua* L.) seed germ. *Food Chem*, 102:1368–1374.
- Dewanto V, Wu X, Adom KK, Liu RH. 2002. Thermal processing enhances the nutritional value of tomatoes by increasing total antioxidant activity thermal processing enhances the nutritional value of tomatoes by increasing total antioxidant activity. *J Agric Food Chem*, 50:3010–3014.

- Dkhil MA, Moneim AEA, Al-Quraishy S. 2016. Indigofera oblongifolia ameliorates lead acetate-induced testicular oxidative damage and apoptosis in a rat model. *Biol Trace Elem Res*, 173:354–361.
- Dorostghoal M, Seyyednejad SM, Jabari A. 2014. Protective effects of *Fumaria parviflora* L. on lead-induced testicular toxicity in male rats. *Andrologia*, 46:437–446.
- Elgawish RAR, Abdelrazek HMA. 2014. Effects of lead acetate on testicular function and caspase-3 expression with respect to the protective effect of cinnamon in albino rats. *Toxicol Reports*, 1:795–801.
- El-Nekeety AA, El-Kady AA, Soliman MS, Hassan NS, and Abdel-Wahhab MA. 2009. Protective effect of *Aquilegia vulgaris* (L.) against lead acetate-induced oxidative stress in rats. *Food Chem Toxicol*, 47:2209–2215.
- Gandhi J, Hernandez RJ, Chen A, Smith NL, Sheynkin YR, Joshi G, Khan SA. 2017. Impaired hypothalamic-pituitary-testicular axis activity, spermatogenesis, and sperm function promote infertility in males with lead poisoning. *Zygote*, 25: 103-110.
- Glade MJ and Smith K. 2015. Oxidative stress, nutritional antioxidants, and testosterone secretion in men. *Ann Nutr Disord Ther*, 2:1019.
- Hamadouche NA, Nesrine S, Abdelkeder A. 2013. Lead toxicity and the hypothalamic-pituitary-testicular axis. *Not Sci Biol*, 5:1–6.
- Hammond PB. 1977. Exposure of humans to lead. *Annu Rev Pharmacol Toxicol*, 17:197–214.
- Harris ED. 1992. Regulation of antioxidant enzymes. *FASEB J*, 6:2675–2683.
- He X, Wu J, Yuan L, Lin F, Yi J, Li J, Yuan H, Shi J, Yuan T, Zhang S, Fan Y. 2017. Lead induces apoptosis in mouse TM3 Leydig cells through the Fas/FasL death receptor pathway. *Environ Toxicol Pharmacol*, 56:99–105.
- He Y, Zou Q, Chen H, Weng S, Luo T. 2016. Lead inhibits human sperm functions by reducing the levels of intracellular calcium, cAMP, and tyrosine phosphorylation. *Tohoku J Exp Med*, 238:295–303.
- Huang BM, Liu MY. 2004. Inhibitory actions of lead on steroidogenesis in MA-10 mouse Leydig tumor cells. *Syst Biol Reprod Med*, 50:5–9.
- Jung K-A, Kwak M-K. 2010. The Nrf2 system as a potential target for the development of indirect antioxidants. *Molecules*, 15:7266–7291.
- Kanter M. 2010. Protective effects of melatonin on testicular torsion/detorsion-induced ischemia-reperfusion injury in rats. *Exp Mol Pathol*, 89:314–320.
- Krassas E, Papadopoulou F, Tziomalos K, Zeginiadou T, Pontikides N. 2008. Hypothyroidism has an adverse effect on human spermatogenesis: a prospective, controlled study. *Thyroid*, 18:1255.
- Kumazawa S, Taniguchi M, Suzuki Y, Shimura M, Kwon MS, Nakayama T. 2002. Antioxidant activity of polyphenols in carob pods. *J Agric Food Chem*, 50:373–377.
- Lindl KA, Jordan-Sciutto KL. 2008. Examining the endogenous antioxidant response through immunofluorescent analysis of Nrf2 in tissue. *Methods Mol Biol*, 477:229–243.
- Liu MY, Lai HY, Yang BC, Tsai ML, Yang HY, Huang BM. 2001. The inhibitory effects of lead on steroidogenesis in MA-10 mouse Leydig tumor cells. *Life Sci*, 68:849–859.
- Loraine JA, Bell ET. 1971. Hormone assays and their clinical application. pp. 44-91 and 541-553, Edinburgh, UK: E. & S. Livingstone.
- Mabrouk A, Ben Cheikh H. 2014. Thymoquinone supplementation ameliorates lead-induced testis function impairment in adult rats. *Toxicol Ind Health*, 32:1114–1121.
- Mishra M, Acharya UR. 2004. Protective action of vitamins on the spermatogenesis in lead-treated Swiss mice. *J Trace Elem Med Biol*, 18:173–178.
- Mokhtari M, Sharifi E, Azadian S. 2012. The effects of hydro alcoholic extract of *Ceratonia siliqua* L. seeds on pituitary - Testis hormones and spermatogenesis in rat. *Adv Environ Biol*, 6:2778–2783.
- Moneim AEA. 2015. The neuroprotective effect of berberine in mercury-induced neurotoxicity in rats. *Metab Brain Dis*, 30:935–942.
- Moniem AEA, Dkhil MA, Al-quraishy S. 2010. Protective role of flaxseed oil against lead acetate induced oxidative stress in testes of adult rats. *Afr J Biotechnol*, 9:7216–7223.

Carob fruit ameliorates Pb toxicity in reproductive system of male mice

- Naghmouchi S, Khouja ML, Romero A, Tous J, Boussaid M. 2009. Tunisian carob (*Ceratonia siliqua* L.) populations: Morphological variability of pods and kernel. *Sci Horti*, 121:125–130.
- Nishikimi M, Rao NA, Yagi K. 1972. The occurrence of superoxide anion in the reaction of reduced phenazine methosulfate and molecular oxygen. *Biochem Biophys Res Commun*, 2:849–854.
- Owen RW, Haubner R, Hull WE, Erben G, Spiegelhalder B, Bartsch H, Haber B. 2003. Isolation and structure elucidation of the major individual polyphenols in carob fibre. *Food Chem Toxicol*, 41:1727–1738.
- Özcan MM, Arslan D, Gökçalik H. 2007. Some compositional properties and mineral contents of carob (*Ceratonia siliqua*) fruit, flour and syrup. *Int J Food Sci Nutr*, 58:652–658.
- Oziyci HR, Tetik N, Turhan I, Yatmaz E, Uçgun K, Akgül H, Gubbuk H, Karhan M. 2014. Mineral composition of pods and seeds of wild and grafted carob (*Ceratonia siliqua* L.) fruits. *Sci Horti*, 167:149–152.
- Paglia DE, Valentine WN. 1967. Studies on the quantitative and qualitative characterization of erythrocyte glutathione peroxidase. *J Lab Clin Med*, 70:158–169.
- Papagiannopoulos M, Wollseifen HR, Mellenthin A, Haber B, Galensa R. 2004. Identification and quantification of polyphenols in carob fruits (*Ceratonia siliqua* L.) and derived products by HPLC-UV-ESI/MSn. *J Agric Food Chem*, 52:3784–3791.
- Patra RC, Rautray AK, Swarup D. 2011. Oxidative stress in lead and cadmium toxicity and its amelioration. *Vet Med Int*, 2011.
- Pedersen H. 2001. Late-flowering dune populations of *Dactylorhiza incarnata* (Orchidaceae): Variation patterns and taxonomic inferences. *Nord J Bot*, 21:177–186.
- Placer ZA, Cushman LL, Johnson BC. 1996. Estimation of product of lipid peroxidation (malonyl dialdehyde) in biochemical systems. *Anal Biochem*, 16:359–364.
- Rahman MA, H. Salem FE, Bauomy AA, Khalifa MA. 2016. Ameliorative effect of carob aqueous extract on water pipe smoke induced-toxicity in adult male albino rats. *Int J Pharm Pharm Sci*, 9:246.
- Reshma Anjum M, Madhu P, Pratap Reddy K, Sreenivasula Reddy P. 2017. The protective effects of zinc in lead-induced testicular and epididymal toxicity in Wistar rats. *Toxicol Ind Health*, 33:265–276.
- Reshma Anjum M, Sainath SB, Suneetha Y, Sreenivasula Reddy P. 2011. Lead acetate induced reproductive and paternal mediated developmental toxicity in rats. *Ecotoxicol Environ Saf*, 74:793–799.
- Rtibi K, Jabri MA, Selmi S, Souli A, Sebai H, El-Benna J, Amri M, Marzouki L. 2015. Gastroprotective effect of carob (*Ceratonia siliqua* L.) against ethanol-induced oxidative stress in rat. *BMC Complement Altern Med*, 15:1–8.
- Rtibi K, Selmi S, Jabri MA, El-Benna J, Amri M, Marzouki L, Sebai H. 2016a. Protective effect of *Ceratonia siliqua* L. against a dextran sulfate sodium-induced alterations in liver and kidney in rat. *J Med Food*, 19:882–889.
- Rtibi K, Selmi S, Jabri MA, Mamadou G, Limas-Nzouzi N, Sebai H, El-Benna J, Marzouki L, Eto B, Amri M. 2016b. Effects of aqueous extracts from: *Ceratonia siliqua* L. pods on small intestinal motility in rats and jejunal permeability in mice. *RSC Adv*, 6:44345–44353.
- Said AA, Kairy MH, Shalaby AME, Ismail HT. 2017. The effect of carob pods and fig fruits ether extracts against lead induced hematological and biochemical changes in *Oreochromis niloticus*. *Zagazig Vet J*, 45:19–28.
- Shotyk W, Le Roux G. 2005. Biogeochemistry and cycling of lead. *Met Ions Biol Syst*, 43:239–275.
- Sigman M. 2007. Medications that impair male fertility. *SRM*, 11–15.
- Simmons SO, Fan C-Y, Yeoman K, Wakefield J, Ramabhadran R. 2011. NRF2 Oxidative Stress Induced by Heavy Metals is Cell Type Dependent. *Curr Chem Genomics*, 5:1–12.
- Sinha AK. 1972. Colorimetric assay of catalase. *Anal Biochem*, 47:389–394.
- Solei manzadeh A, Kian M, Moradi S, Malekifard F. 2018. Protective effects of hydro-alcoholic extract of *Quercus brantii* against lead-induced oxidative stress in the reproductive system of male mice. *Avicenna J Phytomedicine*, 8:448–456.

- Suvarna, KS, Layton C, Bancroft, J.D. eds., 2018. Bancroft's Theory and Practice of Histological Techniques E-Book, pp. 40-139, Elsevier Health Sciences.
- Suzek H, Celik I, Dogan A. 2017. Nephroprotective hepatoprotective potential and antioxidant role of carob pods (*Ceratonia siliqua* L.) against carbon tetrachloride-induced toxicity in rats. *Indian J Pharm Educ Res*, 51:312–320.
- Tejada RI, Mitchell JC, Norman A, Marik JJ, Friedman S. 1984. A test for the practical evaluation of male fertility by acridine orange (AO) fluorescence. *Fertil Steril*, 42:87–91.
- Tetik N, Turhan I, Oziyci HR, Gubbuk H, Karhan M, Ercisli S. 2011. Physical and chemical characterization of *Ceratonia siliqua* L. germplasm in Turkey. *Sci Hortic*, 129:583–589.
- Vafaei A, Mohammadi S, Fazel A, Soukhtanloo M, Pour AM, Beheshti F. 2018. Effects of carob (*Ceratonia siliqua*) on sperm quality, testicular structure, testosterone level and oxidative stress in busulfan-induced infertile mice. *Pharm Sci*, 24:104-111.
- Wang Y, Fang J, Huang S, Chen L, Fan G, Wang C. 2012. The chronic effects of low lead level on the expressions of Nrf2 and Mrp1 of the testes in the rats. *Environ Toxicol Pharmacol*, 35:109–116.
- Wong EWP and Cheng CY. 2011. Impacts of environmental toxicants on male reproductive dysfunction. *Trends Pharmacol Sci*, 32:290–299.
- World Health Organisation. 2010. WHO laboratory manual for the examination and processing of human semen. World Health Organization.
- Yu B, Lin H, Yang L, Chen K, Luo H, Liu J, Gao X, Xia X, Huang Z. 2012. Genetic variation in the Nrf2 promoter associates with defective spermatogenesis in humans. *J Mol Med*, 90:1333–1342.
- Zhang H, Wei K, Zhang M, Liu R, Chen Y. 2014. Assessing the mechanism of DNA damage induced by lead through direct and indirect interactions. *J Photochem Photobiol B Biol*, 136:46–53.