

Original Research Article

Silymarin inhibits Toll-like receptor 8 gene expression and apoptosis in Ramos cancer cell line

Nasrin Ranjbar^{1,3}, Ramin Saravani^{2,3,*}, Zohreh Faezizadeh⁴

¹Student Scientific Research Center, Department of Clinical Biochemistry, School of Medicine, Zahedan University of Medical Sciences, Zahedan, Iran

²Cellular and Molecular Research Center, Department of Clinical Biochemistry, School of Medicine, Zahedan University of Medical Sciences, Zahedan, Iran

³Department of Clinical Biochemistry, School of Medicine, Zahedan University of Medical Sciences, Zahedan, Iran

⁴Department of Laboratory Sciences, Borujerd Branch, Islamic Azad University, Borujerd, Iran

Article history:

Received: Dec 10, 2018

Received in revised form:

May 14, 2019

Accepted: Jul 08, 2019

Vol. 10, No. 2, Mar-Apr 2020,
161-169.

* Corresponding Author:

Tel: +98-543329892

Fax: +98-543329892

saravaniramin@zaums.ac.ir

saravaniramin@yahoo.com

Keywords:

Toll-like receptor

TLR8

Gene expression

Silymarin

Apoptosis

Ramos cells

Abstract

Objective: Silymarin is a herbal extract containing flavonolignans, and it has inhibitory effects against the growth of different cancer cell lines by inducing apoptosis. Toll-like receptors are suggested as a novel and attractive target to treat cancer. The current study aimed at examining the mechanism of silymarin-induced apoptosis in Ramos cells and investigating its effects on *TLR8* expression.

Materials and Methods: The half maximal inhibitory concentration (IC_{50}) of silymarin in Ramos cells was determined via MTT viability test while the type of cell death was tested by annexin V/propidium iodide (PI) double staining method. The activity of caspase-3 and expression of *TLR8* were measured in a time-dependent manner (in IC_{50}) by colorimetric assay and real-time polymerase chain reaction (RT-PCR), respectively.

Results: The results of MTT showed that IC_{50} of silymarin in Ramos cells was 100 $\mu\text{g/ml}$ after 48 hr treatment ($p < 0.01$). Flow cytometry by annexin V/PI, showed that silymarin induced early/late apoptosis in this cell line ($p < 0.05$ to $p < 0.01$). In addition, the caspase-3 colorimetric method showed that caspase-3 increased in the Ramos cell line after treatment ($p < 0.01$). This treatment led to a reduction in *TLR8* mRNA expression in a time-dependent manner ($p < 0.01$).

Conclusion: The results indicated a new mechanism in the anticancer activity of Toll-like receptor (TLR) signaling after silymarin treatment in Ramos cancer cell line. This plant could be used to develop anticancer agents inhibiting TLRs.

Please cite this paper as:

Ranjbar N, Saravani R, Faezizadeh Z. Silymarin inhibits Toll-like receptor 8 gene expression and apoptosis in Ramos cancer cell line. Avicenna J Phytomed, 2020; 10(2): 161-168.

Introduction

Compounds derived from certain plants induce apoptosis in different cancerous cell lines (Unitt and Hornigold, 2011). It was

reported that herbal agents suppress toll-like receptors (TLRs) and may be potential targets of chemotherapeutic compounds. The fruits and seeds of the medicinal plant

Silybum marianum L., are used to obtain a bioactive extract named silymarin, which mainly contains silibinin and partially silibinin stereoisomers (Ramasamy and Agarwal, 2008; Wen et al., 2008). Extensive studies indicated strong chemopreventive and anticancer effects of silymarin on many types of cancers (Wen et al., 2008; Tyagi et al., 2007). There are more than 20 types of lymphatic system cancers. Burkitt lymphoma, a germinal center B-cell-derived cancer, is an important human oncogene found more than three decades ago. This cancer, which is a form of non-Hodgkin's lymphoma, begins in immune cells called B-lymphocytes (Burkitt, 1958). The most tenable hypothesis consistent with all the observed facts is that tumor formation results from interactions between some virus or viruses and a reticuloendothelial system altered by chronic and heavy infection by malarial or other parasites (Burkitt, 1969).

TLRs are transmembrane pattern recognition receptors capable of recognizing generic pathogen-associated molecular patterns (PAMPs) (Barton and Kagan, 2009). Their activation induces intracellular signaling pathways that result in the production of a variety of nuclear factor (NF)- κ B-mediated cytokines (e.g. tumor necrosis factor) and type I interferon (Kawai and Akira, 2011), which prevent cell death by expressing anti-apoptotic proteins such as Bcl-2 and induce chronic inflammation by producing cyclooxygenase-2 (Grimm et al., 2010; Bowie and O'Neill, 2000). Humans have ten known functional TLRs, of which TLR 3, 7, 8, and 9 are expressed on intracellular vesicular membranes and are commonly involved in recognizing nucleic acids; *TLR8* detects viral single-stranded RNA (Kawai and Akira, 2006). Recently, it was observed that TLRs are expressed in tumor cells (Sato et al., 2009). The role of TLRs expression in tumor cells and its relationship with tumor development are exciting subjects in tumor immunity.

TLR8 is activated in different cancers including cervical, pancreatic, and lung

cancer cells (Zhang et al., 2014; Cherfils-Vicini et al., 2010a; Grimmig, et al., 2015). Cherfils-Vicini et al. reported that *TLR8* was expressed in human lung cancer cells, and PolyU (*TLR8* ligand) could upregulate B-cells lymphoma-2 (Bcl-2) and promote the survival of lung cancer cells. They also proposed that Bcl-2 plays a crucial role in autophagy and apoptosis pathways (Cherfils-Vicini et al., 2010b). Therefore, *TLR8* is considered a key marker of tumor genesis and is a highly selective candidate to develop anti-proliferative and anti-malignant agents. To date, there is no report on *TLR8* gene expression and the molecular mechanism involved in the anti-proliferative effect of silymarin in Ramos (P₅₃-mutant) cancer cell line. Also, there are data on mutations in the P₅₃ that associated with malignancy condition such as proliferation, migration, and invasion

The current study aimed at examining the effect of silymarin on *TLR8* gene expression and induction of apoptosis in Ramos cancer cell line.

Materials and Methods

Chemical reagents

All the materials were of analytical grade. RPMI 1640 medium, trypan blue and trypsin were purchased from INOCLON (G. Innovative Biotech Co Inoclon), Iran). Fetal bovine serum (FBS), penicillin/streptomycin and amphotericin B were purchased from Gibco (Rockville, MD, USA). Annexin V/PI (propidium iodide), apoptosis detection kit was obtained from BioVision (San Francisco, CA, USA). 3-(4, 5-dimethylthiazol-2-yl)-2, 5-diphenyltetrazolium bromide (MTT) and dimethyl sulfoxide (DMSO) were purchased from Sigma-Aldrich (St. Louis, MO, USA). Reverse transcription kit and real-time PCR kit were purchased from TaKaRa Bio Inc. (Dalian, China).

Cell culture

The P₅₃-mutant Ramos cells (ATCC® Number: CRL-1596™; Organism: Homo

Silymarin induces apoptosis and reduces *TLR8* expression

sapiens, human; Cell Type: B lymphocyte; and Disease: Burkitt's lymphoma) were maintained in a medium that contained RPMI 1640 medium, fetal bovine serum (10%), penicillin (100 U/ml), and streptomycin (100 µg/ml). The P₅₃-mutant Ramos cell line was grown under cell culture conditions at 37°C, with 95% humidity, with an atmosphere of 5% CO₂.

MTT viability assay

To evaluate the cytotoxic effects of silymarin on Ramos cell line, cells were plated at a density of 6×10^3 cell/well in a 96-well plate and exposed to various concentrations of silymarin (0, 25, 50, 75 and 100 µg/ml) for 24, 48 and 72 hr. Then, the viability of the cells was measured by MTT assay. Briefly, after the aforementioned periods, 20 µl of MTT dye was added and the solution was incubated at 37°C for 3 hr. Then, 200 µl of dimethyl sulfoxide (DMSO) was added and the solution was placed in a dark box for 15 min. Finally, all the plates were read at 570 nm wavelength using a microplate reader (Stat Fax 2100; Awareness Technology, Los Angeles, CA, USA).

Flow cytometry assay using annexin V/PI staining

Apoptosis was detected using annexin/PI staining kit. Ramos cells were seeded (1×10^5 cell/well) in 6-well plates and incubated for 12 hr before treatment with increasing concentrations of silymarin. After 48 hr of treatment, both treated and untreated cells were trypsinized, washed with cold phosphate-buffered saline (PBS), and resuspended in PBS. Apoptosis was measured using an annexin V/PI double staining kit (BioVision, San Francisco, CA, USA). According to the manufacturer's instructions, first of all, the cell pellets were stained with 250 µl 1X binding buffer and later, incubated with 2.5 µl annexin V for 15 min in the dark at room temperature. Before analysis, 2.5 µl of propidium iodide (PI) was added and incubated for 10 min. Finally, the working solution (1000 µl) was added to the samples before analysis by a

fluorescence-activated cell sorting (FACS) cytometer (BD Biosciences, San Jose, CA, USA).

Caspase-3 activity

Caspase-3 activity was measured according to the manufacturer's instructions using colorimetric Assay Kits (R&D systems Co. Germany). Briefly, cells were treated with silymarin (IC₅₀) at different times (0 [control], 6, 12 and 24 hr); next, cells were washed twice in cold PBS and lysed using lysis buffer for 10 min. Then, cells were centrifuged at 10000 g for 10 min and then 50 µl reaction buffer and 5 µl of caspase-3 substrate were added to supernatants and incubated at 37°C for 1 hr. The results of absorbance as measured by a microplate reader (Stat Fax 2100; Awareness Technology, Los Angeles, CA, USA) at 405 nm wavelength are expressed as fold changes of caspase-3 activity in treated cells divided by the caspase-3 activity of adjacent untreated cells.

TLR8 mRNA expression

Cells at a density of 12×10^4 cells/well were seeded in 96-well plates and then they were incubated without or with silymarin (IC₅₀) at different times (0 [control], 4, 8, 12, 24, and 48 hr) and RNAs were extracted by RNX (SinaClon, Tehran, Iran). After synthesis of cDNAs by reverse transcription kit (TAKARA Bio Inc. Dalian, China) based on the extracted RNAs, the mRNA expression levels of *TLR8* gene were evaluated by real-time PCR (Applied Biosystems, Foster City, CA, USA) using specific primers for both target genes [*TLR8* mRNA] and *GAPDH* used as housekeeping gene (Table 1).

Statistical analysis

All data were analyzed by IBM SPSS statistics version 22 (SPSS Inc., Chicago, IL, USA). To compare data, One-way analysis of variance (ANOVA) with the Dunnett's test was used. A p-value of less than 0.05 was considered statistically different.

Table 1. Sequences of the primer pairs for *TLR8* and reference gene.

Target name	Amplicon length	Primer sequence forward	Primer sequence reverse
<i>TLR8</i>	184 bp	5'CAGAGCATCAACCAAAGCAA3'	5'GCTGCCGTAGCCTGAAATA3'
<i>GAPDH</i>	150 bp	5'GAGCCACATCGCTCAGACAC3'	5'CATGTAGTTGAGGTCAATGAAGG3'

Results

Anti-proliferative effect of silymarin on human Ramos cancer cells

The effect of silymarin on the Ramos cell line was examined. The cells in different concentrations of silymarin and at different times were exposed, the cell line viability was examined by using an MTT assay. The most significant inhibitory effects of silymarin in Ramos cancer cell line were $69\pm4.5\%$, $50\pm2.1\%$ and 19 ± 1.5 at concentration $100\ \mu\text{g/ml}$ after treatment for 24, 48 and 72 hr, respectively. Concentration of $100\ \mu\text{g/ml}$ at time 48hr revealed decreased growth by 50% in Ramos cancer cell line ($p<0.01$). Figure 1 shows the results of MTT in cancer cell line.




Figure 1. Effects of silymarin on suppression of Ramos cell growth. Cells were exposed for 24, 48 and 72 hr to different concentrations of silymarin, and cell proliferation was examined by MTT assay. This compound decreased cell proliferation in time- and dose-dependent fashions. $**p<0.01$ show significant differences compared to control cells.

Detection of apoptosis by flow cytometry

To study whether silymarin induced apoptosis in the cell line, both treated and untreated Ramos cells were double-stained

with annexin V-FITC/PI; then, cells were analysed by flow cytometry. The obtained results showed a significant increase in the percentage of both early and late apoptosis in a concentration-dependent manner ($p<0.05$ to $p<0.01$); Figures 2 and 3).




Figure 2. Flow cytometric evaluation of apoptosis in Ramos cells after 48 hr treatment with silymarin by staining with annexin-V and propidium iodide (PI). Both early and late apoptosis were increased in cells in comparison with untreated cell line. $*p<0.05$ and $**p<0.01$ show significant differences compared to control cells.

Caspase-3 activation assay

To examine the role of caspase-3 in silymarin-induced apoptosis, the activity of caspase-3 was measured. Treatment of Ramos cancer cells with $100\ \mu\text{g/ml}$ of silymarin markedly increased the activity of caspase-3 in a time-dependent manner ($p<0.05$ to $p<0.01$). Figure 4 shows Colorimetric assay of caspase-3 activity in cancer cell line.

RT- PCR assay

The expression levels of *TLR8* were examined by real-time RT-PCR using SYBR Green in Ramos cancer cells. As shown in Figure 5, *TLR8* mRNAs in cancer cell line was detected, and also the effects of silymarin on expression of the *TLR8* gene was performed at different times (o[control], 4, 8, 12, 24 and 48 hr). The results show that *TLR8* gene expression level decreased significantly in Ramos cancer cell line in the presence of silymarin ($100\ \mu\text{g/ml}$) in comparison to untreated cells ($p<0.01$).

Silymarin induces apoptosis and reduces *TLR8* expression




Figure 3. Silymarin induced apoptosis in Ramos cell line. Cells were treated with different concentrations of silymarin (0 [control], 25, 50, and 100 µg/ml). After 48 hr of treatment, Ramos cells were stained with annexin/PI. All measurements done by flow-cytometry were performed independently in triplicates per experimental point, and representative results are shown.




Figure 4. Colorimetric assay of caspase-3 activity after treatment with silymarin 100 µg/ml. Activity of caspase-3 was enhanced in Ramos cancer cells. * $p < 0.05$ and ** $p < 0.01$ show significant differences compared to control cells.




Figure 5. Real-time RT-PCR analysis of TLR8 mRNA expression on human Ramos cancer cell line following silymarin (100 µg/ml) treatment in different times. ** $p < 0.01$ show significant differences compared to control cells.

Discussion

The current study focused on the effect of silymarin on *TLR8* expression inhibition, cell viability, and induction of apoptosis in Ramos cancer cell line. The results of the current study showed a significant positive correlation between Ramos cancer cell line with silymarin and reduced expression of *TLR8* and apoptosis in a concentration- and time-dependent fashion; silymarin also increased caspase-3 activity.

Silymarin is composed of several flavonolignans including silybin A, silybin B, isosilybin, and silychristin. The main components of *Silybum marianum* such as polyphenolic fraction and silybin have significant antioxidant and cytoprotective

effects (Křen and Walterova, 2005; Pradhan and Girish, 2006). Previous investigations showed that silymarin induces apoptosis in different cell lines such as HepG2 (Ramakrishnan et al., 2009), PC-3 (Deep et al., 2008), C33A, HeLa (Huang et al., 2005; Yu et al., 2012), and FaDu cell line, as well as oral cancer (Su et al., 2013). So far, extensive studies demonstrated that herbal compounds could inhibit inflammatory responses by the modulation of TLRs (Romano et al., 2013) and finally, induce apoptosis (Marx, 2004; Kelly et al., 2006). TLRs play a pivotal role to maintain tissue homeostasis by regulating tissue repair and inflammatory responses to injuries (Chuang and Ulevitch, 2000).

Through activating the NF- κ B pathway, TLRs play an essential role in inflammatory processes (Grimmig et al., 2015). Thus, targeting TLRs should be considered a potential strategy to treat cancers (Grimmig et al., 2015). *TLR8* gene is mapped to chromosome Xp22 (Wang et al., 2008). Recently, overexpression of *TLR8* was reported in a variety of tumor cells (Pradere et al., 2014; Rivlin et al., 2011) such as pancreatic cancer (Grimm et al., 2010). Previous studies demonstrated that the expression levels of *TLR7/TLR8* genes were associated with colorectal cancer (tumor progression) and that survival rate was lower among patients with *TLR7/TLR8* overexpression (Grimm et al., 2010). Mutations in the *P53* gene are observed in almost 10% of hematopoietic malignancies (Brosh and Rotter, 2009). Also, *P53*-mutant cells show different results on growth inhibition as compared with wild-type *P53* (Farrell et al., 1991). On the other hand, studies revealed that binding to human papillomavirus 16 E6 oncoproteins leads to rapid proteolytic degradation of *P53* and induces the ubiquitin pathway in cells. However, Ramos cells are resistant to E6-directed degradation and are Epstein-Barr virus-negative (EBV-negative). Also, these cells express high levels of *TLR8* and could be used as a suitable model to investigate

Silymarin induces apoptosis and reduces *TLR8* expression

the functions and mechanisms of TLRs in *in vitro* studies (Vousden et al., 1993). However, mutant P₅₃ downstream signaling pathway is not clear because there are different P₅₃-mutant variants (Rivlin et al., 2011). TLRs ligand binding induces myeloid differentiation primary response 88 (MYD88) adapter molecules, which leads to the activation of mitogen-activated protein kinase (MAPK) signaling pathway, and in turn, inhibition of apoptosis via expression of anti-apoptotic proteins such as Bcl-2 (Grimm et al., 2010; Bowie and O'Neill, 2000). In the current study, it was observed that different concentrations (25–100 µg/ml) of silymarin induce apoptosis in Ramos cell line (P₅₃-mutant). A study revealed that silymarin induces apoptosis and causes drug sensitivity in anaplastic lymphoma cells (Molavi et al., 2016). It is also reported that silymarin induces apoptosis in silymarin-treated K562 cells (Faezizadeh et al., 2012).

Caspase-3 plays a vital role to induce apoptosis (Hussein, 2005). The activation of caspase-3 from its zymogen form (procaspase-3) is probably triggered through two mechanisms including the release of cytochrome c (intracellular way) and activation of caspase-8 (extracellular way) (Wang et al., 1999). Induction of apoptosis is considered an anti-cancer mechanism for many bio-active agents. *Epimedium* contains Icariin, the hydrolytic product of which exhibits induction of cell apoptosis by activation of caspases in human Raji Burkitt lymphoma and P3HR-1 cell line (Li et al., 2014). The application of plant compounds, particularly silymarin, as chemotherapeutic compounds for cancers is recommended since they do not impose any adverse effects on normal lymphocytes (Agarwal et al., 2006; Shanmugam et al., 2011). Therefore, targeting TLRs could be a promising strategy to expand anticancer agents.

In summary, the current study data indicated downregulation of *TLR8* expression after silymarin treatment in Ramos cancer cells. It was also discovered

that silymarin effectively suppressed cell proliferation, induced apoptosis in Ramos cancer cells in a concentration-dependent fashion, and caused activation of caspase-3. Ultimately, according to the obtained results, *TLR8* signaling may be a potential target for Burkitt lymphoma therapy.

Acknowledgment

This paper was based on the MSc. Thesis of Nasrin Ranjbar (grant No. K/92); Zahedan University of Medical Sciences provided the fund.

Conflicts of interest

The authors have declared that there is no conflict of interest.

References

- Agarwal R, Agarwal C, Ichikawa H, Singh RP, Aggarwal BB. 2006. Anticancer potential of silymarin: from bench to bed side. *Anticancer Res*, 26: 4457-4498.
- Barton GM, Kagan JC. 2009. A cell biological view of Toll-like receptor function: regulation through compartmentalization. *Nat. Rev. Immunol*, 9:535-542.
- Bowie A, O'Neill L. 2000. The interleukin-1 receptor/Toll-like receptor superfamily: signal generators for pro-inflammatory interleukins and microbial products. *J Leukoc Biol*, 67: 508-514.
- Brosh R, Rotter V. 2009. When mutants gain new powers: news from the mutant p53 field. *Nat Rev Cancer*, 9: 701.
- Burkitt D. 1958. A sarcoma involving the jaws in African children. *Br J Surg*, 46:218-23.
- Burkitt DP. 1969. Etiology of Burkitt's lymphoma—an alternative hypothesis to a vectored virus. *J Natl Cancer Inst*, 42: 19-28.
- Cherfils-vicini J, Platonova S, Gillard M, Laurans L, Validire P, Caliandro R, Magdeleinat P, Mami-chouaib F, Dieu-nosjean MC, Fridman WH. 2010a. Triggering of *TLR7* and *TLR8* expressed by human lung cancer cells induces cell survival and chemoresistance. *J Clin Invest*, 120:1285.
- Cherfils-vicini J, Platonova S, Gillard M, Laurans L, Validire P, Caliandro R, Magdeleinat P, Mami-chouaib F, Dieu-nosjean MC, Fridman WH. 2010b.

- Triggering of TLR7 and TLR8 expressed by human lung cancer cells induces cell survival and chemoresistance. *J Clin Invest*, 120:1285-1297.
- Chuang TH, Ulevitch RJ. 2000. Cloning and characterization of a sub-family of human toll-like receptors: hTLR7, hTLR8 and hTLR9. *Eur Cytokine Netw*, 11: 37.
- Deep G, Oberlies NH, Kroll DJ, Agarwal R. 2008. Identifying the differential effects of silymarin constituents on cell growth and cell cycle regulatory molecules in human prostate cancer cells. *Int J Cancer*, 123: 41-50.
- Faezizadeh Z, Mesbah-namin SAR, Allameh A. 2012. The effect of silymarin on telomerase activity in the human leukemia cell line K562. *Planta Med*, 78, 899-902.
- Farrell PJ, Allan GJ, Shanahan F, Vousden K, Crook T. 1991. p53 is frequently mutated in Burkitt's lymphoma cell lines. *EMBOJ*, 10: 2879-2887.
- Grimm M, Kim M, Rosenwald A, Heemann U, Germer CT, Waaga-gasser AM, Gasser M. 2010. Toll-like receptor (TLR) 7 and TLR8 expression on CD133+ cells in colorectal cancer points to a specific role for inflammation-induced TLRs in tumourigenesis and tumour progression. *Eur J Cancer*, 46: 2849-2857.
- Grimmig T, Matthes N, Hoeland K, Tripathi S, Chandraker A, Grimm M, Moench R, Moll EM, Friess H, Tsaur I. 2015. TLR7 and TLR8 expression increases tumor cell proliferation and promotes chemoresistance in human pancreatic cancer. *Int J Oncol*, 47: 857-866.
- Huang Q, Wul J, Tashiro SI, Onodera S, Li LH, Ikejima T. 2005. Silymarin augments human cervical cancer HeLa cell apoptosis via P38/JNK MAPK pathways in serum-free medium. *J Asian Nat Prod RES*, 7: 701-709.
- Hussein MR. 2005. Apoptosis in the ovary: molecular mechanisms. *Hum Reprod Update*, 11: 162-178.
- Kawai T, Akira S. 2006. TLR signaling. *Cell Death Differ*, 13: 816-825.
- Kawai T, Akira S. 2011. Toll-like receptors and their crosstalk with other innate receptors in infection and immunity. *Immunity*, 34: 637-650.
- Kellym G, Alvero AB, Chen R, Silasi DA, Abrahams VM, Chan S, Visintin I, Rutherford T, Mor G. 2006. TLR-4 signaling promotes tumor growth and paclitaxel chemoresistance in ovarian cancer. *Cancer Res*, 66: 3859-3868.
- Kren V, Walterova D. 2005. Silybin and silymarin-new effects and applications. *Biomedical Pap*, 149:29-41.
- Li ZJ, Yao C, Liu SF, Chen L, Xi YM, Zhang W, Zhang GS. 2014. Cytotoxic effect of icaritin and its mechanisms in inducing apoptosis in human burkitt lymphoma cell line. *Biomed Res Int*, 2014.
- Marx J. 2004. Inflammation and cancer: the link grows stronger. *Science*, 306: 966.
- Molavi O, Samadi N, Wu C, Lavasanifar A, Lai R. 2016. Silibinin suppresses NPM-ALK, potently induces apoptosis and enhances chemosensitivity in ALK-positive anaplastic large cell lymphoma. *Leuk Lymphoma*, 57: 1154-1162.
- Pradere JP, Dapito DH, Schwabe RF. 2014. The Yin and Yang of Toll-like receptors in cancer. *Oncogene*, 33: 3485.
- Pradhan S, Girish C. 2006. Hepatoprotective herbal drug, silymarin from experimental pharmacology to clinical medicine. *Indian J Med Res*, 124: 491.
- Ramakrishnan G, Lo Muzio L, Elinos-baez CM, Jagan S, Augustine TA, Kamaraj S, Anandakumar P, Devaki T. 2009. Silymarin inhibited proliferation and induced apoptosis in hepatic cancer cells. *Cell Proliferation*, 42: 229-240.
- Ramasamy K, Agarwal R. 2008. Multitargeted therapy of cancer by silymarin. *Cancer Lett*, 269, 352-362.
- Rivlin N, Brosh R, Oren M, Rotter V. 2011. Mutations in the p53 tumor suppressor gene: important milestones at the various steps of tumorigenesis. *Genes Cancer*, 2: 466-474.
- Romano B, Pagano E, Montanaro V, Fortunato AL, Milic N, Borrelli F. 2013. Novel insights into the pharmacology of flavonoids. *Phytother Res*, 27: 1588-1596.
- Sato Y, Goto Y, Narita N, Hoon DS. 2009. Cancer cells expressing toll-like receptors and the tumor microenvironment. *Cancer Microenviron*, 2: 205-214.
- Shanmugam MK, Kannaiyan R, Sethi G. 2011. Targeting cell signaling and apoptotic pathways by dietary agents: role in the prevention and treatment of cancer. *Nutr Cancer*, 63: 161-173.
- Su CH, Chen LJ, Liao J F, Cheng JT. 2013. Increase of phosphatase and tensin homolog by silymarin to inhibit human pharynx

Silymarin induces apoptosis and reduces *TLR8* expression

- squamous cancer. *J Med Food*, 16:778-784.
- Tyagi A, Raina K, Singh RP, Gu M, Agarwal C, Harrison G, Glode LM, Agarwal R. 2007. Chemopreventive effects of silymarin and silibinin on N-butyl-N-(4-hydroxybutyl) nitrosamine-induced urinary bladder carcinogenesis in male ICR mice. *Mol Cancer Ther*, 6: 3248-3255.
- Unitt J, Hornigold D. 2011. Plant lectins are novel Toll-like receptor agonists. *Biochem Pharmacol*, 81: 1324-1328.
- Vousden KH, Crook T, Farrell PJ. 1993. Biological activities of p53 mutants in Burkitt's lymphoma cells. *J Gen Virol*, 74: 803-810.
- Wang CY, Guttridge DC, Mayo MW, Baldwin S. 1999. NF- κ B induces expression of the Bcl-2 homologue A1/Bfl-1 to preferentially suppress chemotherapy-induced apoptosis. *Mol Cell Biol*, 19: 5923-5929.
- Wang R, Miyahara Y, Wang H. 2008. Toll-like receptors and immune regulation: implications for cancer therapy. *Oncogene*, 27: 181.
- Wen Z, Dumas TE, Schrieber SJ, Hawke RL, Fried MW, Smith PC. 2008. Pharmacokinetics and metabolic profile of free, conjugated, and total silymarin flavonolignans in human plasma after oral administration of milk thistle extract. *Drug Metab Dispos*, 36: 65-72.
- Yu HC, Chen LJ, Cheng KC, Li YX, Yeh CH, Cheng JT. 2012. Silymarin inhibits cervical cancer cell through an increase of phosphatase and tensin homolog. *Phytother Res*, 26: 709-715.
- Zhang Y, Yang H, Barnie PA, Yang P, Su Z, Chen J, Jiao Z, Lu L, Wang S, Xu H. 2014. The expression of Toll-like receptor 8 and its relationship with VEGF and Bcl-2 in cervical cancer. *Int J Med Sci*, 11: 608.