

Review Article

A review study on the effect of Iranian herbal medicines against *in vitro* replication of herpes simplex virus

Mohammad-Taghi Moradi¹, Mahmoud Rafieian-Kopaei², Ali Karimi^{2*}

¹Students Research Committee, Shahrekord University of Medical Science, Shahrekord, Iran

²Medical Plants Research Center, Shahrekord University of Medical Science, Shahrekord, Iran

Article history:

Received: Jul 01, 2015

Received in revised form:

Sep 20, 2015

Accepted: Nov 06, 2015

Vol. 6, No. 5, Sep-Oct 2016,
506-515.

*** Corresponding Author:**

Tel: +98383 3346692

Fax: +98383330709

rakarimi1342@gmail.com

Keywords:

Herpes Simplex Virus

Herbal medicine

Iran

In-vitro

Abstract

Objective: There are a number of published data indicating *in vitro* anti-HSV activity of some of Iranian herbal extracts with no systematic review to discuss these results. Therefore, this article was aimed to review and discuss the methods carried out and the phytochemistry and bioactivity of the extracts used and also conclusions provided in these publications.

Materials and Methods: Published articles both in English (from Medline, Science Direct, EMBASE, Scopus, Pro Quest, Google scholar, Cochrane Library) and in Persian (from SID, Iran Medex and Magiran) databases, from 1966 to October 2014 were incorporated in this review. The *in vitro* studies that lacked CC₅₀, IC₅₀, were excluded.

Results: Only 42 published reports were found to examine Iranian herbs against HSV replication *in vitro*. Seventeen out of 42 studies in which 23 kinds of medicinal plants were subjected to crude extraction were included. The review of data showed that some of the herbal extracts including *Hyssopus officinalis* methanolic extract, *Melissa officinalis* aqueous extract, *Quercus persica* L. hydroalcoholic extract and *Securigeras ecuridaca* methanolic extract with selective index (SI) of 234, 877, >778 and 250, respectively were highly effective against HSV *in vitro*.

Conclusion: More comprehensive studies using more advanced methods are needed to be done to achieve promising anti-HSV agents from the bioactive compounds isolated from these herbs.

Please cite this paper as:

Moradi MT, Rafieian-Kopaei M, Karimi A. A review study on the effect of Iranian herbal medicines against *in vitro* replication of herpes simplex virus. Avicenna J Phytomed, 2016; 6 (5): 506-515.

Introduction

Belonging to the family of Herpesviridae and subfamily of Alphaherpesvirinae, herpes simplex virus type-1 (HSV-1) and type 2 (HSV-2) are two of the most common human pathogens. Infections caused by these

viruses are public health concern worldwide. These infections range from inapparent to severe life threatening infections such as encephalitis (Koelle and Corey, 2008). Although HSV-1 and HSV-2 are usually transmitted via different routes and infections caused by these two

viruses involve different areas of the body, there is a great deal of overlapping features between the epidemiology and clinical manifestations of their infections. It is generally accepted that HSV is involved in oral diseases as HSV-1 is associated with recurrent aphthous ulceration (Malvy et al., 2007). HSV-1 infection commonly occurs in the mouth and lips resulting in gingivostomatitis (Malvy et al., 2007).

During the past two decades, the mechanisms of replication and pathogenesis of HSV-1 and thus the potential antiviral targets in this virus have been widely understood leading to development of antiviral compounds that target these mechanisms (Koelle and Corey, 2008). The antiviral drugs that are most commonly used to treat HSV-1 infections are nucleoside analogues such as Acyclovir (ACV) and Penciclovir of which Acyclovir is widely used as a drug of choice (Villarreal, 2003). However, a major problem associated with the use of ACV, is the development of drug-resistant HSV strains, particularly in AIDS patients (Elion, 1993). This kind of drug-resistance may occur after long-term treatment and is mainly due to mutations in HSV-1 thymidine kinase and/or DNA polymerase genes (Hill et al., 2009).

As potential alternatives to these agents, antiviral agents from medicinal plants with effective compounds exhibiting different modes of action against HSV infections are urgently needed. These natural active compounds are potential source of novel antiviral agents that could be widely used against these viruses. Some of these compounds have been identified to be active against HSV infections. They have high chemical diversity with biochemical specificity and have biologically active small molecules with drug-like properties that can be absorbed and metabolized in the body (Kitazato et al., 2007). Furthermore, they are easily available and no laborious pharmaceutical work is needed for their synthesis. Therefore, these natural products may be good candidates

for development of special anti-HSV agents (Girish and Pradhan, 2008; Li et al., 2009).

To the best of our knowledge, so far, no systematic review has been published regarding Iranian medicinal plants. Therefore, this article was aimed to review and summarize published results reporting Iranian medicinal plants with *in vitro* activity against HSV replication. This is to provide some recommendations for future research.

Materials and Methods

To provide this systematic review, a number of databases including Medline, Science Direct, EMBASE, Scopus, Pro Quest, Google scholar, Cochrane Library Database and Persian databases such as SID, Iran Medex and Magiran, from 1966 to October 2014 were searched. The terms Herpes simplex virus, anti-viral, plant (herb), Iran, and natural or herbal medicine, were used as keywords. The databases used were limited to those Iranian articles published both in English and in Persian. In this review, *in-vitro* studies lacking CC₅₀, IC₅₀, were excluded. All selected articles were studied by two reviewers to examine inclusion criteria and including common and scientific names of herbs, and their outcome.

Results

Using on line database search, only 42 published reports that examined Iranian herbs for *in vitro* inhibition of HSV replication were found. In this systematic review, 17 studies in which 23 kinds of medicinal plants were subjected to crude extraction were included and the remaining that used no standard methods to evaluate CC₅₀, IC₅₀, and SI, were excluded. The review of data showed that some of the herbal extracts including *Hyssopus officinalis* methanolic extract, *Melissa officinalis* aqueous extract,

Quercus persica L. hydro alcoholic extract and *Securigeras ecuridaca* methanolic extract with SI of 234, 877, >778 and 250, respectively were highly effective against HSV, *in vitro* (Astani et al., 2014;

Behbahani, 2009; Behbahani et al., 2013a; Karimi et al., 2013). Data of these published studies are summarized in Table 1.

Table 1. *In vitro* anti-HSV characteristics of some of Iranian herbal medicines

No	Botanical name	Extract	Common name	Parts	Family	Type of virus	IC ₅₀ (µg/ml)	CC ₅₀ (µg/ml)	SI	references
1.	<i>Aloe vera</i>	hot glycerine extract	Sabrezard	Leaf	Xanthorrhoeaceae	HSV-2	536	3238	7.56	(Zandi et al., 2007)
2.	<i>Avicenna marina</i>	methanol extract	Harra	Leaf	Avicenniaceae	HSV-2	10.5	262.5	25	(Behbahani et al., 2013b)
3.	<i>Avicenna marina</i>	Methanolic extract	Harra	Leaf	Avicenniaceae	HSV-1	80	>200	8.67	(Namazi et al., 2013)
4.	<i>Avicenna marina</i>	Ethanol extract	Harra	Leaf	Avicenniaceae	HSV-1	192	200	0.17	(Namazi et al., 2013)
5.	<i>Avicenna marina</i>	Water extract	Harra	Leaf	Avicenniaceae	HSV-1	>200	>200	1.21	(Namazi et al., 2013)
6.	<i>Avicenna marina</i>	Chloroformic extract	Harra	Leaf	Avicenniaceae	HSV-1	180	110	0.001	(Namazi et al., 2013)
7.	<i>Avicenna marina</i>	n-hexanic extract	Harra	Leaf	Avicenniaceae	HSV-1	176	35	0.007	(Namazi et al., 2013)
8.	<i>Avicenna marina</i>	crude hot glycerine extract	Harra	Leaf	Avicenniaceae	HSV-1	137	5751	41.9	(Zandi et al., 2009)
9.	<i>Cuminumcyminum</i>	Methanolic extract	Zirehsabz	Seed	Apiaceae.	HSV-1	180	450	2.5	(Motamedifar et al., 2010)
10.	<i>Thymus kotschyanus</i>	aqueous extracts	Avishan	Flower + leaf	Lamiaceae	HSV-1	100	800	8	(Farahani, 2013)
11.	<i>Echinacea purpurea</i>	aqueous extracts	Sarkharghol	Root	Asteraceae or Compositae	HSV-1	500	900	1.8	(Farahani, 2013)
12.	<i>Camellia sinensis</i>	aqueous extract	Chai	Leaf	Theaceae	HSV-1	50	1000	20	9Farahani, 2013)
13.	<i>Echiumamoenum</i> L.	aqueous extract	Goleghavzaban	Flower	Boraginaceae	HSV-1	500	1000	2	(Farahani, 2013)
14.	<i>Hyssopus officinalis</i>	Methanol96% extract	Zofa	Leaf	Lamiaceae	HSV-1	4.1	960	234	(Behbahani, 2009)
15.	<i>Melissa officinalis</i>	aqueous extract	Badrangboyeh	Leaf	Lamiaceae	HSV-1	0.4	350	875	(Astani et al., 2014)
16.	<i>Euphorbia spinidens</i>	methanolic extract	Shirmal	aerial parts	Euphorbiaceae	HSV-1	5072	320	15.85	(Mohammadi-Kamalabadi et al., 2014)
17.	<i>Eucalyptus amigdalina</i>	hydroalcoholic extract	okaliptus	Leaf	Myrtaceae	HSV-1	180	650	3.6	(Karimi, 2012)
18.	<i>Quercuspersica</i>	hydroalcoholic extract	Baloot	Seed	Fagaceae	HSV-1	257	>200000	>778	(Karimi et al., 2013)
19.	<i>Myrtuscommunis</i>	hydroalcoholic extract	Moord	Leaf	Myrtaceae	HSV-1	3100	4960	1.6	(Moradi et al., 2011)
20.	<i>Curcuma longa</i>	-	Zardchbeh	Root	Zingiberaceae	HSV-1	33.0	484.2	14.6	(Zandi et al., 2010)
21.	<i>oleaeuropaea</i>	hydroalcoholic extract	zeyton	Leaf	Oleaceae	HSV-1	660	1750	2.6	(Motamedifar et al., 2007)
22.	<i>Securigeras ecuridaca</i>	Methanolic extract	Adasallmolok	Seed	Leguminosae	HSV-2	1.6	130	81.2	(Sayedipour et al., 2012)
23.	<i>Securigeras ecuridaca</i>	Methanolic extract	Adasallmolok	Seed	Leguminosae	HSV-1	2	500	250	(Behbahani et al., 2013a)
24.	<i>Glycyrrhiza glabra</i>	-	Shirinbayan	Root	Leguminosae	HSV-1	500	800	1.6	(Monavari et al., 2008)
25.	<i>Zataria Multiflora</i> Boiss	essential oil	Avishansirazi	Leaf	Lamiaceae	HSV-1	0.0059%	0.067%	11.7	(Mardani et al., 2012)

IC₅₀: 50% inhibition of viral cytopathic, CC₅₀: 50% cytotoxic concentration, SI: Selectivity index (CC₅₀/IC₅₀), HSV-1: herpes simplex virus type 1, HSV-2: herpes simplex virus type 2

Since each extract with $SI \geq 4$ should be considered as a potential antiviral agent (Tsuchiya et al., 1985), the characteristics of some of the extracts isolated from some Iranian medicinal plants are discussed below.

Avicenna marina

Avicenna marina (commonly known as grey mangrove or white mangrove), is one of the mangrove species plants which is native to Southern Africa (Duke, 1991). This plant grows in Hara forests of Persian Gulf and is used in Persian folk medicine. Leaf extracts of *A. marina* contain compounds such as iridoid glucosides, fatty acids, sterols and hydrocarbons (Hogg and Gillan, 1984; König and Rimpler, 1985).

Based on the results of a study, the methanol extract of this herbal medicine showed higher antiviral activity against HSV, *in vitro*. The most polar fraction (D) of the methanol extract showed significant anti-HSV activity. This fraction inhibited the HSV replication following entry of the virus into the target cells. The phytochemical examinations of this extract have shown that fraction D contains flavonoid compounds. It has been shown that *A. marina* contains active flavonoid compounds which inhibit the replication of HSV (KOS strain) following penetrating into the target cells (Namazi et al., 2013).

Behbahani et al showed that crude methanol extract of *A. marina* and the corresponding fractions LMEG and luteolin inhibit HSV-2 replication with selective index (SI) of 25, 25 and 5.3, respectively. The results indicated that LMEG was the most active sub-fraction of *A. marina* with inhibitory effect on the early stage of HSV-2 replication (Behbahani et al., 2013b). Zandi et al showed that crude hot glycerin extract of fresh leaves of *A. marina* has antiviral activity against HSV-1 replication, *in vitro*, mostly before adsorption of the virus to the cells (Zandi et al., 2009).

Aloe vera

Aloe vera is a member of the Liliaceae family. Historically, it has been used for medical purposes such as treatment of wounds and burns (Perfect et al., 2005). This medicinal plant is the source of two herbal preparations namely, aloe gel and aloe latex. Aloe gel is often referred to a clear gel or mucilaginous substance produced by parenchymal cells located in the central region of its leaves. Diluted aloe gel is commonly referred to as *A. vera* extract. The main components of this gel are water (99%), monosaccharides, and polysaccharides (25% of the dry weight of the gel) (Shelton, 1991). There are some published reports indicating that the extracts of *A. vera* and particularly its anthraquinone compounds have antiviral activity (Andersen et al., 1991; Sydiskis et al., 1991). It has been shown that crude hot glycerine extract of *Aloe vera* gel, grown in southwest of Iran, Bushehr, inhibits HSV-2 replication, both before and after attachment to the Vero cell line (Zandi et al., 2007).

Hyssopus officinalis

Hyssopus officinalis is a member of the Lamiaceae family. Some members of this family such as *Origanum vulgare* are known to be poisonous and traditionally used as an arrow poison. Growing in different geographical regions of Iran, *H. officinalis*, mainly its leaves, has been used in the traditional Iranian medicine (TIM) for treatment of various ailments (Kazazi et al., 2007). Methanolic extract of the leaves of *H. officinalis*, has exhibited remarkable activity against both wild type and resistant strains of HSV, *in vitro*. The extract, also, significantly inhibited plaque formation of both of the two strains of HSV-1 in Vero E6 cells with minimal cell cytotoxicity ($CC_{50} = 960 \mu\text{g/ml}$). An oral dose of the extract (125 mg/kg) significantly delayed the onset of HSV-1 infection by over 50%. It also increased the mean survival time of infected mice by 55-65% compared to the non-treated mice

($p < 0.05$). The mortality rate for mice treated with the extract was also significantly reduced by 90% in comparison with the non-treated mice that exhibited 100% mortality (Behbahani, 2009).

Melissa officinalis

Leaves of *Melissa officinalis* (lemon balm), a member of the Lamiaceae family (Saller et al., 2001; Wolbling and Leonhardt, 1994) contain polyphenols such as rosmarinic acid and flavonoids (Carnat et al., 1998). *M. officinalis* has been evaluated for its activity against HSV, *in vitro* (Geuenich et al., 2008; Nolkemper et al., 2006). Aqueous extract of *M. officinalis* has been shown to interact directly with free viral particles of two acyclovir-resistant and one acyclovir-sensitive strains of HSV-1 at low IC_{50} values of 0.13, 0.23 and 0.4 $\mu\text{g/ml}$ and high selectivity indices of 2692, 1522 and 875, respectively with inhibiting attachment of these strains to the host cells in a dose-dependent manner. The results of this work indicated that anti-HSV-1 activity of Melissa extract was mainly due to its rosmarinic acid contents (Astani et al., 2014).

Zataria multiflora

Zataria multiflora Boiss (Avishan-e-Shirazi in Persian and Sa'atar or Zaatar in ancient Iranian medical books) is a thyme-like plant and a member of Lamiaceae family that wildly grows in central and southern parts of Iran (Amin, 1991). It is used in traditional folk remedies for its antiseptic, analgesic (pain-relieving) and carminative (anti-flatulence and intestine soothing) properties (Amin, 1991; Mozaffarian, 1996). In addition to antibacterial activity (Eftekhar et al., 2011), essential oil of *Zataria multiflora* Boiss has been shown to have significant inhibitory effect on HSV-1 replication with IC_{50} values and selectivity indices of 0.0059% and 11.7, respectively (Mardani et al., 2012).

Quercus persica

Q. persica "oak" is a tree of which about 600 species exist around the world. *Quercus* species contain high levels of tannins in both hydrolysable and condensed (Rakić et al., 2007) forms. Oak fruit (acorn) comprises starch, protein, oil, and tannins with high amounts of amylose, amylopectin, high molecular weight and viscosed substances (Saffarzadeh et al., 1999). This fruit is useful in treatment of anemia, diarrhea and is used in livestock feeding. There are some published reports indicating antiviral (Karimi et al., 2013; Muliawan et al., 2006), antibacterial (Khosravi and Behzadi, 2006), and anti-inflammatory (Kaur et al., 2004) potentials of *Quercus* species. These effects have been attributed to its phenolic contents, particularly to its flavonoid and tannin compounds (Chang et al., 2002). It has been suggested that tannin-based components of oak fruits have remarkable effect on virus replication with different mechanisms and with the minimum cytotoxicity (Buzzini et al., 2008). In our previous study, *Q. persica* L. had highly anti-HSV-1 activity and very low cytotoxicity *in vitro*, as exhibited no cytotoxic effect on this cell line up to concentration of 200 mg/ml. Using inhibition assay, this extract inhibited HSV-1 replication by reduction of its cytopathic effect (CPE) before and after cellular attachment in concentrations of 1.02 and 0.257 mg/ml, respectively (Karimi et al., 2013).

Thymus kotschyanus

Lamiaceae (formerly Labiatae) is one of the most important plant families in which *Thymus* with about 215 species, is a significant genus (Zaidi and Crow, 2005). The plants of this genus are well known as medicinal plants because of their biological and pharmacological properties. Fourteen species of this genus are represented in Iranian flora, of which *T. daenensis* Celak, *T. kotschyanus* Boiss, and *T. Hohen* are widely used as medicinal

plants. Infusion and decoction of aerial parts of these species are used in Iranian traditional medicine (ITM) as tonic, carminative, digestive, antispasmodic, anti-inflammatory, anti-tussive and expectorant and for the treatment of common colds (Zargari, 1990). Aqueous extract of *T. kotschyanus* has been shown to inhibit HSV-1 *in vitro* with IC₅₀ value and selectivity indices of 100 µg/ml and 8, respectively (Farahani, 2013).

Securigeras ecuridaca

Securigeras ecuridaca (family Fabaceae), also called goat pea, is an annual herb used against a variety of diseases such as malaria, gastric reflux, epilepsy, hypertension and hyperlipidemia in eastern folk medicine (Garjani et al., 2009; Pouramir et al., 2011). Compounds such as cardenolides, steroidal and pentacyclic have been detected in extract of *S. securidaca* seeds (Zatula et al., 1969). Kaempferol and kaempferol-7-O-glucoside have also been isolated from *S. securidaca* with potential activity against HSV-1 and HSV-2 replication (Behbahani et al., 2013a; Sayedipour et al., 2012). Behbahani et al, evaluated anti HSV-1 activity of 20 µg/ml of both crude methanol extract and twelve associated fractions of *S. securidaca*. Their results showed that IC₅₀ values of crude methanol extract, kaempferol, kaempferol-7-O glycoside and Acyclovire (ACV) were 2±0.12, 0.2±0.01, 0.1±0.01 and 0.1±0.01 µg/ml, respectively. Also, the calculated selective index (SI) for crude methanol extract, kaempferol and kaempferol-7-O-glycoside were 250, 300 and 2500, respectively (Behbahani et al., 2013a). It has also been shown that crude methanol extract of *S. securidaca* had inhibitory effect on HSV-2 in cell culture with IC₅₀ values and selectivity indices of 1.6 µg/ml and 81.2, respectively (Sayedipour et al., 2012).

Curcuma longa

Curcuma longa L. has been used in Asian traditional medicine (ATM) for a

long time to treat gastrointestinal upset, arthritic pain, urinary tract infection, and liver ailments (Dixit et al., 1988; Luper, 1999). One of its main component, Thcurcumin has been shown to have several pharmacological activities such as anti-inflammatory, anti-oxidant, anti-microbial properties (Maheshwari et al., 2006), and antiviral activity against hepatitis B virus (HBV) by decreasing the transcription of HBV X (HBx) gene (Kim et al., 2009). Antiviral activity of different concentrations of curcumin, gallium-curcumin and Cu-curcumin were tested against HSV-1 and the results showed that curcumin and its new derivatives had remarkable antiviral effects on replication of this virus in cell culture. The IC₅₀ values of curcumin, gallium-curcumin and Cu-curcumin were 33.0, 13.9 and 23.1 µg/ml, respectively and their calculated SI values were 14.6, 18.4 and 14.1, respectively (Zandi et al., 2010).

Camellia sinensis

Tea is a cultivated evergreen plant and its green, oolong and black types are all made from the same plant species, *Camellia sinensis* L. Their differences in appearance, organoleptic taste, and chemical content as well as flavour are due to their fermentation process. The chemical components of its leaves include polyphenols (catechins and flavonoides), alkaloids (caffeine, theobromine, the ophylline, etc.), volatile oils, polysaccharides, amino acids, lipids, vitamins (e.g., vitamin C), inorganic elements (e.g., aluminium, fluorine and manganese), etc. The beneficial properties of tea are attributed to its polyphenols. Its flavonoides have antioxidant, anti-inflammatory, anti-allergic and anti-microbial effects. The polyphenols content of green tea and black tea varies from 30% to 40% and 3% to 10%, respectively. Tea and particularly, green tea has been evaluated for its medicinal properties and a number of studies suggested that chemical constituents of tea play important role in

human health (Sharangi, 2009). It was shown that *in vitro* replication of HSV-1 is inhibited by aqueous extract of *C. sinensis* with IC₅₀ value and selectivity index of 50 and 20 µg/mL, respectively (Farahani, 2013).

Euphorbia spinidens

Plants in the family of Euphorbiaceae are well known for the chemical diversity of their isoprenoid constituents (Jury et al., 1987). Members of the *Euphorbia* genus are known to contain substances with inhibitory action against some bacteria. This inhibitory action has been attributed to their large amounts of phenolic compounds (Li et al., 2009). *E. spinidens* is an indigenous plant naturally growing in different geographical regions of Iran and is increasingly used in traditional medicine. Crude methanol extract of *E. spinidens* and the corresponding fractions n-Hexane, Chloroform, and n-Butanol, with SI values of 15.85, 2.062, 8.2, and 25.37, respectively, inhibited early stages of HSV-1 replication, *in vitro* (Mohammadi-Kamalabadi et al., 2014).

Conclusion

Searching for new bioactive compounds in order to develop new anti-viral agents seems to be urgent. Research on some medicinal plants may potentially lead to isolation of active compounds which could be effective against HSV. To achieve this goal, both experimental and clinical investigations are needed to provide a better understanding of mechanisms of action, therapeutic effects, and the safety profile of these compounds. Iran with a diverse climate and wide variety of herbal flora could be considered as potentially a great source of these herbs from which isolation of these components is possible. Therefore, this systematic review which covered the published results of research on some Iranian medicinal plants and the corresponding isolated active phytochemicals, conclusively, would highlight the need for more comprehensive

studies using more advanced methodology to develop promising anti-HSV agents from these bioactive compounds.

Acknowledgements

This research was supported by Research and Technology Deputy of Shahrekord University of Medical Sciences.

Conflict of interest

The authors declare no conflict of interests.

References

- Amin GR. 1991. Popular medicinal plants of Iran. Tehran: Iranian Research Institute of Medicinal Plants, pp. 40.
- Andersen DO, Weber ND, Wood SG, Hughes BG, Murray BK, North JA. 1991. In vitro virucidal activity of selected anthraquinones and anthraquinone derivatives. *Antiviral Res*, 16:185-196.
- Astani A, Navid MH, Schnitzler P. 2014. Attachment and penetration of acyclovir-resistant herpes simplex virus are inhibited by *Melissa officinalis* extract. *Phytother Res*, 28:1547-1552.
- Behbahani M. 2009. Anti-viral activity of the methanolic leaf extract of an Iranian medicinal plant "*Hyssopus officinalis*" against herpes simplex virus. *J Med Plant Res*, 3:1118-1125.
- Behbahani M, Shanehsazzadeh M, Shokoohinia Y, Soltani M. 2013a. Evaluation of Anti-Herpetic Activity of Methanol Seed Extract and Fractions of *Securigerasecuridaca* In vitro. *J Antivir Antiretrovir*, 5:072-076.
- Behbahani M, Zadeh MS, Mohabatkar H. 2013b. Evaluation of antiherpetic activity of crude extract and fractions of *Avicenna marina*, in vitro. *Antiviral Res*, 97:376-380.
- Buzzini P, Arapitsas P, Goretti M, Branda E, Turchetti B, Pinelli P, Ieri F, Romani A. 2008. Antimicrobial and antiviral activity of hydrolysable tannins. *Mini Rev Med Chem*, 8:1179-1187.
- Carnat AP, Carnat A, Fraisse D, Lamaison JL. 1998. The aromatic and polyphenolic composition of lemon balm (*Melissa*

Iranian herbal medicines against *in vitro* replication of HSV

- officinalis L. subsp. officinalis) tea. Pharm Acta Helv, 72:301-305.
- Chang C-C, Yang MH, Wen HM, Chern JC. 2002. Estimation of total flavonoid content in propolis by two complementary colorimetric methods. J Food Drug Anal, 10:178-182.
- Dixit VP, Jain P, Joshi SC. 1988. Hypolipidaemic effects of Curcuma longa L and Nardostachysjatamansi, DC in triton-induced hyperlipidaemic rats. Indian J Physiol Pharmacol, 32:299-304.
- Duke N. 1991. A systematic revision of the mangrove genus Avicennia (Avicenniaceae) in Australasia. AustSyst Bot, 4:299-324.
- Eftekhari F, Zamani S, Yusefzadi M, Hadian J, NejadEbrahimi S. 2011. Antibacterial activity of Zataria multiflora Boiss essential oil against extended spectrum β lactamase produced by urinary isolates of Klebsiella pneumonia. Jundishapur J Microbiol, 4: S43-S49.
- Elion GB. 1993. Acyclovir: discovery, mechanism of action, and selectivity. J Med Virol 41(Suppl 1):2-6.
- Farahani M. 2013. Inhibition of HSV-1 multiplication by five species of medicinal plants. J Microbiol, Biotechnol Food Sci, 3:69-71.
- Garjani A, Fathiazad F, Zakheri A, Akbari NA, Azarmie Y, Fakhrjoo A, Andalib S, Maleki-Dizaji N. 2009. The effect of total extract of Securigerasecuridaca L. seeds on serum lipid profiles, antioxidant status, and vascular function in hypercholesterolemic rats. J Ethnopharmacol, 126:525-532.
- Geuenich S, Goffinet C, Venzke S, Nolkemper S, Baumann I, Plinkert P, Reichling J, Keppler OT. 2008. Aqueous extracts from peppermint, sage and lemon balm leaves display potent anti-HIV-1 activity by increasing the virion density. Retrovirology, 5:27.
- Girish C, Pradhan SC. 2008. Drug development for liver diseases: focus on picroliv, ellagic acid and curcumin. Fundam Clin Pharmacol, 22:623-632.
- Hill C, McKinney E, Lowndes CM, Munro H, Murphy G, Parry JV, Gill ON. 2009. Epidemiology of herpes simplex virus types 2 and 1 amongst men who have sex with men attending sexual health clinics in England and Wales: implications for HIV prevention and management. Euro Surveill, 14:34-9.
- Hogg RW, Gillan FT. 1984. Fatty acids, sterols and hydrocarbons in the leaves from eleven species of mangrove. Phytochemistry, 23:93-97.
- Jury SL, Reynolds T, Cutler DF, Evans F. 1987. The Euphorbiales. Chemistry, taxonomy and economic botany. London: (Bot. J. Linn. Soc. 94.) Academic Press, pp, 3-46
- Karimi A, Moradi MT, Saedi M, Salimzadeh L, Rafieian-Kopaei M. 2012. The Inhibitory Effects of Eucalyptus Extract on Herpes Simplex Virus-1 Replication in Baby Hamster Kidney Cells. Qom Univ Med Sci J, 6:3-8.
- Karimi A, Moradi MT, Saeedi M, Asgari S, Rafieian-Kopaei M. 2013. Antiviral activity of Quercuspersica L.: High efficacy and low toxicity. Adv Biomed Res, 2:36.
- Kaur G, Hamid H, Ali A, Alam MS, Athar M. 2004. Antiinflammatory evaluation of alcoholic extract of galls of Quercusinfectoria. J Ethnopharmacol, 90:285-292.
- Kazazi H, Rezaei K, Ghotb-Sharif SJ, Emam-Djomeh Z, Yamini Y. 2007. Supercritical fluid extraction of flavors and fragrances from *Hyssopus officinalis* L. cultivated in Iran. Food chem, 105:805-811.
- Khosravi A, Behzadi A. 2006. Evaluation of the antibacterial activity of the seed hull of Quercusbrantii on some gram negative bacteria. Pak J Med Sci, 22:429-432.
- Kim HJ, Yoo HS, Kim JC, Park CS, Choi MS, Kim M, Choi H, Min JS, Kim YS, Yoon SW, Ahn JK. 2009. Antiviral effect of Curcuma longa Linn extract against hepatitis B virus replication. J Ethnopharmacol, 124:189-196.
- Kitazato K, Wang Y, Kobayashi N. 2007. Viral infectious disease and natural products with antiviral activity. Drug Discov Ther, 1:14-22.
- Koelle DM, Corey L. 2008. Herpes simplex: insights on pathogenesis and possible vaccines. Annu Rev Med, 59:381-395.
- König G, Rimpler H. 1985. Iridoid glucosides in Avicennia marina. Phytochemistry, 24:1245-1248.
- Li A, Xie Y, Qi F, Li J, Wang P, Xu S, Zhao L. 2009. Anti-virus effect of traditional Chinese medicine Yi-Fu-Qing granule on

- acute respiratory tract infections. *Biosci Trends*, 3:119-123.
- Luper S. 1999. A review of plants used in the treatment of liver disease: part two. *Altern Med Rev*, 4:178-188.
- Maheshwari RK, Singh AK, Gaddipati J, Srimal RC. 2006. Multiple biological activities of curcumin: A short review. *Life Sci*, 78:2081-2087.
- Malvy D, Ezzedine K, Lancon F, Halioua B, Rezvani A, Bertrais S, Chanzy B, Malkin JE, Morand P, De Labareyre C, Hercberg S, El Hasnaoui A. 2007. Epidemiology of orofacial herpes simplex virus infections in the general population in France: results of the HERPIMAX study. *J Eur Acad Dermatol Venereol*, 21:1398-1403.
- Mardani M, Motamedifar M, Hoseinipour R. 2012. A Study of the Antiviral Effect of the Essential oil of *Zataria Multiflora Boiss* on Herpes Simplex Type 1 in Vero Cell Culture. *J Dent Shiraz Univ Med Sci*, 13:s414-s420.
- Mohammadi-Kamalabadi M, Karimi A, Rafieian-Kopaei M, Amjad L. 2014. Phytochemical study and antiviral effect evaluation of methanolic extract with fractions of aerial parts of *euphorbia spinidens*. *J Babol Univ Med Sci*, 16:25-34.
- Monavari S, ShamsiShahrabadi M, Mortazkar P. 2008. The study of antiviral effects of glycyrrhizaglabra extract on HSV. *J Med Pharm*, 4: 81-86.
- Moradi MT, Karimi A, Rafieian M, Kheiri S, Saedi M. 2011. The inhibitory effects of myrtle (*Myrtuscommunis*) extract on Herpes simplex virus-1 replication in Baby Hamster Kidney cells. *J Shahrekord Univ Med Sci*, 12:54-61.
- Motamedifar M, Ghafari N, Talezadeh Sp. 2010. The effect of cumin seed extracts against herpes simplex virus type 1 in vero cell culture. *Iran J Med Sci*, 35:304-309.
- Motamedifar M, Nekoeian A, Moatari A. 2007. The Effect of Hydroalcoholic Extract of Olive Leaves against Herpes Simplex Virus Type 1. *Iran J Med Sci*, 32:222-226.
- M. Motamedifar, A.A. Nekoeian, A. Moatari Mozaffarian V. 1996. A dictionary of Iranian plant names: Latin, English, Persian. Tehran: Farhang Moaser.
- Muliawan SY, Kit LS, Devi S, Hashim O, Yusof R. 2006. Inhibitory potential of *Quercuslusitanica* extract on dengue virus type 2 replication. *Southeast Asian J Trop Med Public Health*, 37:132-135.
- Namazi R, Zabihollahi R, Behbahani M, Rezaei A. 2013. Inhibitory Activity of *Avicennia marina*, a Medicinal Plant in Persian Folk Medicine, against HIV and HSV Iran *J Pharm Res*, 12:435-443.
- Nolkemper S, Reichling J, Stintzing FC, Carle R, Schnitzler P. 2006. Antiviral effect of aqueous extracts from species of the Lamiaceae family against Herpes simplex virus type 1 and type 2 in vitro. *Planta med*, 72:1378-1382.
- Perfect MM, Bourne N, Ebel C, Rosenthal SL. 2005. Use of complementary and alternative medicine for the treatment of genital herpes. *Herpes*, 12:38-41.
- Pouramir M, Shahaboddin ME, Moghadamnia A-A, Parastouei K. 2011. To study the effects of *Securigerasecuridaca* (L.) seed against alloxan-induced hyperglycemia. *J Med Plants Res*, 5:3188-3191.
- Rakić S, Petrović S, Kukić J, Jadranin M, Tešević V, Povrenović D, Šiler-Marinković S. 2007. Influence of thermal treatment on phenolic compounds and antioxidant properties of oak acorns from Serbia. *Food Chem*, 104:830-834.
- Saffarzadeh A, Vincze L, Csapo J. 1999. Determination of the chemical composition of acorn (*Quercusbranti*), *Pistaciaatlantica* and *PistaciaKhinjnk* seeds as non-conventional feedstuffs. *Acta Agraria Kaposváriensis*, 3:59-69.
- Saller R, Buechi S, Meyrat R, Schmidhauser C. 2001. Combined herbal preparation for topical treatment of Herpes labialis. *Forsch Komplementarmed Klass Naturheilkd*, 8:373-382.
- Sayedipour SS, Behbahani M, Moshtaghian SJ. 2012. Evaluation of Anti-Herpes Simplex Virus Type 2 (HSV-2) Activity of Methanol Extract of *Securigerasecuridaca* by Cell Culture Method. *Genetics in the 3rd millennium*, 10:2802-2809.
- Sharangi AB. 2009. Medicinal and therapeutic potentialities of tea (*Camellia sinensis* L.) – A review. *Food Res Int*, 42:529-535.
- Shelton RM. 1991. Aloe vera. Its chemical and therapeutic properties. *Int J Dermatol*, 30:679-683.
- Sydiskis RJ, Owen DG, Lohr JL, Rosler KH, Blomster RN. 1991. Inactivation of enveloped viruses by

- anthraquinones extracted from plants. *Antimicrob Agents Chemother*, 35:2463-2466.
- Tsuchiya Y, Shimizu M, Hiyama Y, Itoh K, Hashimoto Y, Nakayama M, Horie T, Morita N. 1985. Antiviral activity of natural occurring flavonoids *in vitro*. *Chem Pharm Bull (Tokyo)*, 33:3881-3886.
- Villarreal EC. 2003. Current and potential therapies for the treatment of herpesvirus infections. *Prog Drug Res*, 60:263-307.
- Wolbling RH, Leonhardt K. 1994. Local therapy of herpes simplex with dried extract from *Melissa officinalis*. *Phytomedicine*, 1:25-31.
- Zaidi MA, Crow SA, Jr. 2005. Biologically active traditional medicinal herbs from Balochistan, Pakistan. *J Ethnopharmacol* 96:331-334.
- Zandi K, Ramedani E, Mohammadi K, Tajbakhsh S, Deilami I, Rastian Z, Fouladvand M, Yousefi F, Farshadpour F. 2010. Evaluation of antiviral activities of curcumin derivatives against HSV-1 in Vero cell line. *Nat Prod Commun*, 5:1935-1938.
- Zandi K, Taherzadeh M, Yaghoubi R, Tajbakhsh S, Rastian Z, Sartavi K. 2009. Antiviral activity of *Avicennia marina* against herpes simplex virus type 1 and vaccine strain of poliovirus (An *in vitro* study). *J Med Plant Res* 3:771-775.
- Zandi K, Zadeh MA, Sartavi K, Rastian Z. 2007. Antiviral activity of *Aloe vera* against herpes simplex virus type 2: An *in vitro* study. *Afr J Biotechnol*, 6:1770-1773.
- Zargari A. 1990. Medicinal plants. Tehran: Tehran University Press.
- Zatula VV, Kovalev IP, Kolesnikov DG. 1969. Configuration of securigenin and securigenol. *Chem Nat Compd*, 5:111-112.