

Original Research Article

## The antioxidant activity of *Beta vulgaris* leaf extract in improving scopolamine-induced spatial memory disorders in rats

Shadieh Hajhosseini<sup>1</sup>, Mahbubeh Setorki<sup>2\*</sup>, Zahra hooshmandi<sup>1</sup>

<sup>1</sup> Department of Biology, Sanandaj Branch, Islamic Azad University, Sanandaj, Iran

<sup>2</sup> Department of Biology, Izeh Branch, Islamic Azad University, Izeh, Iran

**Article history:**

Received: Aug 11, 2016

Received in revised form:

May 05, 2017

Accepted: May 11, 2017

**\* Corresponding Author:**

Tel: +09133121589

Fax: +9877654098

doctor.setorgi@gmail.com

**Keywords:**

*B. vulgaris*

Leaf extract

Antioxidant activity

Spatial memory

**Abstract**

**Objective:** Medicinal plants have attracted global attention due to their safety as well as their considerable antioxidant content that helps to prevent or ameliorate various disorders including memory impairments. This study was conducted to investigate the effect of beet root (*Beta vulgaris*) leaf extract on scopolamine-induced spatial memory impairments in male Wistar rats.

**Materials and Methods:** Male Wistar rats were randomly divided into 5 groups (n=10): Control (C), scopolamine 1 mg/kg/day (S), scopolamine+50 mg/kg *B. vulgaris* leaf extract (S+B 50), scopolamine+100 mg/kg *B. vulgaris* leaf extract (S+B 100) and scopolamine+200 mg/kg *B. vulgaris* leaf extract (S+B 200). Morris water maze task was used to assess spatial memory. Serum antioxidant capacity and malondialdehyde (MDA) level were also measured.

**Results:** Group S spent significantly less time in the target quadrant compared to the control group, and the administration of *B. vulgaris* leaf extract (100 and 200 mg/kg) significantly increased this time ( $p < 0.05$ ). Scopolamine decreased serum antioxidant capacity and increased serum MDA level yet insignificantly. *B. vulgaris* extract (200 mg/kg) significantly increased the antioxidant capacity and decreased serum MDA level in scopolamine-treated rats ( $p < 0.05$ ).

**Conclusion:** Our results suggested that *B. vulgaris* leaf extract could ameliorate the memory impairments and exhibited protective effects against scopolamine-induced oxidation. Further investigation is needed to isolate specific antioxidant compounds from *B. vulgaris* leaf extract with protective effect against brain and memory impairments.

Please cite this paper as:

Hajhosseini S, Setorki M, Hooshmandi Z. The antioxidant activity of *Beta vulgaris* leaf extract in improving scopolamine-induced spatial memory disorders in rats. Avicenna J Phytomed 2017 Epub

## Introduction

Learning and memory play key roles in daily human life. Learning is the basis of all teaching and training programs. By identifying compounds that can improve

these two processes, mental and memory status can be greatly improved. Studies showed that many drugs can affect memory and learning, for example,

cholinergic drugs have a positive effect on memory, while anticholinergic and anesthetic drugs have negative effects on memory (Uttara et al., 2007; Moopanar et al., 2005; Hosseini et al., 2015). Scopolamine is an anticholinergic drug that blocks the action of acetylcholine within the nervous system. It has also been shown that scopolamine-induced cholinergic blockade induces significant memory impairment. Scopolamine administration is also associated with enhanced brain oxidative stress (Warner et al., 2004).

High rate of reactive oxygen species and free radicals production due to oxidative stress can lead to peroxidation of biomolecules such as lipids, proteins and damage to DNA that cause memory impairments (Sulakhiya et al., 2016). Brain tissue is more sensitive to the effect of free radicals due to its high oxygen demands, abundant lipid content, and relative paucity of antioxidant enzymes compared to other organs (Rasoolijazi et al., Vincent et al., 2004).

The use of medicinal plants with high levels of antioxidant components is one of the most efficient ways to minimize the adverse effects of free radicals and treat diseases. While synthetic drugs may be unsafe for human and the environment, medicinal plants and their derivatives are growingly accepted worldwide due to their safety (Sulakhiya et al., 2016; Zaidi et al., 2014). In this regard, it has been frequently shown that antioxidant agents reverse scopolamine-induced memory impairments (Hosseini et al., 2015; Yang et al., 2009).

Beetroot (*Beta vulgaris*) is a plant from the family Chenopodiaceae. *B. vulgaris* is native to Mediterranean regions and is extensively cultivated in different countries including Iran (Sulakhiya et al., 2016). Roots and leaves of *B. vulgaris* are used in traditional medicine, to treat different diseases. *B. vulgaris* leaves have diuretic, purgative, laxative, and aphrodisiac activity (Sulakhiya et al.,

2016; Jain et al., 2011; El Gamal et al., 2014). *B. vulgaris* leaves contain various phytoconstituents such as betalains, flavonoids, polyphenols, vitamins, and minerals (Sulakhiya et al., 2016; Jain et al., 2012; Bolkent et al., 2007; Mroczek et al., 2012). *B. vulgaris* leaves have antioxidant, anticancer, hepatoprotective, nephroprotective, wound healing, and anti-inflammatory activities (Sulakhiya et al., 2016; El Gamal et al., 2014; Babu et al., 2010; Jain et al., 2011; Sacan et al., 2010), but its potential neuroprotective effects have not yet been tested.

Considering the important role of reactive oxygen species and free radicals in brain oxidative damage and memory impairments as well as the antioxidant potential of *B. vulgaris* leaf extract, this study was conducted to evaluate the effect of *B. vulgaris* leaf extract on scopolamine-induced brain tissue damage in rats.

## Materials and Methods

### Preparation of ethanolic *B. vulgaris* leaf extract

Fresh *B. vulgaris* leaves were shade-dried, pulverized, and macerated with 97% ethanol for 5 hr. The resulting extract was then filtered, and the filtrate was concentrated and dried using vacuum distillation at 40°C (Al-Jassabi et al., 2013)

### Determination of antioxidant activity of *B. vulgaris* leaf extract

Briefly, 2 ml of *B. vulgaris* leaf extract or butylated hydroxytoluene (BHT) at different concentrations (prepared in ethanol) were added to 2 mL of DPPH<sup>1</sup> solution (0.1 Mm in ethanol). After 15 min at room temperature in the dark, the absorbance of samples were measured at 517 nm wavelength. The mixture of ethanol (2 mL) and DPPH solution (2 mL) served as control. The scavenging activity

---

1 1,1-Diphenyl-2-picryl-hydrazyl

percentage was determined by the following equation (Rabiei et al., 2014)

$$(AA\%) = 100 \times (A_{\text{control}} - A_{\text{sample}}) / A_{\text{control}}$$

AA; scavenging activity

$A_{\text{control}}$ : absorbance of control

$A_{\text{samples}}$ : absorbance of samples

#### **Determination of total phenolic content of *B. vulgaris* leaf extract**

Briefly, 0.1 ml of diluted extract (0.01 g in 10 ml of 60°C methanol) was mixed with 0.5 ml of Folin-Ciocalteu reagent. After 3-5 min, 0.4 mL of 7.5% sodium carbonate solution was added to the mixture and left at room temperature for 30 min. The mixture absorbance was measured at 750 nm wavelength against distilled water blank. A standard calibration curve was plotted using different concentrations of gallic acid. The phenolic content was expressed as “mg gallic acid equivalents (GAEs)/g of the sample” (Rabiei et al., 2014).

#### **Determination of total flavonoid contents of *B. vulgaris* leaf extract**

The total flavonoid content of *B. vulgaris* leaf extract was determined using colorimetric method. The diluted extract (0.5 ml) was mixed with 0.5 ml of 2% aluminum chloride and 3 ml of 5% potassium acetate. After 40 min of incubation at room temperature, the absorbance of the reaction mixture was measured at 415 nm wavelength. A standard curve was prepared using different concentrations of rutin solution. The total flavonoid content was expressed as “mg of rutin equivalents/g of dried extract” by using a standard curve of rutin (Rabiei et al., 2014).

#### **Determination of total flavonolic content of *B. vulgaris* leaf extract**

The diluted extract (0.1 ml) was mixed with 0.5 ml of 2% aluminum chloride and 3 ml of 5% potassium acetate and left at room temperature for 2.5 hr. After incubation, the mixture absorbance was

measured at 415 nm wavelength. Total flavonolic content was expressed in terms of rutin equivalent (mg/g), which is a common reference compound (Rabiei et al., 2014).

#### **Animals and treatments**

Adult male Wistar rats (weighing 200–250 g) were kept in experimental animal care center of Islamic Azad University of Sanandaj, Sanandaj, Iran. Rats were maintained under standard laboratory conditions of  $25 \pm 2^\circ\text{C}$  temperature and  $55 \pm 5\%$  humidity with 12 hr light-dark cycle with free access to water and standard laboratory food. The rats were randomly divided into 5 groups of 10 each. Control group (C) received distilled water alone intraperitoneally (ip); scopolamine-treated group (S) received distilled water 30 min after scopolamine injection (1 mg/kg/day, ip); Extract-treated groups received *B. vulgaris* leaf extract (50, 100, and 200 mg/kg/day, ip) 30 min after scopolamine injection for 14 days. The doses of scopolamine and *B. vulgaris* extract were selected based on the previous studies (Hosseini et al., 2015; Sulakhiya et al., 2016)

After treatment, spatial memory task was performed using Morris water maze. Then, the animals were put under deep anesthesia using chloral hydrate and then, cardiac blood samples were collected. The blood samples were centrifuged and their serum was isolated. Then, the biochemical analysis was carried out.

#### **Spatial memory**

To evaluate spatial memory, the rats were tested in Morris water maze which is a black circular tank with a diameter of 136 cm and height of 60 cm high, filled with  $24 \pm 1^\circ\text{C}$  water to a depth of 25 cm. A hidden circular platform (diameter: 10 cm) was located in the center of the Southwest quadrant and submerged 1 cm below the surface of the water. The maze was located in a room containing many visual cues such as bookshelves,

refrigerator, and poster. Each rat experienced two sessions of four trials per day for 5 consecutive days. During each trial, the rats were individually placed in the pool and released in the release positions randomly determined for each trial by computer. The rats were allowed to swim until they found the platform and remained on it for 30 s. If 60 s passed and the rats did not find the platform, they were gently guided to the platform. After each session, the rats were returned to their cages for resting. The time latency to reach the platform was recorded by a video tracking system. On day 5, to conduct a probe trial, platform was elevated above the water surface and time spent by the rats in the target quadrant (Q1) where platform was located, was recorded on days 1–4 (Gray *et al.*, 2003).

#### Measurement of serum antioxidant capacity

Three solutions were used to measure serum antioxidant level: Solution 1 consisted of 1.5 ml of sodium acetate and 8 ml of concentrated acetic acid, diluted to 500 mL with distilled water; solution 2 consisted of 270 mg of iron (III) chloride, dissolved in 50 ml of distilled water and solution 3 consisted of 47 mg of Treeazin, dissolved in 40 ml of HCl. Working solution was prepared by mixing 10 ml of solution 1, 1 ml of solution 2 and 1 ml of solution 3. Thereafter, 25  $\mu$ l of serum samples was added to 5.1 ml of the working solution. The resulting mixture was incubated at 37°C for 15 min; then, the absorbance was measured at 593 nm wavelength (Rabiei *et al.*, 2014).

#### Measurement of serum malondialdehyde (MDA) level

Briefly, 50  $\mu$ L of the sera was mixed with 50  $\mu$ L of 0.05% BHT, 400  $\mu$ L of 0.44 M H<sub>3</sub>PO<sub>4</sub> and 100  $\mu$ L of 42 Mm TBA. The mixture was vortexed and then, heated in a boiling water bath for 1 hr. After cooling at 0°C for 5 min, 250  $\mu$ L of n-butanol was added to the mixture, vortexed, and then,

centrifuged at 14000 rpm for 5 min. The supernatant absorbance was measured at 532 nm wavelength (Biondo *et al.*, 2014; Rabiei *et al.*, 2014)

#### Statistical analysis

SPSS 18 was used to conduct data analysis. All data were expressed as mean  $\pm$  SD. One-way ANOVA followed by Duncan's test was used to compare the means among experimental groups. A  $p < 0.05$  was considered statically significant.

## Results

### Standardization of *B. vulgaris* leaf extract

To standardize the plant extract, total phenolic flavonoid and flavonol compounds present in *B. vulgaris* leaf extract were measured. Total phenolic content of *B. vulgaris* leaf extract was 51 mg/g GAE expressed as mg phenol/g of dry matter. Total flavonoid and flavonolic contents were 17.3 mg/g and 2.5 mg/g, respectively, expressed as mg rutin equivalent/g of dry matter. The percentage of DPPH radical scavenging of extract is shown in Table 1.

Table 1: DPPH radical scavenging activities for various concentrations of *Beta vulgaris* leaf extract.

samples	concentration	% of DPPH radical scavenging
<i>Beta vulgaris</i> leaf extract	10	10.39
	20	18.01
	50	21.93
	100	39.95(IC50)
	250	81.52

### Morris Water Maze Test

The time spent in the target quadrant during the 60-sec probe trial is illustrated in Figure 1. The control group spent significantly more time in the target quadrant (when the hidden platform was there during the previous days) compared to the scopolamine-treated group ( $p < 0.05$ ). Administration of *B. vulgaris* leaf extract

*Beta vulgaris* leaves extract ameliorates scopolamine-induced spatial memory impairments

at 100 and 200 mg/kg doses significantly increased the time spent in target quadrant in comparison with the scopolamine group ( $p < 0.05$ ).

The latencies to reach the platform during 4 consecutive days are shown in Figure 2. In the control group, the latency to find the platform during 4 consecutive days were shorter than scopolamine-treated group yet insignificantly.

Treatment with *B. vulgaris* leaf extract at doses of 50, 100 and 200 mg/kg resulted in shorter latency time compared to the scopolamine group during the study period yet insignificantly. In rats treated with *B. vulgaris* extract at 200mg/kg, the latency to reach the platform on day 3 was significantly lower than that in scopolamine-treated rats ( $p < 0.05$ ).




Figure 1. The time spent in target quadrant during the probe trial; the data are expressed as mean  $\pm$  SD;  $n = 10$  in each group; SCO+BV= scopolamine+ *B. vulgaris*. \* shows significant difference compared to control group ( $*P < 0.05$ ); + shows significant difference compared to scopolamine group ( $+++p < 0.001$ ;  $+P < 0.05$ ).




Figure 2. Comparison of latency time (sec) to find the platform among different experimental groups; the data are expressed as mean  $\pm$  SD;  $n = 10$  in each group; SCO+BV= scopolamine+ *B. vulgaris*. + Shows significant difference compared to scopolamine group ( $+P < 0.05$ )

**Effect of *B. vulgaris* leaf extract on serum antioxidant capacity**

As Figure 3 illustrates, there was no significant difference in serum antioxidant capacity between control and scopolamine-treated rats. Administration of *B. vulgaris* leaf extract at concentrations of 50, 100 and 200 mg/kg to scopolamine-treated rats, increased serum antioxidant capacity, and the difference was significant at the dose of 200 mg/kg compared to the scopolamine group ( $p < 0.05$ ).

The effect of *B. vulgaris* leaf extract on serum MDA level in rats is shown in Figure 4. The scopolamine-treated rats presented higher serum MDA levels when compared to control group ( $p > 0.05$ ). Administration of *B. vulgaris* leaf extract 50, 100 and 200 mg/kg decreased serum MDA level in comparison with scopolamine group, and 200 mg/kg of *B. vulgaris* leaf extract caused the lowest serum MDA level with a significant difference compared to other concentrations.

**Effect of *B. vulgaris* leaf extract on serum MDA level**




Figure 3. The effect of *Beta vulgaris* leaf extract on serum antioxidant capacity; the data are expressed as mean  $\pm$  SD; n = 10 in each group; SCO+BV= scopolamine+ *B. vulgaris*; <sup>+</sup> shows significant difference compared to scopolamine group.




Figure 4. The effect of *Beta vulgaris* leaf extract on serum MDA level in rats; the data are expressed as mean  $\pm$  SD; n = 10 in each group; SCO+BV= scopolamine+ *B. vulgaris*; <sup>+</sup> shows significant difference compared to scopolamine group.

## Discussion

The results of the present study indicated that scopolamine led to learning and memory impairments in rats as assessed by Morris water maze test. This result is in agreement with the previous studies (Chuong et al., 2014; Rabiei et al., 2015). Scopolamine is considered a cholinergic antagonist which interferes with the transmission of acetylcholine in the central nervous system leading to memory deficits (Misane et al., 2003).

The results also indicated that the treatment of scopolamine-treated rats with *B. vulgaris* leaf extract at 100 and 200 mg/kg significantly increased the time spent in target quadrant of Morris water maze ( $p < 0.05$ ). *B. vulgaris* extract (100 mg/kg) also significantly increased latency to reach the platform on day 3. These results showed the protective effects of *B. vulgaris* extract on scopolamine-induced memory deficits.

The results of the present study showed that scopolamine resulted in reduction of serum antioxidant capacity, but it increased serum MDA levels yet insignificantly. These results are in agreement with studies that reported increased levels of MDA after treatment with scopolamine (Rabiei et al., 2015; Wilson, 1997). Increase in MDA, the most well-known secondary product of lipid peroxidation, is a valid marker for *in vivo* lipid peroxidation and can be used to determine the severity of oxidative damage to cells and tissues (Grotto et al., 2009)

In our study, administration of *B. vulgaris* leaf extract at 200 mg/kg significantly increased the antioxidant capacity and decreased serum MDA levels compared to scopolamine group. Jain et al. (2012) also reported that treatment with *B. vulgaris* leaf extract significantly decreased hepatic MDA level and increased GSH, suggesting that the antioxidant effect of *B. vulgaris* leaf extract plays an important protective role against ethanol-mediated toxicity (Jain et al., 2012).

*B. vulgaris* leaf also could improve antioxidant status in mice fed with high-fat diet (Jain et al., 2012). The results of this study indicated that *B. vulgaris* leaf extract exhibited protective activity against the adverse effects of scopolamine; therefore, it could reduce the brain damage and improve brain and memory activities in scopolamine-treated rats. These findings may be associated with antioxidant activity of *B. vulgaris* leaf extract and inhibition of oxidative stress in the rats' brains.

In the present study, we observed that *B. vulgaris* leaf extract at 100  $\mu\text{g/ml}$  could inhibit 39.95% of DPPH radicals that reached 81% when the dose of 250  $\mu\text{g/ml}$  of this extract was administered. These results indicate remarkable antioxidant and free radical-scavenging activity of *B. vulgaris* leaf extract which is comparable to those of other antioxidant agents. The free radical-scavenging activity of *B. vulgaris* leaf extract can be due to the presence of bioactive compounds with antioxidant properties such as polyphenols, flavonoids, and vitamin C. The total phenolic content in this plant was 51 mg GAE/1 g of the dried extract. Jain et al., (2012) reported that the main constituents of *B. vulgaris* leaf are sterols, triterpenoids, phenols, tannins, flavonoids, alkaloids, glycosides, and saponins (Jain et al., 2012). Polyphenols, flavonoids, saponins, glycolipid, phospholipids, fatty acids, folic acid, and ascorbic acid are some of the major components of *B. vulgaris* leaf extract that give antioxidant property to this extract. In addition, folic acid, iron, calcium, phosphorus, zinc, and vitamins A, B, and C can play important roles in brain development and motor function (BolKent et al., 2007; Mroczek et al., 2012; Mokhtari-Dehkordi et al., 2014). Sulakhiya et al., (2016) reported that *B. vulgaris* leaf extract had desirable effect on depression which was attributed to the high levels of

antioxidant compounds such as polyphenols (betalains and betaine), flavonoids, and vitamin C in *B. vulgaris* leaf.

The protective effect of *B. vulgaris* leaf extract on scopolamine-induced memory impairment may be related to increases in acetylcholine levels in the brain. We found no previous study on cholinergic effect of *B. vulgaris* leaf extract; therefore, it is recommended to further evaluate the cholinergic effect of *B. vulgaris* extract.

The results of this study indicated that *B. vulgaris* leaf extract exhibited protective activity against scopolamine-induced brain and memory impairments. This effect may be associated with increased activity of antioxidant defense system and inhibition of oxidative stress in the brains of rats. *B. vulgaris* leaf extract can be used as a beneficial medicinal herb for improving brain and memory complications due to high levels of polyphenolic antioxidant compounds.

### Conflict of interest

The authors have no conflicts of interest to declare.

### References

Al-Jassabi S, Abdullah M. 2013. Purification and characterization of antioxidant fractions from *Zizyphus spina-christi*. *Fruits*, 5: 13-24.

Babu VA, Gowri R. 2010. Evaluation of antioxidant activity of *Beta vulgaris* root extract in rats. *Asian J Chem*, 22: 3385-90.

Biondo I PB, Boeing I SJ, Barizão I EO, Souza I NE, Matsushita I M, Oliveira I C, Boroski I M, Visentainer I JV. 2014. Evaluation of beetroot (*Beta vulgaris* L.) leaves during its developmental stages: a chemical composition study. *Food Sci Technol*, 34: 94-101.

Bolkent Ş, Yanardağ R, Tabakoğlu-Oğuz A, Özsoy-Saçan Ö. 2007. Effects of chard (*Beta vulgaris* L. var. *cicla*) extract on pancreatic B cells in streptozotocin-diabetic rats: a

morphological and biochemical study. *J Ethnopharmacol*, 73: 251-259.

Chuong NN, Trung BH, Luan TC, Hung TM, Dang NH, Dat NT. 2014. Anti-amnesic effect of alkaloid fraction from *Lycopodiella cernua* (L.) Pic. Serm. on scopolamine-induced memory impairment in mice. *Neurosci Lett*, 575:42-46.

El Gamal AA, AlSaid MS, Raish M, Al-Sohaibani M, Al-Massarani SM, Ahmad A, et al. 2014. Beetroot (*Beta vulgaris* L.) extract ameliorates gentamicin-induced nephrotoxicity associated oxidative stress, inflammation, and apoptosis in rodent model. *Mediators Inflamm*, 22: 133-145.

Gray C, Nukada H, Jackson DM, McMorran PD, Wu A, Ma F. 2003. Neuroprotective effects of nitron radical scavenger S-PBN on reperfusion nerve injury in rats. *Brain Res*, 982:179-185.

Grotto D, Maria LS, Valentini J, Paniz C, Schmitt G, Garcia SC. 2009. Importance of the lipid peroxidation biomarkers and methodological aspects for malondialdehyde quantification. *Quimica Nova*, 32:169-174.

Hosseini M, Mohammadpour T, Karami R, Rajaei Z, Sadeghnia HR, Soukhtanloo M. 2015. Effects of the hydro-alcoholic extract of *Nigella Sativa* on scopolamine-induced spatial memory impairment in rats and its possible mechanism. *Chinese J Integ Med*, 21: 438-444.

Jain NK, Singhai AK. 2012. Protective role of *Beta vulgaris* L. leaves extract and fractions on ethanol-mediated hepatic toxicity. *Acta Pol Pharm*, 69: 945-950.

Jain S, Garg VK, Sharma PK. 2011. Anti-inflammatory activity of aqueous extract of *Beta vulgaris* L. *J Basic Clin Pharm*, 2: 83-89.

Misane I, Ogren SO. 2003. Selective 5-HT1A antagonists WAY 100635 and NAD-299 attenuate the impairment of passive avoidance caused by scopolamine in the rat. *Neuropsychopharmacol*, 28:253-264.

Mokhtari-Dehkordi S, Hojjati M, Rouhi L, Rabiei Z, Alibabaei Z. 2014. Effect of ethanolic extract of beet roots and leaves on motor coordination in male Wistar rats. *J Shahrekord Uni Med Sci*, 16: 14-23.



***Beta vulgaris* leaves extract ameliorates scopolamine-induced spatial memory impairments**

- Moopanar TR, Allen DG. 2005. Reactive oxygen species reduce myofibrillar Ca<sup>2+</sup> sensitivity in fatiguing mouse skeletal muscle at 37 C. *J Physiol*, 564: 189-199.
- Mroczek A, Kapusta I, Janda B, Janiszowska W. 2012. Triterpene saponin content in the roots of red beet (*Beta vulgaris* L.) cultivars. *J Agric Food Chem*, 60: 12397-12402.
- Rabiei Z, Alibabaei Z, Rafieian-Kopaei M. 2015. Determining the Antioxidant properties of chamomile and investigating the effects of chamomile ethanol extract on motor coordination disorders in rats. *J Babol Univ Med Sci*, 17:44-50.
- Rabiei Z, Rafieian-kopaei M, Heidarian E, Saghaei E, Mokhtari S. 2014. Effects of *Zizyphus jujube* extract on memory and learning impairment induced by bilateral electric lesions of the nucleus basalis of meynert in rat. *Neurochem Res*, 39:353-360.
- Rasoolijazi H, Mehdizadeh M, Soleimani M, Nikbakhte F, Farsani ME, Ababzadeh S. 2015. The effect of rosemary extract on spatial memory, learning and antioxidant enzymes activities in the hippocampus of middle-aged rats. *Med J Islam Repub Iran*, 29:187-193.
- Sacan O, Yanardag R. 2010. Antioxidant and antiacetylcholinesterase activities of chard (*Beta vulgaris* L. var. cicla). *Food Chem Toxicol*, 48: 1275-1280.
- Sulakhiya K, Patel VK, Saxena R, Dashore J, Srivastava AK, Rathore M. 2016. Effect of *Beta vulgaris* Linn. leaves extract on anxiety- and depressive-like behavior and oxidative stress in mice after acute restraint stress. *Pharm Res*, 8: 11-12.
- Uttara B, Singh AV, Zamboni P, Mahajan RT. 2007. Oxidative stress and neurodegenerative diseases: A review of upstream and downstream antioxidant therapeutic options. *Current Neuropharmacol*, 7: 65-74.
- Vincent AM, Russell JW, Low P, Feldman EL. 2004. Oxidative stress in the pathogenesis of diabetic neuropathy. *Endoc Rev*, 25: 612-28.
- Warner DS, Sheng H, Batinić-Haberle I. 2004. Oxidants, antioxidants and the ischemic brain. *J Exp Biol*, 207: 3221-31.
- Wilson JX. 1997. Antioxidant defense of the brain: a role for astrocytes. *Canadian J Physiol Pharmacol*, 75: 1149-63.
- Yang MH, Yoon KD, Chin Y-W, Park JH, Kim SH, Kim YC, et al. 2009. Neuroprotective effects of *Dioscorea opposita* on scopolamine-induced memory impairment in in vivo behavioral tests and in vitro assays. *J Ethnopharm*, 121:130-134.
- Zaidi SK, Hoda MN, Tabrez S, Ansari SA, Jafri MA, Shahnawaz Khan M. 2014. Protective effect of *Solanum nigrum* leaves extract on immobilization stress induced changes in rat's brain. *J Evid Based Complementary Altern Med*, 3: 22-34.