

Original Research Article

## Naringin mitigates testicular injury and associated neuronal toxicity in lead-exposed cockerel chicks

Oluwaseun Olanrewaju Esan<sup>1</sup>, Chinomso Gift Ebirim<sup>1</sup>, Moses Olusola Adetona<sup>2</sup>, Temitayo Olabisi Ajibade<sup>3,\*</sup>, Ademola Adetokunbo Oyagbemi<sup>3</sup>, Temidayo Olutayo Omobowale<sup>1</sup>, Omolade Abodunrin Oladele<sup>1</sup>, Adeolu Alex Adedapo<sup>4</sup>, Momoh Audu Yakubu<sup>5</sup>, Evaristus Nwulia<sup>6</sup>, Oluwafemi Omoniyi Oguntibeju<sup>7</sup>

<sup>1</sup>Department of Veterinary Medicine, Faculty of Veterinary Medicine, University of Ibadan, Nigeria

<sup>2</sup>Department of Anatomy, Faculty of Basic Medical Sciences, University of Ibadan, Nigeria

<sup>3</sup>Department of Veterinary Physiology and Biochemistry, Faculty of Veterinary Medicine, University of Ibadan, Nigeria

<sup>4</sup>Department of Veterinary Pharmacology and Toxicology, Faculty of Veterinary Medicine, University of Ibadan, Nigeria

<sup>5</sup>Department of Environmental & Interdisciplinary Sciences, College of Science, Engineering & Technology, COPHS, Texas Southern University, Houston, TX, USA

<sup>6</sup>Howard University, College of Medicine, Department of Psychiatry and Behavioral Sciences, Howard University Hospital, 2041 Georgia Avenue, Washington, DC 20060, USA

<sup>7</sup>Department of Biomedical Sciences, Faculty of Health and Wellness Sciences, Cape Peninsula University of Technology, Bellville 7535, South Africa

### Article history:

Received: Oct 20, 2022

Received in revised form:

Feb 25, 2023

Accepted: Feb 25, 2023

Epub ahead of print

### \*Corresponding Author:

Tel: +2349071842142

toajibade@gmail.com

### Keywords:

Lead toxicity

Oxidative stress

Testicular apoptosis

Neuroinflammation

### Abstract

**Objective:** Lead (Pb) poisoning affects multiple organs including the reproductive system. The experiment was performed to explore the protective effect of naringin on testicular apoptosis, neuronal dysfunction and markers of stress in cockerel chicks.

**Materials and Methods:** Thirty-six cockerel chicks were used for this study, and randomly grouped into six chicks per group viz. control, Pb only (600 ppm), Pb and naringin (80 mg/kg), Pb and Naringin (160 mg/kg), naringin only (80 mg/kg) and naringin only (160 mg/kg), respectively, for eight weeks. Pb was administered via drinking water while naringin was administered via oral gavage. Oxidative stress indices in the brain and testes were assessed, and immunohistochemistry of TNF- $\alpha$  and caspase 3 was done in the brain and testes, respectively.

**Results:** Lead administration induced inflammatory and testicular apoptosis cascade accompanied with increased oxidative stress and upregulation of brain and testicular antioxidant enzymes in comparison to the control and Pb-only-treated cockerels. Immunohistochemistry showed significant immunoreactivity of testicular caspase 3 and TNF- $\alpha$  in the brain.

**Conclusion:** Treatment of Pb-exposed chickens with naringin offered protection to Pb acetate-induced testicular oxidative stress, apoptosis, and neuroinflammation in cockerel chicks.

Please cite this paper as:

Esan O.O, Ebirim C.G, Adetona M.O, Ajibade T.O, Oyagbemi A.A, Omobowale T.O, Oladele O.A, Adedapo A.A, Yakubu M.A, Nwulia E, Oguntibeju O.O. Naringin mitigates testicular injury and associated neuronal toxicity in lead-exposed cockerel chicks. Avicenna J Phytomed, 2023. Epub ahead of print.

## Introduction

Lead (Pb) is a common, non-biodegradable contaminant that has a significant impact on human and animal health. Pb is naturally found in small amounts, nevertheless, industrial activities have been reported to considerably increase the concentration of Pb (Al-Megrin *et al.*, 2020). Human exposure to Pb can be through ingestion and inhalation. Pb toxicity is very harmful on people, animals, and plants (Yousef *et al.*, 2019). Pb poisoning is induced by free radical generation and inhibition of antioxidant enzymes. Reactive oxygen species (ROS) can chelate atoms of oxygen and nitrogen causing numerous enzymes and proteins to malfunction (Abdel Moneim *et al.*, 2011). Oxidative insults also affect DNA structure and prevent damaged DNA from being repaired, which can lead to aberrant gene expression.

Pb has also been discovered to elicit inflammation by promoting immune and non-immune cells, causing them to over-secrete inflammatory mediators, which may drive apoptotic signaling pathway (Chen *et al.*, 2019). Reproductive organs are extremely susceptible to stressful situations and can be damaged by any hazardous element with Pb having a direct impact on testicular shape (Khodabandeh *et al.*, 2021). Heavy metals have been reported to inhibit enzymatic function and impact sperm properties such as motility, morphology, and concentration (El-Khadragy *et al.*, 2020). Pb is thought to have an indirect effect on male reproductive function through disruption of the endocrine system (Abiodun and Adeleye, 2020). It also causes spermatozoa death and genetic instability as well as activating apoptosis in spermatogenic cells. It also inhibits spermatogenic activity by lowering sperm calcium levels and decreasing sperm protein phosphorylation. Lead exposure has been reported to have a negative effect on hormone secretion by suppressing

luteinizing hormone and follicle stimulating secretion from the anterior pituitary (El-Khadragy *et al.*, 2020; Kumar and Devi, 2018).

Umbelliferone has been found to prevent testicular injury induced by Pb toxicity via suppressing oxidative damage, inflammation, and cell death, and boosting antioxidant defenses, Nrf2/HO-1 signaling and pituitary-gonadal axis (Alotaibi *et al.*, 2020). Khodabandeh *et al.* (2021) reported that quercetin preserved testes architecture and mitigated apoptosis against Pb-acetate toxicity. In another study, Olaniyan *et al.* (2021) documented that *Cocos nucifera* L. oil alleviates lead acetate-induced reproductive toxicity in sexually matured male Wistar rats. The amino acid L-carnitine was reported to ameliorate chronic toxicity associated with Pb toxicity (Abdel-Emam and Ahmed, 2021). Interestingly, Ibrahim *et al.* (2021) demonstrated that *Chlorella vulgaris* and *Spirulina platensis* mitigated Pb acetate-induced testicular oxidative stress and apoptosis, thereby corroborating the earlier report of Khodabandeh *et al.* (2021). Again, *Allium triquetrum* was found to significantly alleviate Pb acetate-induced reproductive injuries by boosting sperm characteristics and ameliorating oxidative stress markers (Kahalerras *et al.*, 2022). The antioxidative, anti-inflammatory and anti-apoptotic actions of ellagic acid, resveratrol, and *Asparagus officinalis* against Pb acetate-induced testicular and hepato-renal oxidative damages have also been reported (Bidanchi *et al.*, 2022; Mitra *et al.*, 2022; Alyami *et al.*, 2022). Recently, Bentaiba *et al.* (2023) reported that *Withania frutescens* root extract restored testicular damage induced by Pb toxicity.

The damage to the central nervous system (CNS) induced by Pb pollution requires immediate and urgent attention. Pb exposure has been linked to behavioral abnormalities as well as cognitive and mental development deficiencies in a growing number of studies (Ogundele, 2018). Furthermore, in human and animal

## Protection by naringin in lead-exposed chicks

studies, Pb exposure could cause emotional abnormalities or even neuropsychiatric problems (Zhang et al., 2021). Alterations in neurotransmitter system and synaptic signaling, mitochondrial malfunction, and oxidative stress, are all plausible pathways for Pb neurotoxicity, with oxidative stress being a focus of attention (Zhang et al., 2021). ROS are frequently involved in negative effects of oxidative, exaggerated increase in free radicals' generation and reduction in the anti-oxidative status (Ahmad et al., 2018). Pb is known to compete with  $Fe^{2+}$ , and the buildup of free iron causes oxidative stress and neurotoxicity as reported by Ferreira et al. (2017).

The nervous system is particularly sensitive to oxidative stress due to the presence of high metabolic activity, inadequate antioxidant systems, and fatty acid content (Yu et al., 2021). Pb as a neurotoxin interferes with and diminishes ion calcium communication in nerve processes, thereby affecting the cerebral cortex, cerebellum, and hippocampus. There has been a positive correlation between peripheral nerve degeneration and apoptosis in the neural tissue of encephalopathy induced by Pb intoxication (Al-Khafaf et al., 2021).

Naringin (NAR) is a polyphenolic chemical compound found in high concentration in citrus and grapefruits. In experimental trials, it has been shown to have several therapeutic effects against a variety of diseases, including neurodegenerative disorders (Varshney and Garabadu, 2021). In *in-vivo* and *in-vitro* experimental models, NAR was documented to possess antioxidant property (Qin et al., 2016). In numerous animal models, NAR has been shown to improve memory development (de-Andrade Teles et al., 2016). In the animal brain, it has been shown to promote cholinergic function while inhibiting acetylcholinesterase activity (Varshney and Garabadu, 2021).

In a cell culture investigation, it was discovered that NAR improves mitochondrial activity and apoptosis, possibly through heme oxygenase-1 (HO-1)-mediated action (Varshney and Garabadu, 2021). NAR pretreatment has also been shown to increase learning and memory formation in rats (Ghofrani et al., 2015). Recently, it was shown that NAR has therapeutic effects in experimental animals via the renin-angiotensin system (RAS) as previously described by Wang et al. (2019). NAR has also been reported to improve neuroprotection by lowering oxidative stress and enhancing neuronal survival (Gindri et al., 2021). NAR has been documented to undergo enterohepatic biotransformation to release naringenin which can cross the blood brain barrier (BBB) and trigger a variety of protective activities (Wang et al., 2019).

In this work, our aim was to understand the protective actions of NAR on the reproductive and central nervous systems and explore its possible molecular mechanism of action. We also explored the metal-chelating antioxidant activity of NAR on the toxicity of Pb. We hypothesized that the inclusion of NAR in poultry feeds might be beneficial as an antioxidant with metal binding capacity.

## Materials and Methods

### Chemicals

Pb acetate trihydrate (( $C_2H_3O_2$ ) $2Pb.3H_2O$ ); CAS No.: 6080-56-4, naringin; CAS No.: 6751AF, O-dianisidine; CAS No.: 119-90-4, H<sub>2</sub>O<sub>2</sub>; CAS No.: 7722-84-1, xylenol orange (XO); CAS No.: F943, potassium hydroxide; CAS No.: 1993AH, GSH; CAS No.: 70-18-8, oxidized glutathione (GSSG); CAS No.: 27025-41-8, thiobarbituric acid (TBA); CAS No.: 504-17-6, 1,2-dichloro-4-nitrobenzene; CAS No.: 99-54-7, from Sigma (St. Louis, MO, USA). Normal goat serum, Biotinylated, and antibody 2-step plus Poly-HRP Anti Mouse/Rabbit IgG Detection System with

DAB (E-IR-R217) solution were purchased from Elabscience Biotechnology®, China), anti-tumor necrosis factor (TNF- $\alpha$ ) polyclonal antibody (E-AB-63550: 1:500 Dilution) for brain and caspase 3 monoclonal antibody (E-AB-22213: 1:500 Dilution) for the testes.

### **Animals housing and experimental design**

Thirty-six (36) day old cockerel chickens aged ten weeks were distributed into six groups of six chickens each. Group A served as the Control, Group B (Pb; 600 parts per million; ppm), Group C (Pb and 80 mg/kg Naringin), Group D (Pb and 160 mg/kg Naringin), Group E (80 mg/kg Naringin) and Group F (160 mg/kg Naringin). Pb administration was done based on previously reported studies (Wang *et al.*, 2012; Oladapo *et al.*, 2021). The chicks were liberally supplied poultry feed from Top Feeds Limited, Ibadan, and water *ad libitum*. The vehicle used for this work was distilled water for Pb and NAR. Pb and NAR were administered orally. The study was approved with UI-ACUREC/021-0421/16 approval number.

### **Preparation of testes and brain cytosolic fractions**

For the preparation of tissue homogenates, the brain and testes were quickly excised, rinsed, blotted with filter paper, weighed, chopped into bits, and homogenized with homogenizing buffer (0.1 M phosphate buffer, pH 7.4) using a Teflon homogenizer for thirty strokes each. The resulting homogenate was centrifuged at 10,000  $\times g$  for 10 min with a cold centrifuge at  $-4^{\circ}\text{C}$  to obtain post mitochondrial fractions (PMFs). The supernatants (PMFs) were used for biochemical assays.

### **Determination of *in vivo* antioxidant status**

The superoxide dismutase (SOD) assay was carried out by the method of

Oyagbemi *et al.* (2015). Briefly, 100 mg of epinephrine was dissolved in 100 ml distilled water and acidified with 0.5 ml concentrated hydrochloric acid. Thirty microliters of PMF were added to 2.5 ml 0.05 M carbonate buffer (pH 10.2) followed by the addition of 300  $\mu\text{l}$  of 0.3 mM adrenaline. The increase in absorbance at 480 nm was monitored every 30 sec for 150 sec. The one unit of SOD activity was given as the amount of SOD necessary to cause 50% inhibition of the autooxidation of adrenaline to adrenochrome. Beutler *et al.* (1963) study was used to determine glutathione peroxidase (GPx), and glutathione S-transferase (GST) was estimated by the method of Habig *et al.* (1974) and Beutler *et al.* (1963). The reaction mixtures contained 0.5 ml of potassium phosphate buffer (pH 7.4), 0.1 ml of sodium azide 0.2 ml of GSH solution, 0.1 ml of  $\text{H}_2\text{O}_2$ , 0.5 ml of sample and 0.6 ml of distilled water. The mixture was incubated in a water bath at  $37^{\circ}\text{C}$  for 5 min and 0.5 ml of trichloroacetic acid was added and centrifuged at 4,000 rpm for 5 min. A volume of 1 ml of the supernatant was taken and added 2 ml of  $\text{K}_2\text{PHO}_4$  and 1 ml of Ellman's reagent. The absorbance was read at 412 nm using distilled water as blank. The reduced glutathione (GSH) content was estimated by an established method (Salbitani *et al.*, 2017). Here, 0.5 ml of 4% sulfo-salicylic acid (precipitating agent) was added to 0.5 ml of sample and centrifuged at 4,000 rpm for 5 min. To 0.5 ml of the resulting supernatant, 4.5 ml of Ellman's reagent (0.04 g of DTNB in 100 ml of 0.1 M phosphate buffer, pH 7.4) was added. The absorbance was read at 412 nm against distilled water as blank.

### **Determination of oxidative stress indices**

The content of  $\text{H}_2\text{O}_2$  generated was quantified by Wolff's method (Wolff, 1994). To 50  $\mu\text{l}$  of sample, 2.5 ml 0.1M potassium phosphate buffer (pH 7.4), 250  $\mu\text{l}$  of ammonium ferrous sulfate, 100  $\mu\text{l}$  of

## Protection by naringin in lead-exposed chicks

sorbitol, 100 µl of xanthine oxidase and 25 µl of H<sub>2</sub>SO<sub>4</sub> were added and the mixture was thoroughly vortexed until a light pink color of the reaction mixture was observed. The reaction mixture was subsequently incubated at room temperature for 30 min. The optical density was read at absorbance at 560 nm. The H<sub>2</sub>O<sub>2</sub> generated was extrapolated from H<sub>2</sub>O<sub>2</sub> standard curve.

Malondialdehyde (MDA) content was determined following the protocol of Varshney and Kale (1990). Malondialdehyde (MDA), a lipid peroxidation by-product was quantified in the PMFs of neural and testicular tissues according to the method of Varshney and Kale (1990). To 1.6 ml of Tris-KCl, 0.5 ml of 30% TCA, 0.4 ml of sample, and 0.5 ml of 0.75% thiobarbituric acid prepared in 0.2 M HCl were added and mixed well. The reaction mixture was incubated in a water bath at 80°C for 45 min to generate a pink colored malondialdehyde and centrifuged at 4,000 rpm for 15 min. The absorbance was measured against a blank at 532 nm. Lipid peroxidation levels were extrapolated with a molar extinction coefficient of  $1.56 \times 10^5$  M/cm.

### Protein concentration and nitric oxide content

Protein contents were determined by the method of Wilson and Walker (2010). Mixture was incubated at room temperature for 30 min, and values were read with spectrophotometer at 540 nm using distilled water as blank. The final value for total protein was extrapolated from total protein standard curve using Bovine Serum Albumin.

### Histopathology

Testes and brain were fixed in 10% formalin and Bouin's solution, respectively, embedded in paraffin wax, and sections of 5-6 µm in thickness were made and thereafter stained with Haematoxylin and Eosin (H&E) as

previously described by Rotimi et al. (2000).

### Immunohistochemistry

Immunostaining was performed as earlier described for caspase 3 monoclonal antibody in the testis and TNF-α polyclonal antibody in the brain tissues (Oyagbemi et al., 2022). Sections were observed with light microscope (Leica LAS-EZ®) using Leica software application suite version 3.4 equipped with a digital camera.

### Data analysis

One-Way Analysis of Variance (ANOVA) was used to test statistical significance, and the results are expressed as mean ± standard deviation (SD). Tukey post-hoc test was used to test for the difference in the variances among different groups. Significant difference between the means of two groups was compared by the student's t test.

## Results

### Oxidative stress indices and antioxidant status

The results from this study indicated a statistically significant ( $p < 0.05$ ) increase in the activities of testicular GPx, GST, and SOD in chicks intoxicated with Pb compared to the control. The recorded high testicular antioxidant enzymes could be linked to toxicant-induced adaptive response. Co-treatment with NAR however improved testicular antioxidant status (Figure 1). In Figure 2, indices of testicular oxidative stress (H<sub>2</sub>O<sub>2</sub> and MDA) are noticed to increase in Pb-exposed chicks compared to the untreated animals. Also, significant reduction in values of MDA was observed in the NAR co-treated with Pb acetate groups (Figure 2). In another experiment, a noticeable lower activity of GPx in the brain, however, increased activities of GST, and SOD of Pb-intoxicated chicks were observed in comparison to the control.

NAR co-treatment caused observable improvement in the brain antioxidant activities (Figure 3). We observed higher values in the activities of SOD and GST, and this could be associated with adaptive response. Again, Pb administration enhanced the production of markers of oxidative stress together with depletion of

GSH content in comparison to the control and chicks treated with NAR as observed in Figure 4. The metal-chelating and antioxidant power of NAR was demonstrated by reduction in neuronal H<sub>2</sub>O<sub>2</sub> generation and MDA content as well as appreciable improvement in GSH content (Figure 4).




Figure 1. Testicular antioxidant defense status: Superscript (\*) indicates significant difference at  $p < 0.05$  when compared with Group A (Control), while superscript (#) indicates significant difference when compared with Group B (Pb alone). Group A: Control, Group B: Pb (600 ppm), Group C: Pb and Naringin (80 mg/kg), Group D: Pb and Naringin (160 mg/kg), Group E: Naringin (80 mg/kg), and Group F: Naringin (160 mg/kg). Sample size ( $n = 6$ ). Abbreviations: GPx; Glutathione peroxidase, GST; Glutathione S-transferase, SOD; Superoxide dismutase; Pb= lead acetate; Pb+NAR1=lead acetate plus naringin 80 mg/kg; Pb+NAR2= lead acetate plus naringin 160 mg/kg; NAR 1= 80 mg/kg; NAR2= 160 mg/kg.




Figure 2. Testicular markers of oxidative stress: Superscript (\*) indicates significant difference at  $p < 0.05$  when compared with Group A (Control), while superscript (#) indicates significant difference when compared with Group B (Pb alone). Group A: Control, Group B: Pb (600 ppm), Group C: Pb and Naringin (80 mg/kg), Group D: Pb and Naringin (160 mg/kg), Group E: Naringin (80 mg/kg), and Group F: Naringin (160 mg/kg). Sample size ( $n = 6$ ). Abbreviations: H<sub>2</sub>O<sub>2</sub>; hydrogen peroxide, MDA; Malondialdehyde, GSH; Reduced glutathione; Pb= lead acetate; Pb+NAR1=lead acetate plus naringin 80 mg/kg; Pb+NAR2= lead acetate plus naringin 160 mg/kg; NAR 1= 80 mg/kg; NAR2= 160 mg/kg.

## Protection by naringin in lead-exposed chicks

### Histopathology

The histology of the testes revealed Sertoli cells swelling and degeneration of seminiferous tubules in chicks intoxicated with Pb as indicated in Figure 5 compared to the control. Interestingly, cockerel

chicks co-administered with Pb and NAR (80 mg/kg), Pb and NAR (160 mg/kg), showed mild swelling of Sertoli cells and degeneration of seminiferous tubules relative to the chicks intoxicated with Pb (Figure 5).




Figure 3. Brain antioxidant defense status: Superscript (\*) indicates significant difference at  $p < 0.05$  when compared with Group A (Control), while superscript (#) indicates significant difference when compared with Group B (Pb alone). Group A: Control, Group B: Pb (600 ppm), Group C: Pb and Naringin (80 mg/kg), Group D: Pb and Naringin (160 mg/kg), Group E: Naringin (80 mg/kg), and Group F: Naringin (160 mg/kg). Sample size ( $n = 6$ ). Abbreviations: GPx; Glutathione peroxidase, GST; Glutathione S-transferase, SOD; Superoxide dismutase; Pb= lead acetate; Pb+NAR1=lead acetate plus naringin 80 mg/kg; Pb+NAR2= lead acetate plus naringin 160 mg/kg; NAR 1= 80 mg/kg; NAR2= 160 mg/kg.




Figure 4. Brain markers of oxidative stress: Superscript (\*) indicates significant difference at  $p < 0.05$  when compared with Group A (Control), while superscript (#) indicates significant difference when compared with Group B (Pb alone). Group A: Control, Group B: Pb (600 ppm), Group C: Pb and Naringin (80 mg/kg), Group D: Pb and Naringin (160 mg/kg), Group E: Naringin (80 mg/kg), and Group F: Naringin (160 mg/kg). Sample size ( $n = 6$ ). Abbreviations: H<sub>2</sub>O<sub>2</sub>; hydrogen peroxide, MDA; Malondialdehyde, GSH; Reduced glutathione; Pb= lead acetate; Pb+NAR1=lead acetate plus naringin 80 mg/kg; Pb+NAR2= lead acetate plus naringin 160 mg/kg; NAR 1= 80 mg/kg; NAR2= 160 mg/kg.

NAR in a dose-dependently ameliorated anatomical anarchy precipitated Pb toxicity. Again, we discovered that swelling of neurons with loss of arborization and acute necrosis of neurons in brain of Pb exposed chicks (Figure 6). The neuroprotective effect of NAR was demonstrated with no visible lesions in brain tissues of chicks co-administered with NAR (Figure 6).

### Immunohistochemistry

The immuno-localization of testicular caspase 3 and brain TNF- $\alpha$  revealed higher expressions in Pb exposed chicks relative to control and groups co-treated with NAR (Figures 7 and 8). The reduction in the expressions of caspase 3 and TNF- $\alpha$  was indicative of anti-apoptotic, anti-inflammatory, and neuroprotective effects of NAR against Pb acetate-induced testicular cell death and associated neuro-inflammation.




Figure 5. The histopathology of the testes. A (Control), B (Lead acetate; 600 ppm), C (Lead acetate + Naringin 80 mg/kg), D (Lead acetate + Naringin 80 mg/kg), E (Naringin 80 mg/kg), and F (Naringin 100 mg/kg). Slides were stained with Haematoxylin and Eosin. (Magnification x 400).




Figure 6. The histopathology of the brain. A (Control), B (Lead acetate; 600 ppm), C (Lead acetate + Naringin 80 mg/kg), D (Lead acetate + Naringin 80 mg/kg), E (Naringin 80 mg/kg), and F (Naringin 100 mg/kg). Slides were stained with Haemtoxylin and Eosin. (Magnification x 400).



## Protection by naringin in lead-exposed chicks




Figure 7a. The immunohistochemistry of testicular caspase 3. A (Control), B (Lead acetate; 600 ppm), C (Lead acetate + *Naringin* 80 mg/kg), D (Lead acetate + *Naringin* 80 mg/kg), E (*Naringin* 80 mg/kg), and F (*Naringin* 100 mg/kg). Slides were stained with high-definition Haemtoxylin. (Magnification x 100).




Figure 8a. The immunohistochemistry of neuronal Tumor Necrosis Factor alpha (TNF alpha). A (Control), B (Lead acetate; 600 ppm), C (Lead acetate + *Naringin* 80 mg/kg), D (Lead acetate + *Naringin* 80 mg/kg), E (*Naringin* 80 mg/kg), and F (*Naringin* 100 mg/kg). Slides were stained with high-definition Haematoxylin. (Magnification x 100).

## Discussion

Pb poisoning has contributed significantly to public health risk, thereby negatively impacting the avian ecosystem health. Pb is a male reproductive toxicant that causes reproductive dysfunction and infertility (Al-Megrin et al., 2020). The testicular cells are easily subjected to oxidative damage due to their high contents of polyunsaturated fatty acids (PUFA) making testicular tissues prone to lipid peroxidation, oxidative stress, and apoptosis (El-khadragy et al., 2020). The pro-apoptotic gene of caspase-3

overexpression in Pb toxicity has been found to stimulate apoptotic processes (Descalzo et al., 2021). In this study, we proposed that Pb intoxication might activate testicular programmed cell death. Pb toxicity has been shown to enhance testicular cell death by raising caspase-3 expression, with concomitant testicular toxicity (Matuq Al-Yasi et al., 2021). However, from our results, NAR co-administration lowered caspase 3 expression. Pb exposure has also been linked to testicular damage and spermatogenic cell death in previous

investigation (El-Khadragy *et al.*, 2020). Similarly, Pb administration has been documented to cause exaggerated increase in oxidative stress, inflammation, and apoptosis (Yardim *et al.*, 2020). In the present study, we observed induction of oxidative as indicated with exaggerated raise in MDA content, hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) generation, and reduction in reduced glutathione content in testes and brain tissues. Oxidative stress is the imbalance between antioxidant and ROS generation in favor of ROS (Ileriturk *et al.*, 2021). Pb toxicity has been reported to cause oxidative stress (Oyem *et al.*, 2021). Furthermore, oxidative stress induction has been associated with heavy metal toxicity and contamination in poultry as described by Touyz *et al.* (2020). Our findings are consistent with previous reports on oxidative stress, inflammation and apoptosis following Pb intoxication (Wang *et al.*, 2022; Bhattacharya, 2022). The anti-apoptotic and antioxidant property of NAR as recorded in this study could be of a great economic benefit to the global poultry industry. The testicular apoptosis observed in our study was reversed in NAR-treated chicks. Therefore, NAR could be incorporated into poultry feeds to improve reproductive toxicity associated with infertility in breeder poultry farms contaminated with heavy metals such as Pb. Hence, the combination of anti-inflammatory, antioxidant, and metal-chelating properties of NAR could be explored in poultry feeds contaminated with heavy metals.

Pb is one of several neurotoxins that interferes with and diminishes intracellular communication in nerve processes. Neurodegeneration appears to be linked to degenerative alterations and apoptosis in the neural tissue, and this could occur due to ability of Pb to cross the blood brain barrier (Yu *et al.*, 2021). Lead toxicity produces free radicals and causes dysfunctional homeostatic milieu (Gadde and Betharia, 2021). Nerve cells have lower capacity to detoxify ROS, making

them prone to increases in ROS levels, which can be caused by Pb poisoning, thereby precipitating neurological processes (Omobowale *et al.*, 2016). From our study, Pb-treated chicks were seen to have marked oxidative stress levels and depletion in the antioxidant defense systems which corroborates other studies (Zhang *et al.*, 2021). However, NAR administration was able to quench oxidative stress indices thereby restoring the antioxidant defense system as indicated in the improvement in *in vivo* antioxidant machinery (Wang *et al.*, 2019). The improvement in antioxidant capacity of NAR can be linked to its observed anti-oxidative and anti-apoptotic properties. Nitric oxide (NO) concentrations in the brain could be used as an indicator of neuronal nitrosative stress (Rafaiee *et al.*, 2021). Without a doubt, NO-induced oxidative imbalance has been reported to trigger inflammatory processes (Oladapo *et al.*, 2021). It is worth noting that previous research has linked NAR's neuroprotective properties to its ability to suppress iNOS expression. Upregulation of NO in the brain has been correlated with neuro-inflammation and oxidative stress as previously described elsewhere (Ajibade and Ogundero, 2021). From our data, co-administration of Pb with NAR significantly attenuated neuro-inflammation induced by Pb intoxication (Umukoro *et al.*, 2018). This therefore supports the reported documentations on the anti-inflammatory action of NAR (Hadjira *et al.*, 2022). Therefore, the present study is in consonance with anti-inflammatory and neuroprotective actions of NAR (Akintunde *et al.*, 2020). We found that Pb exposure caused a considerable increase in the level of TNF- $\alpha$ , as evidenced by immunohistochemistry. The involvement of TNF- $\alpha$  signaling pathway in neuro-inflammation has been reported (Wang *et al.*, 2020). However, we discovered that NAR reversed Pb toxic effects which corroborates the neuroprotective effects of NAR in

## Protection by naringin in lead-exposed chicks

accordance with other studies (Varshney and Garabadu, 2021). We hypothesized that the anti-inflammatory action of NAR might be through suppression and or mitigation of TNF- $\alpha$  signaling pathway. The link between testicular damage and neuroinflammation as recorded in the present study, might be connected to the hypothalamic-pituitary-gonadal axis. The disruption of this axis by Pb toxicity could be responsible for the brain and testicular damage observed from our findings. Therefore, NAR could open a novel therapeutic window for the management of conditions associated with neuroinflammation precipitated by Pb toxicity.

It is therefore recommended that NAR could be incorporated into poultry feed as a metal-chelating antioxidant, anti-stress, and heavy metal binder as an alternative method of improving poultry health. The major limitation of this study was inadequate funding precipitated by global COVID-19 pandemic. In the current post-pandemic era, we hope to source for more funds to gain deeper insights into the molecular mechanism of NAR against lead toxicity. This enables us to unravel the mechanism of Pb toxicity and its negative impact on hypothalamic-pituitary-gonadal axis and the reproductive system.

### Acknowledgment

The authors deeply thank the African union through the the Pan African University Institute of Earth and Life Sciences Institute (PAULESI) for funding this work and Cardio-renal laboratory, Uuniversity of Ibadan for the facilities used.

### Conflicts of interest

The authors have declared that there is no conflict of interest.

### References

Abdel Moneim AE, Dkhil MA, Al-Quraishy S. 2011. The protective effect of flaxseed oil on lead acetate-induced renal toxicity in

rats. *J Hazard Mater*, 194: 250-255.

Abdel-Emam RA, Ahmed EA. 2021. Ameliorative effect of l-carnitine on chronic lead-induced reproductive toxicity in male rats. *Vet Med Sci*, 7: 1426-1435.

Abiodun EM, Adeleye MM. 2020. Association of copper-to zinc ratio with sperm concentration among males investigated for infertility. *J Infertil Reprod Biol*, 8: 49-52.

Ahmad F, Salahuddin M, Alsamman K, AlMulla AA, Salama KF. 2018. Developmental lead (Pb)-induced deficits in hippocampal protein translation at the synapses are ameliorated by ascorbate supplementation. *Neuropsychiatr Dis Treat*, 14: 3289-3298.

Ajibade AJ, Ogundero SA. 2021. Neurohistological changes in the cerebellar cortex of adult wistar rats following lead acetate induced cortical damage. *GSC Biol Pharm Sci*, 15: 182-188

Akintunde JK, Akintola TE, Adenuga GO, Odugbemi ZA, Adetoye RO, Akintunde OG. 2020. Naringin attenuates bisphenol-A mediated neurotoxicity in hypertensive rats by abrogation of cerebral nucleotide depletion, oxidative damage and neuroinflammation. *Neurotoxicology*, 81: 18-33.

Hosseinzadeh H, Parvardeh S, Nassiri-Asl M, Mansouri MT. 2005b. Intracerebroventricular administration of thymoquinone, the major constituent of *Nigella sativa* seeds, suppresses epileptic seizures in rats. *Med Sci Monit*, 4: 106-110.

Al-Khafaf AI, Ismail HK, Al-Saidya AM. 2021. Histopathological effects of experimental exposure to lead on nervous system in albino female rats. *Iraqi J Vet*, 35: 45-48.

Al-Megrin WA, Alomar S, Alkhuriji AF, Metwally DM, Mohamed SK, Kassab RB, Abdel Moneim AE, El-Khadragy MF. 2020. Luteolin protects against testicular injury induced by lead acetate by activating the nrf2/HO-1 pathway. *IUBMB Life*, 72: 1787-1798.

Alotaibi MF, Al-Joufi F, Abou Seif HS, Alzoghbi MA, Djouhri L, Ahmeda AF, Mahmoud AM. 2020. Umbelliferone inhibits spermatogenic defects and testicular injury in lead-intoxicated rats by suppressing oxidative stress and inflammation, and improving nrf2/HO-1 signaling. *Drug Des Devel Ther*, . 14:

- 4003-4019.
- Alyami NM, Almeer R, Alyami HM. 2022. Protective effects of *asparagus officinalis* (asparagus) against lead toxicity in mice. *Environ Sci Pollut Res Int*, 30: 18718-18730.
- Bentaiba K, Belhocine M, Chougrani F, Bououdina M, Mostari A, Fernini M, Bouzouina M. 2023. Effectiveness of *withania frutescens* root extract on testicular damage induced by lead acetate in adult albino rats. *Reprod Toxicol*, 115: 102-110.
- Beutler E, Duron O, Kelly MB. 1963. Improved method for determination of blood glutathione reduced. *J Lab Clin Med*, 61: 882-888.
- Bhattacharya S. 2022. Essential trace metals as countermeasure for lead toxicity. *J Environ Pathol Toxicol Oncol*, 41: 61-67.
- Bidanchi RM, Lalrindika L, Khushboo M, Bhanushree B, Dinata R, Das M, Nisa N, Lalrinzuali S, Manikandan B, Saeed-Ahmed L, Sanjeev S, Murthy MK, Roy VK, Gurusubramanian G. 2022. Antioxidative, anti-inflammatory and anti-apoptotic action of ellagic acid against lead acetate induced testicular and hepato-renal oxidative damages and pathophysiological changes in male Long Evans rats. *Environ Pollut*, 302: 119048.
- Chen C, Lin B, Qi S, He J, Zheng H. 2019. Protective effects of salidroside on lead acetate-induced oxidative stress and hepatotoxicity in sprague-dawley rats. *Biol Trace Elem Res*, 191: 426-434.
- de Andrade Teles RB, Diniz TC, Costa Pinto TC, de Oliveira Júnior RG, Gama E Silva M, de Lavor ÉM, Fernandes AWC, de Oliveira AP, de Almeida Ribeiro FPR, da Silva AAM, Cavalcante TCF, Quintans Júnior LJ, da Silva Almeida JRG. 2018. Flavonoids as therapeutic agents in alzheimer's and parkinson's diseases: a systematic review of preclinical evidences. *Oxid Med Cell Longev*, 2018: 7043213.
- Descalzo E, Camarero PR, Sánchez-Barbudo IS, Martínez-Haro M, Ortiz-Santaliestra ME, Moreno-Opo R, Mateo R. 2021. Integrating active and passive monitoring to assess sublethal effects and mortality from lead poisoning in birds of prey. *Sci Total Environ*, 750: 142260.
- El-Khadragy MF, Al-Megrin WA, Al-Sadhan NA, Metwally DM, El-Hennamy RE, Salem FEH, Kassab RB, Abdel Moneim AE. 2020. Impact of coenzyme Q10 administration on lead acetate induced testicular damage in rat. *Oxid Med Cell Longev*, 2020: 4981386.
- Ellman GL. 1959. Tissue sulfhydryl groups. *Arch Biochem Biophys*, 82: 70-77.
- Ferreira MC, Zucki F, Duarte JL, Iano FG, Ximenes VF, Buzalaf MA, Oliveira RC. 2017. Influence of iron on modulation of the antioxidant system in rat brains exposed to lead. *Environ Toxicol*, 32: 813-822.
- Gadde R, Betharia SN. 2021. N-bis-(2-mercaptoethyl) isophthalamide (NBMI) exerts neuroprotection against lead-induced toxicity in U-87 MG cells. *Arch Toxicol*, 95: 2643-2657.
- Ghofrani S, Joghataei MT, Mohseni S, Baluchnejadmojarad T, Bagheri M, Khamse S, Roghani M. 2015. Naringenin improves learning and memory in an Alzheimer's disease rat model: Insights into the underlying mechanisms. *Eur J Pharmacol*, 764: 195-201.
- Gindri DB, Peres KC, Scortegagna CM, Moura MR, Mateus HR, Dos-Santos RK, Maciel AP, Matté C. 2021. Naringin supplementation during pregnancy induces sex and region-specific alterations in the offspring's brain redox status. *Int J Environ Res Public Health*, 18: 4805.
- Habig WH, Pabst MJ, Jakpoby WB. 1974. Glutathione S-transferase, a first enzymatic step in mercapturic acid formation. *J Biol Chem*, 249: 7130-7139.
- Hadjira S, Mansour A, Seghiri R, Menad A, Benayache F, Benayache S, Ameddah S. 2022. Neuroinflammation and behavioral deficit in rotenone-induced neurotoxicity in rats and the possible effects of butanolic Extract of *centaurea africana*. *Recent Adv Inflamm Allergy Drug Discov*, 15: 35-43.
- Ibrahim IA, Shalaby AA, Abd Elaziz RT, Bahr HI. 2021. *Chlorella vulgaris* or *spirulina platensis* mitigate lead acetate-induced testicular oxidative stress and apoptosis with regard to androgen receptor expression in rats. *Environ Sci Pollut Res Int*, 28: 39126-39138.
- Ilerturk M, Benzer F, Aksu EH, Yildirim S, Kandemir FM, Dogan T, Dortbudak MB, Genc A. 2021. Chrysin protects against testicular toxicity caused by lead acetate in rats with its antioxidant, anti-inflammatory, and antiapoptotic properties. *J Food*

## Protection by naringin in lead-exposed chicks

- Biochem, 45: 13593.
- Kahalerras L, Otmani I, Abdennour C. 2022. Wild Garlic *Allium triquetrum* L. Alleviates lead acetate-induced testicular injuries in rats. *Biol Trace Elem Res*, 200: 2205-2222.
- Khodabandeh Z, Dolati P, Zamiri MJ, Mehrabani D, Bordbar H, Alae S, Jamhiri I, Azarpira N. 2021. Protective effect of quercetin on testis structure and apoptosis against lead acetate toxicity: a stereological study. *Biol Trace Elem Res*, 199: 3371-3381.
- Kumar RS, Devi SA. 2018. Lead toxicity on male reproductive system and its mechanism: a review. *Res J Pharm Technol*, 11: 1228.
- Matuq Al-Yasi H, El-Shazly SA, Ahmed EF, Alamer KH, Hessini KY, Attia HA, Alkafafy ME, Mohamed AA, Hassan FA. 2021. Protective effects of taif rosewater against testicular impairment induced by lead intoxication in rats. *Andrologia*, 53: 14045.
- Mitra S, Patra T, Saha D, Ghosh P, Mustafi SM, Varghese AC, Murmu N. 2022. Sub-chronic cadmium and lead compound exposure induces reproductive toxicity and development of testicular germ cell neoplasia in situ in murine model: attenuative effects of resveratrol. *J Biochem Mol Toxicol*, 36: 23058.
- Ogundele MO. 2018. Behavioural and emotional disorders in childhood: A brief overview for paediatricians. *World J Clin Pediatr*, 7: 9-26.
- Oladapo OM, Ben-Azu B, Ajayi AM, Emokpae O, Eneni AO, Omogbiya IA, Iwalewa EO. 2021. Naringin confers protection against psychosocial defeat stress-induced neurobehavioral deficits in mice: involvement of glutamic acid decarboxylase isoform-67, oxido-nitric stress, and neuroinflammatory mechanisms. *J Mol Neurosci*, 71: 431-445.
- Olaleye SB, Adaramoye OA, Erigbali PP, Adeniyi OS. 2007. Lead exposure increases oxidative stress in gastric mucosa of HCL/ethanol exposed rat. *World J Gastroenterol*, 13: 5121-5126.
- Olaniyan OT, Ojewale OA, Dare A, Adebayo O, Enyojo JE, Okotie GE, Adetunji CO, Mada BS, Okoli BJ, Eweoya OO. 2021. *Cocos nucifera* L. oil alleviates lead acetate-induced reproductive toxicity in sexually-matured male Wistar rats. *J Basic Clin Physiol Pharmacol*, 33: 297-303.
- Omobowale TO, Oyagbemi AA, Akinleye SA, Ola-Davies OE, Adebowale BS, Olopade JO, Adedapo AA. 2016. Effect of exposure and withdrawal on Lead-induced toxicity and oxidative stress in cardiac tissues of rats. *Toxicol Int*, 23: 12-17.
- Oyagbemi AA, Omobowale TO, Akinrinde AS, Saba AB, Ogunpolu BS, Daramola O. 2015. Lack of reversal of oxidative damage in renal tissues of lead acetate-treated rats. *Environ Toxicol*, 30: 1235-1243.
- Oyagbemi AA, Adejumbi OA, Jarikre TA, Ajani OS, Asenuga ER, Gbadamosi IT, Adedapo ADA, Aro AO, Ogunpolu BS, Hassan FO, Falayi OO, Ogunmiluyi IO, Omobowale TO, Arojoye OA, Ola-Davies OE, Saba AB, Adedapo AA, Emikpe BO, Oyeyemi MO, Nkadimeng SM, McGaw LJ, Kayoka-Kabongo PN, Oguntibeju OO, Yakubu MA. 2022. Clofibrate, a Peroxisome proliferator-activated receptor-alpha (PPAR $\alpha$ ) agonist, and its molecular mechanisms of action against sodium fluoride-induced toxicity. *Biol Trace Elem Res*, 200: 1220-1236.
- Oyagbemi AA, Omobowale TO, Awoyomi OV, Ajibade TO, Falayi OO, Ogunpolu BS, Okotie UJ, Asenuga ER, Adejumbi OA, Hassan FO, Ola-Davies OE, Saba AB, Adedapo AA, Yakubu MA. 2019. Cobalt chloride toxicity elicited hypertension and cardiac complication via induction of oxidative stress and upregulation of COX-2/Bax signaling pathway. *Hum Exp Toxicol*, 38: 519-532.
- Oyem JC, Chris-Ozoko LE, Enaohwo MT, Otabor FO, Okudayo VA, Udi OA. 2021. Antioxidative properties of *ocimum gratissimum* alters lead acetate induced oxidative damage in lymphoid tissues and hematological parameters of adult Wistar rats. *Toxicol Rep*, 10: 215-222.
- Qin S, Chen Q, Wu H, Liu C, Hu J, Zhang D, Xu C. 2016. Effects of naringin on learning and memory dysfunction induced by gp120 in rats. *Brain Res Bull*, 124: 164-171.
- Rafaiee R, Khastar H, Garmabi B, Taleb M, Norouzi P, Khaksari M. 2021. Hydrogen sulfide protects hippocampal CA1 neurons against lead mediated neuronal damage via reduction oxidative stress in male rats. *J Chem Neuroanat*, 112: 101917.
- Rotimi O, Cairns A, Gray S, Moayyedi P,

- Dixon MF. 2000. Histological identification of helicobacter pylori: comparison of staining methods. *J Clin Pathol*, 53: 756-759.
- Salbitani G, Bottone C, Carfagna S. 2017. Determination of reduced and total glutathione content in extremophilic microalga *galdieria phlegrea*. *Bio Protoc*, 7: 2372.
- Touyz RM, Rios FJ, Alves-Lopes R, Neves KB, Camargo LL, Montezano AC. 2020. Oxidative stress: a unifying paradigm in hypertension. *Can J Cardiol*, 36: 659-670.
- Umukoro S, Kalejaye HA, Ben-Azu B, Ajayi AM. 2018. Naringenin attenuates behavioral derangements induced by social defeat stress in mice via inhibition of acetylcholinesterase activity, oxidative stress and release of pro-inflammatory cytokines. *Biomed Pharmacother*, 105: 714-723.
- Varshney R, Kale RK. 1990. Effect of calmodulin antagonists on radiation induced lipid peroxidation in microsomes. *Int J Radiat. Biol*, 58: 733-743.
- Varshney V, Garabadu D. 2021. Naringin exhibits mas receptor-mediated neuroprotection against amyloid beta-induced cognitive deficits and mitochondrial toxicity in rat brain. *Neurotox Res*, 39: 1023-1043.
- Wang J, Yang Z, Lin L, Zhao Z, Liu Z, Liu X. 2012. Protective effect of naringenin against lead-induced oxidative stress in rats. *Biol Trace Elem Res*, 146: 354-359.
- Wang L, Zhang Z, Wang H. 2021. Naringin attenuates cerebral ischemia-reperfusion injury in rats by inhibiting endoplasmic reticulum stress. *Transl Neurosci*, 12: 190-197.
- Wang W, Zhao Y, Shi F, Li S, Liu Z, Wang W, Zhu Y, Zhang Y. 2022. Role of ferroptosis in cerebellar injury of mice following lead exposure. *Wei Sheng Yan J*, 51: 443-469.
- Wang Z, Wang S, Zhao J, Yu C, Hu Y, Tu Y, Yang Z, Zheng J, Wang Y, Gao Y. 2019. Naringenin ameliorates renovascular hypertensive renal damage by normalizing the balance of renin-angiotensin system components in rats. *Int J Med Sci*, 16: 644-653.
- Wilson K, Walker J. 2010. Principles and techniques of practical biochemistry. Cambridge University Press; Cambridge, UK: 2010.
- Wolff SF. 1994. Ferrous ion oxidation in the presence of ferric ion indicator xylenol orange for measurement of hydrogen peroxides. *Methods Enzymol*, 233: 182-189.
- Xia Y, Zweiler JL. 1997. Measurement of myeloperoxidase in leucocyte containing tissue. *Anal Biochem*, 245: 93-96.
- Yardim A, Kandemir FM, Ozdemir S, Kucukler S, Comakli S, Gur C, Celik H. 2020. Quercetin provides protection against the peripheral nerve damage caused by vincristine in rats by suppressing caspase 3, NF- $\kappa$ B, ATF-6 pathways and activating Nrf2, Akt pathways. *Neurotoxicology*, 81: 137-146.
- Yousef AOS, Fahad AA, Abdel Moneim AE, Metwally DM, El-Khadragy MF, Kassab RB. 2019. The neuroprotective role of coenzyme Q10 against lead acetate-induced neurotoxicity is mediated by antioxidant, anti-inflammatory and anti-apoptotic activities. *Int J Environ Res Public Health*, 16: 2895.
- Yu C, Pan S, Zhang J, Li X, Niu Y. 2021. Ferulic acid exerts Nrf2-dependent protection against prenatal lead exposure-induced cognitive impairment in offspring mice. *J Nutr Biochem*, 91: 108603.