

Short-Communication

Carvacrol alleviates the proconvulsive effects of lipopolysaccharide (LPS) and reduces the gene expression of proinflammatory cytokines *interleukin-1* and *tumor necrosis factor- α*

Hesamodin Bagheripoor¹, Mahdiah Mondanizadeh², Mehdi Sadegh^{3,*}

¹Faculty of Medicine, Arak University of Medical Sciences, Arak, Iran

²Department of Biotechnology and Molecular Medicine, Faculty of Medicine, Arak University of Medical Sciences, Arak, Iran

³Department of Physiology, Faculty of Medicine, Arak University of Medical Sciences, Arak, Iran

Article history:

Received: Feb04, 2024

Received in revised form:

Jun12, 2024

Accepted: Jun12, 2024

Epub ahead of print

*** Corresponding Author:**

Tel: +86+34173521-351

Fax: +86+34173520

m.sadegh@arakmu.ac.ir

Keywords:

Inflammation

Interleukin 6 (IL-6)

Interleukin 4 (IL-4)

Seizure

Abstract

Objective: Carvacrol has anti-inflammatory effects. According to the links between inflammatory processes and seizures, this study was designed to investigate the potential effect of carvacrol in reducing seizure severity and involvement of hippocampal pro- and anti-inflammatory cytokines.

Materials and Methods: This research was conducted on 42 adult male Wistar rats. Animals were randomly divided into seven groups (n=6). Seizures were induced by PTZ (Pentylenetetrazol) injection (80 mg/kg). LPS (lipopolysaccharide) was injected (400 μ g/kg) 4 hr before PTZ. Carvacrol (100 mg/kg) was injected immediately after LPS. All injections were intraperitoneal (i.p.). Experimental groups were as follows: 1. Control (Cnt) 2. Carvacrol (Cav); 3. LPS; 4. PTZ, 5. PTZ+LPS; 6. PTZ+Cav; 7. PTZ+LPS+Cav. Seizures were observed for 30 min after the PTZ injection and the occurrence of behavioral stages of seizures was recorded. Following the behavioral study, hippocampal samples were collected for gene expression evaluation using the Real Time-PCR technique to assess IL (interleukin)-1, IL-6, IL-4 and TNF (tumor necrosis factor)- α gene expression.

Results: The current study showed that receiving LPS exacerbated seizures in the studied groups. Carvacrol reduced the severity of seizures in the LPS-receiving groups. In the gene expression study, receiving LPS increased the expression of cytokines TNF- α and IL-1 in the hippocampal tissue. Carvacrol significantly decreased gene expression of TNF- α and IL-1. No significant changes were detected for IL-6, IL-4 gene expression.

Conclusion: There could be a relationship between carvacrol ability to modulate the proconvulsive effects of LPS and its ability to decrease the gene expression of inflammatory cytokines.

Please cite this paper as:

BagheripoorH, MondanizadehM, Sadegh M. Carvacrol alleviates the proconvulsive effects of lipopolysaccharide (LPS) and reduces the gene expression of proinflammatory cytokines interleukin-1 and tumor necrosis factor- α . Avicenna J Phytomed, 2025. Epub ahead of print.

Introduction

Epilepsy is a prevalent neurological disease described by recurrent seizures. It is estimated that 50 million people worldwide have epilepsy (Fiest et al. 2017). It is reported that 30-35% of epilepsy cases do not respond adequately to current anti-epileptic drugs (Loscher et al. 2020). By examining samples obtained from the brains of epilepsy patients, it appears that changes in hippocampal neurons cause the loss of balance between the excitation and inhibition in the limbic system leading to subsequent spontaneous seizures (Hoogland et al. 2000). Traumatic or non-traumatic brain injuries such as seizures trigger biological chains that release cytokines in the space around neurons (Vezzani et al. 2013).

Previous reports suggest that inflammation and immune-related responses such as interleukins in brain tissue could increase the risk of convulsions, making these factors potential biomarkers of seizure susceptibility (Gorter et al. 2006). Cytokines are inflammation-related proteins produced and released in the early stages by glial cells in the neuronal environment, and they can affect neuronal excitability and other glial cell functions (Shimada et al. 2014).

In clinical studies, seizures caused by fever have shown an increase in cytokines interleukin-1 (IL-1), interleukin-6 (IL-6), and tumor necrosis factor- α (TNF- α) in the cerebrospinal fluid (Dinarello 2004). TNF- α is an inflammatory cytokine which is released by activated microglia and causes an increase in glutamate neurotransmission and neuronal excitation (Stellwagen and Malenka 2006). IL-1 is expressed by microglia and causes a reduction in glutamate uptake. The increase in this cytokine after seizure causes dysfunction of neurons and decreases their regeneration capacity (Alyu and Dikmen 2017). IL-6 is usually detectable in small amounts in the nervous system, but the stimulation of astrocytes and microglia cells could increase its production. Studies on the

effect of this cytokine on convulsions are conflicting (Grul 2015). IL-4 is an anti-inflammatory cytokine with a proven effect in reducing the activity of various parts of the immune system by inhibiting the production of pro-inflammatory cytokines (Chen et al. 2020).

Carvacrol (C₁₀H₁₄O) is abundantly found in the *Lamiaceae* family, including thymus, oregano, and savory, which are commonly used as food seasonings. Studies have shown that phenolic plant compounds like carvacrol have anti-inflammatory properties and can reduce pro-inflammatory cytokines (Sadegh and Sakhaie 2018).

In this study, our goal was to investigate if carvacrol is able to alleviate proconvulsive effects of lipopolysaccharide (LPS). To understand the cellular pathways and genes involved in these effects, we investigated the effects of carvacrol on gene expression of proinflammatory cytokines IL-1, IL-4, IL-6 and TNF- α .

Materials and Methods

Animals

Forty-two adult male Wistar rats (180–200 g) were used for this research. All rats were placed in standard cages and standard conditions (temperature of 25°C \pm 12 hr light/12 hr darkness). Food and water were freely accessible. All provision from “Guide for the Care and Use of Laboratory Animals” (8th edition; National Academies Press; 2011) was considered. All efforts were made to reduce the number of animals and minimize their suffering.

Chemicals and seizure induction

Carvacrol, Pentylentetrazol and LPS (obtained from Gram-negative *Escherichia coli* serotype O111:B4) all were from Sigma Aldrich (USA). Carvacrol (100 mg/kg) was dissolved in the DMSO (dimethyl sulfoxide) 1% and was injected intraperitoneally (i.p.) immediately after LPS. According to our previous works DMSO 1% had no effect on seizure

parameters. LPS (400 µg/kg) and PTZ (80 mg/kg) were dissolved in saline (0.9%) and injected i.p. (Akarsu et al. 2006; Zareie et al. 2018).

Behavioral seizure scales were monitored for 30 min following the injection of PTZ. For this, each rat was placed into a Plexiglass made box (40*40*40 cm) immediately after receiving the injections.

Seizure stages were scaled as follows: stage 0, normal behavior; stage 1, sniffing and/or freezing; stage 2, head nodding, clonus in head muscles and forelimbs; stage 3, continuous myoclonic jerk, tail rigidity; stage 4, bilateral clonus in forelimbs and jumping; stage 5, tonic-clonic convulsions, rearing and falling. By observing and recording time using a digital timer, the duration of reaching each stage, the duration of stages 3 to 5, the number of occurrences of each stage and the death rate in animals were measured and recorded for additional analysis.

Experimental groups

The studied rats were divided into seven groups of 6 as follows:

1. Control (Cnt)
2. Carvacrol (Cav), receive only carvacrol;
3. LPS, receive only LPS;
4. PTZ, receive only PTZ,
5. PTZ+LPS, PTZ was injected four hours after receiving LPS;
6. PTZ+Cav, PTZ was injected four hours after carvacrol;
7. PTZ+LPS+Cav, PTZ was injected four hours after carvacrol and LPS.

Seizure stages were measured in those PTZ-treated groups. After conducting behavioral seizure tests, animals were sampled for real-time PCR test. First, animal head was quickly removed under chloroform anesthesia using a guillotine. To induce anesthesia, animal were placed in a see-through box, containing chloroform. Then, the skull was opened along the midline and the brain was extracted. On an ice-cooled cutting board, the left and right hippocampus was dissected from hemispheres, snap frozen in liquid nitrogen

and stored at -70°C for extraction of DNA and RNA.

RNA isolation, DNA digestion, and reverse transcription

RNA was extracted from the hippocampus (both left and right hippocampus were mixed) using RNX-plus kit of Cincalor Company and the work steps were carried out according to the manufacturer's instructions. Spectrophotometric optical density measurement (260 and 280 nm) was used to determine RNA total concentration. For each sample tested, the ratio the spectrophotometric readings ratio at 260 nm and 280 nm (OD₂₆₀/OD₂₈₀) was used as estimation for nucleic acid purity. The ratios for all samples were between 1.7 and 2.0.

Oligo(dT) primers were used for reverse transcriptase reactions. Each reaction tube contained 50 ng/µg of total RNA, 11 µl of cDNA Synthesis MIX, 0.5 µmol/L of random hexamers and diethyl dicarbonate (DEPC)-treated water to volume. A semi-quantitative reverse transcriptase polymerase chain reaction (RT-PCR) was performed to determine the levels of *TNF-α*, *IL-4*, *IL-6*, *IL-1* mRNA expressions. A DNA Thermal Cycler 480 (Perkin Elmer, Branchburg, NJ, U.S.A.) was used for reverse transcriptase reactions. Reactions were completed at 42°C for 20 min and 99°C for 5 min. The final cDNA was transferred to -20°C .

The sequence of *TNF-α*, *IL-4*, *IL-6*, *IL-1* genes was extracted from the NCBI database. The primers used to measure the expression of the mentioned genes were designed by Allele ID software and after performing BLAST in the NCBI database was synthesized by Cincalor.

Statistical analysis

Data were analyzed with GraphPad Prism 5.0. One-way ANOVA and Tukey post-test were used to check the difference among the groups. Chi-square test was used for mortality rate analysis and a $p < 0.05$

was considered significant. All data are presented as a mean ± SD.

Results

Seizure data

Latency to each seizure stage was analyzed among the groups by one-way ANOVA. As demonstrated in Table 1 latencies to stages 3, 4 and 5 in LPS+PTZ were significantly lower than PTZ (stage 3 $p < 0.01$; stages 4 and 5 $p < 0.05$), which means that LPS+PTZ group was more susceptible for seizure compared to the PTZ group. However, carvacrol injection in the LPS+Cav+PTZ group significantly reduced this effect of LPS and significantly increased latencies to stages 3, 4 and 5 compared with LPS+PTZ group (in both stages 3 and 5 $p < 0.01$; in stage 4 $p < 0.05$). In addition, carvacrol injection in the PTZ group provided a significant increase in stage 3 latency ($p < 0.05$) but had no significant effect on other stages. So, it seems that carvacrol was more effective on LPS proconvulsive effect than on PTZ-induced seizures.

Moreover, chi-square analysis for mortality rate following the tonic-clonic convulsion on stage 5, showed a higher death rate in the LPS+PTZ group (3 rats died out of 6 rats) than the PTZ group (1 rat died out of 6 rats) ($p < 0.01$). In other groups, no death happened after the stage 5 (Table 1).

As no significant effect was appeared on latencies to stages 1 and 2, we did not report these data in Table 1.

Gene expression data

As demonstrated in Figure 1, hippocampus gene expression of pro-inflammatory cytokines *TNF-α* (A) and *IL-1* (B) was significantly increased in the LPS and PTZ+LPS groups compared to the control (Cnt) (for *TNF-α* $p < 0.05$ for the LPS group and $p < 0.01$ for the PTZ+LPS group compared to Cnt; for *IL-1* $p < 0.05$ for both LPS and LPS+PTZ groups compared to Cnt), while both *TNF-α* and *IL-1* gene expression significantly decreased in the carvacrol (Cav) group compared to the control ($p < 0.05$ for both genes). In addition, gene expression of these pro-inflammatory cytokines in the PTZ+LPS+Cav group significantly decreased compared to the PTZ+LPS group ($p < 0.01$ for both genes), which means that carvacrol eliminates increasing effect of LPS on gene expression of these two proinflammatory cytokines.

Moreover, anti-inflammatory cytokines *IL-4* and *IL-6* gene expression (Figure 1-C, D) revealed no significant changes in compare with LPS or carvacrol in the Cav, LPS and PTZ+LPS groups compared to the control, although *IL-6* expression in the PTZ+LPS+Cav group was significantly increased compared to the PTZ+LPS group ($p < 0.05$).

Table 1. Latencies to seizure stages 3-5.

Groups	S3L(s)	S4L(s)	S5L(s)	Dead out of 6 rats
PTZ	219±51	404±80	542±82	1
LPS+PTZ	134±30 **	256±64 *	343±88 *	3**
Cav+PTZ	277±56 *	500±144	664±199	0
LPS+Cav+PTZ	208±34 ##	419±133 #	550±107 ##	0

LPS significantly decreased latency (s) to seizure stages 3-5, while carvacrol prevented this effect of LPS. Number of death after stage 5 is demonstrated. Data as Mean±SD; * $p < 0.05$ and ** $p < 0.01$ compared to PTZ; # $p < 0.05$ and ## $p < 0.01$ compared with LPS+PTZ. S3L: latency to stage 3, S4L: latency to stage 4, S5L: latency to stage 5, PTZ: Pentylentetrazol, LPS: Lippolysaccharide, Cav: Carvacrol.




Figure 1. Carvacrol eliminated LPS increasing effect on *TNF- α* and *IL-1* gene expression. *TNF- α* (A) and *IL-1* (B) were significantly increased in the LPS and PTZ+LPS groups compare to the control (Cnt), while significantly decreased in PTZ+LPS+Cav group compare to PTZ+LPS. Data is presented as Mean \pm SD; n=6/groups; *p<0.05 and **p<0.01 compared with Cnt, ##p<0.01 compared with PTZ+LPS.

Discussion

In our study, results demonstrated carvacrol efficiency to reduce LPS proconvulsive effects in the tonic-clonic seizure induction model induced by PTZ. In conditions without LPS, carvacrol also increased the latencies to stages 2–3 of seizure which are similar with focal seizures. Gene expression analysis revealed that the anticonvulsant effect of carvacrol could be accompanied with a decrease in cytokines gene expression in the nervous system. Previous studies have shown that LPS injection facilitates PTZ-induced seizures (Sayyah et al. 2003). The proconvulsant effect of LPS begins approximately 30 min after administration and can last up to 12 hr. Seizure susceptibility increases 4 hr after LPS injection with a dose of 400 μ g/kg in rats (Akarsu et al. 2006). Based on these findings, we also administered LPS 4 hr before PTZ at a dose of 400 μ g/kg.

Consistent with earlier reports, our results indicated that LPS exacerbates acute seizure indexes and acts as a proconvulsant agent. Xiang et al. demonstrated that LPS can activate the toll like receptor-4 (TLR4) signaling pathway, leading to excessive release of inflammatory cytokines (Xiang et al. 2016). Although systemic LPS cannot directly reach the brain, Ho et al. reported that proinflammatory cytokines induced by LPS can cross the blood brain barrier and increase susceptibility to seizures (Ho et al. 2015). In our study, a significant increase in cytokines IL-1, IL-4, IL-6, and TNF- α was observed after LPS administration in the hippocampus.

In seizure models induced by PTZ, an increase in the levels of IL-1 β , IL-6, and TNF- α cytokines in the hippocampus of rats has been reported (Singh et al. 2019). Our results also indicated an increase in the expression of cytokines IL-1, IL-4, IL-6, and TNF- α in the group receiving PTZ compared to the control group. Given the chronic nature of epilepsy, there is

sufficient time for inflammatory reactions to activate. Neuroinflammation, by affecting nerve excitability, can lead to the facilitation of convulsions (Dey et al. 2016; Vezzani et al. 2011). Fiala et al. reported that serum levels of IL-1, TNF- α , and IL-6 were elevated in patients with temporal lobe epilepsy and refractory epilepsy (Fiala et al. 2013). Similarly, levels of cytokines IL-1 α , TNF- α , and IL-6 increased in neurons and glial cells of the nervous system before the onset of seizures (Frantz et al. 2017).

Ho et al. suggested that using anti-inflammatories could be a therapeutic approach to reduce seizures provoked by environmental inflammatory factors (Ho et al. 2015). Carvacrol is a herbal compound with confirmed anti-inflammatory effects in various studies. It has been reported that carvacrol's neuroprotection is primarily related to its antioxidant and anti-inflammatory effects (Azizi et al. 2022). Mahmoodi et al. investigated the effect of carvacrol on white brain tissue in experimental autoimmune encephalitis (EAE) mice and observed that carvacrol modulates the expression of pro-inflammatory and anti-inflammatory cytokines in the CNS (Mahmoodi et al. 2019). Carvacrol has been shown to reduce the production of prostaglandins by suppressing cyclo-oxygenase-2 (Sadegh and Sakhaie 2018). Consistent with our data, previous reports have shown the protective function of carvacrol against the inflammatory and brain damaging effects of LPS. In these studies, carvacrol increased antioxidant factors like malondialdehyde and anti-inflammatory factors like TNF- α and IL-6 (Hakimi et al. 2020; Salmani et al. 2022). Additionally, systemic inflammation and oxidative effects of paraquat were reduced by carvacrol administration in a dose similar to our study, with involvement of MDA, IL-6, and Interferon- γ (Amin et al. 2020a; Amin et al. 2020b).

Furthermore, the anticonvulsant effects of carvacrol have been investigated in

different animal models of seizure. In the study by Khalil et al., carvacrol at a dose of 70 mg/kg suppressed seizures in an electroconvulsive model. They concluded that carvacrol prevented the occurrence of status convulsions due to its antagonistic effect on the transient receptor potential melastatin7 (TRPM7) channels (Khalil et al. 2017). Another study also demonstrated the anticonvulsant effect of carvacrol in a maximal electroshock model (Mishra and Baker 2014). In our PTZ-induced convulsion model, the proconvulsive effects of LPS disappeared after carvacrol administration. In biochemical evaluations, carvacrol decreased the cytokines IL-1 and TNF- α and increased the expression of IL-6 and IL-4 in the LPS group compared to the control group. Carvacrol also produced a similar effect in the PTZ+carvacrol group compared to the PTZ group. Interestingly, results from a clinical trial study showed that daily oral administration of carvacrol (1-2 mg/kg) for one month had no negative effect on physiological indexes of the respiratory and blood system (Ghorani et al. 2019).

The exacerbation of inflammation in the nervous system after LPS and PTZ administration can explain the increase in seizures in the LPS+PTZ group compared to the PTZ group. Although IL-1 and TNF- α are recognized as inflammatory cytokines, there is no definite consensus on the inflammatory or anti-inflammatory effect of IL-4 and 6. In our study, it appears that cytokines IL-1 and TNF- α act as inflammatory cytokines, while IL-4 and IL-6 suppressed inflammation and acted as anti-inflammatory cytokines. Therefore, it can be concluded that carvacrol reduces seizures by increasing anti-inflammatory cytokines IL-4 and IL-6 and suppressing inflammatory cytokines TNF- α and IL-1.

It appears that LPS exacerbates seizures by increasing the expression of inflammatory cytokines in the central nervous system. Our study demonstrated that carvacrol decreases the expression of inflammatory factors IL-1 and TNF- α and

increases the expression of cytokines IL-4 and IL-6 in the central nervous system. This is one potential pathway through which carvacrol suppresses the pro-convulsing effect of LPS. Therefore, carvacrol could be used as an anti-inflammatory adjunct to existing anti-epileptic treatments to improve seizure control. However, further research and experiments are necessary in this field.

Conflicts of interest

No conflict of interest for author's.

Funding

The manuscript was extracted from H. Bagheripoor medical thesis, funded by Arak University of Medical Sciences (No. 6604).

Ethical Considerations

All issues verified by the Review Board and Ethics Committee of Arak University of Medical Sciences.

Code of Ethics

IR.ARAKMU.REC.1401.010

Authors' Contributions

All authors had equal contribution in this research study.

References

- Akarsu ES, Ozdayi S, Algan E, Ulupinar F (2006) The neuronal excitability time-dependently changes after lipopolysaccharide administration in mice: possible role of cyclooxygenase-2 induction. *Epilepsy Res* 71(2-3):181-7 doi:10.1016/j.eplepsyres.2006.06.009
- Alyu F, Dikmen M (2017) Inflammatory aspects of epileptogenesis: contribution of molecular inflammatory mechanisms. *Acta Neuropsychiatr* 29(1):1-16 doi:10.1017/neu.2016.47
- Amin F, Memarzia A, Kazerani HR, Boskabady MH (2020a) Carvacrol and Zataria multiflora influenced the PPARgamma agonist effects on systemic inflammation and oxidative stress induced by inhaled paraquat in rat. *Iran J Basic Med Sci* 23(7):930-936 doi:10.22038/ijbms.2020.45962.10648
- Amin F, Roohbakhsh A, Memarzia A, Kazerani HR, Boskabady MH (2020b) Paraquat-induced systemic inflammation and increased oxidative markers in rats improved by Zataria multiflora extract and carvacrol. *Avicenna J Phytomed* 10(5):513-522 doi:AJP-10-513 [pii]
- Azizi Z, Majlessi N, Choopani S, Naghdi N (2022) Neuroprotective effects of carvacrol against Alzheimer's disease and other neurodegenerative diseases: A review. *Avicenna J Phytomed* 12(4):371-387 doi:10.22038/AJP.2022.19491
- Chen L, Zhu L, Lu D, et al. (2020) Interleukin 4 Affects Epilepsy by Regulating Glial Cells: Potential and Possible Mechanism. *Front Mol Neurosci* 13:554547 doi:10.3389/fnmol.2020.554547
- Dey A, Kang X, Qiu J, Du Y, Jiang J (2016) Anti-Inflammatory Small Molecules To Treat Seizures and Epilepsy: From Bench to Bedside. *Trends Pharmacol Sci* 37(6):463-484 doi:10.1016/j.tips.2016.03.001
- Dinarello CA (2004) Infection, fever, and exogenous and endogenous pyrogens: some concepts have changed. *J Endotoxin Res* 10(4):201-22 doi:10.1179/096805104225006129
- Fiala M, Avagyan H, Merino JJ, et al. (2013) Chemotactic and mitogenic stimuli of neuronal apoptosis in patients with medically intractable temporal lobe epilepsy. *Pathophysiology* 20(1):59-69 doi:10.1016/j.pathophys.2012.02.003
- Fiest KM, Sauro KM, Wiebe S, et al. (2017) Prevalence and incidence of epilepsy: A systematic review and meta-analysis of international studies. *Neurology* 88(3):296-303 doi:10.1212/WNL.0000000000003509
- Frantz AL, Regner GG, Pfluger P, et al. (2017) Manual acupuncture improves parameters associated with oxidative stress and inflammation in PTZ-induced kindling in mice. *Neurosci Lett* 661:33-40 doi:10.1016/j.neulet.2017.09.044
- Ghorani V, Boskabady M, Boskabady MH (2019) Effect of carvacrol on pulmonary function tests, and total and differential white blood cell counts in healthy volunteers: A randomized clinical trial. *Avicenna J Phytomed* 9(2):134-142 doi:AJP-9-134 [pii]
- Gorter JA, van Vliet EA, Aronica E, et al.

- (2006) Potential new antiepileptogenic targets indicated by microarray analysis in a rat model for temporal lobe epilepsy. *J Neurosci* 26(43):11083-110
doi:10.1523/JNEUROSCI.2766-06.2006
- Gruol DL (2015) IL-6 regulation of synaptic function in the CNS. *Neuropharmacology* 96(Pt A):42-54
doi:10.1016/j.neuropharm.2014.10.023
- Hakimi Z, Salmani H, Marefati N, et al. (2020) Protective Effects of Carvacrol on Brain Tissue Inflammation and Oxidative Stress as well as Learning and Memory in Lipopolysaccharide-Challenged Rats. *Neurotox Res* 37(4):965-976
doi:10.1007/s12640-019-00144-5 [pii] 10.1007/s12640-019-00144-5
- Ho YH, Lin YT, Wu CW, Chao YM, Chang AY, Chan JY (2015) Peripheral inflammation increases seizure susceptibility via the induction of neuroinflammation and oxidative stress in the hippocampus. *J Biomed Sci* 22(1):46
doi:10.1186/s12929-015-0157-8
- Hoogland G, Hens Jj Fau - De Wit M, De Wit M Fau - van Veelen CW, et al. (2000) Glutamate and gamma-aminobutyric acid content and release of synaptosomes from temporal lobe epilepsy patients. (0360-4012 (Print))
- Khalil A, Kovac S, Morris G, Walker MC (2017) Carvacrol after status epilepticus (SE) prevents recurrent SE, early seizures, cell death, and cognitive decline. *Epilepsia* 58(2):263-273
doi:10.1111/epi.13645
- Loscher W, Potschka H, Sisodiya SM, Vezzani A (2020) Drug Resistance in Epilepsy: Clinical Impact, Potential Mechanisms, and New Innovative Treatment Options. *Pharmacol Rev* 72(3):606-638
doi:10.1124/pr.120.019539
- Mahmoodi M, Amiri H, Ayoobi F, et al. (2019) Carvacrol ameliorates experimental autoimmune encephalomyelitis through modulating pro- and anti-inflammatory cytokines. *Life Sci* 219:257-263
doi:10.1016/j.lfs.2018.11.051
- Mishra RK, Baker MT (2014) Seizure prevention by the naturally occurring phenols, carvacrol and thymol in a partial seizure-psychomotor model. *Bioorg Med Chem Lett* 24(23):5446-9
doi:10.1016/j.bmcl.2014.10.028
- Sadegh M, Sakhaie MH (2018) Carvacrol mitigates proconvulsive effects of lipopolysaccharide, possibly through the hippocampal cyclooxygenase-2 inhibition. *Metab Brain Dis* 33(6):2045-2050
doi:10.1007/s11011-018-0314-3
- Salmani H, Hakimi Z, Arab Z, et al. (2022) Carvacrol attenuated neuroinflammation, oxidative stress and depression and anxiety like behaviors in lipopolysaccharide-challenged rats. *Avicenna J Phytomed* 12(5):514-526
doi:10.22038/AJP.2022.20005
- Sayyah M, Javad-Pour M, Ghazi-Khansari M (2003) The bacterial endotoxin lipopolysaccharide enhances seizure susceptibility in mice: involvement of proinflammatory factors: nitric oxide and prostaglandins. *Neuroscience* 122(4):1073-80
doi:10.1016/j.neuroscience.2003.08.043
- Shimada T, Takemiya T, Sugiura H, Yamagata K (2014) Role of inflammatory mediators in the pathogenesis of epilepsy. *Mediators Inflamm* 2014:901902
doi:10.1155/2014/901902
- Singh N, Saha L, Kumari P, et al. (2019) Effect of dimethyl fumarate on neuroinflammation and apoptosis in pentylenetetrazol kindling model in rats. *Brain Res Bull* 144:233-245
doi:10.1016/j.brainresbull.2018.11.013
- Stellwagen D, Malenka RC (2006) Synaptic scaling mediated by glial TNF- α . *Nature* 440(7087):1054-9
doi:10.1038/nature04671
- Vezzani A, French J, Bartfai T, Baram TZ (2011) The role of inflammation in epilepsy. *Nat Rev Neurol* 7(1):31-40
doi:10.1038/nrneurol.2010.178
- Vezzani A, Friedman A, Dingledine RJ (2013) The role of inflammation in epileptogenesis. *Neuropharmacology* 69:16-24
doi:10.1016/j.neuropharm.2012.04.004
- Xiang NL, Liu J, Liao YJ, et al. (2016) Abrogating CIC-3 Inhibits LPS-induced Inflammation via Blocking the TLR4/NF- κ B Pathway. *Sci Rep* 6:27583
doi:10.1038/srep27583
- Zareie P, Sadegh M, Palizvan MR, Moradi-Chameh H (2018) Anticonvulsive effects of endocannabinoids; an investigation to determine the role of regulatory components of endocannabinoid metabolism in the Pentylenetetrazol induced tonic-clonic seizures. *Metab*

Carvacrol anticonvulsion in proinflammation

Brain Dis 33(3):939-948
doi:10.1007/s11011-018-0195-5.