

Short-Communication

Investigation of the neuroprotective effect of crocin against electromagnetic field-induced cerebellar damage in male Balb/c mice

Mehrdad Hajinejad¹, Abdolreza Narouiepour^{1,2}, Fatemeh Alipour¹, Alireza Ebrahimzadeh-bideskan^{1,3,*}

¹Department of Anatomy and Cell Biology, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

²Department of Anatomy, School of Medicine, Iranshahr University of Medical Sciences, Iranshahr, Iran

³Applied Biomedical Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

Article history:

Received: Feb 18, 2023

Received in revised form:

Jul 13, 2023

Accepted: Jul 15, 2023

AJP, Vol. 14, No. 3, May-Jun

2024, 289-296.

[https://dx.doi.org/10.22038/](https://dx.doi.org/10.22038/AJP.2023.23005)

AJP.2023.23005

* Corresponding Author:

Tel: +98-5138002486

Fax: +98-5138002487

Ebrahimzadehba@mums.ac.ir

Keywords:

Electromagnetic field (EMF)

Crocin

Cerebellum

Purkinje cell

Astrocytes

Abstract

Objective: Mobile devices are sources of electromagnetic fields (EMFs) that cause increasing concern among scientists about human health, especially with the long-term use of mobile phones. With regard to this issue, the potential adverse health effects, particularly on brain function have raised public concern. There is considerable evidence that natural compounds have neuroprotective effects due to their antioxidant and anti-inflammatory properties. Growing evidence suggests that crocin as a natural bioactive compound can be considered a potential therapeutic agent against various neurologic disorders. Therefore, the present study investigated the effects of crocin on the cerebellum after exposure to EMF.

Materials and Methods: Twenty-four Male Balb/c mice were divided into control group, EMF group (2100 MHz), EMF +Crocin group (2100 MHz+50 mg/kg), and crocin group (50 mg/kg). The animals in the EMF and EMF+Crocin groups were exposed continuously for 30 days to an EMF 120 min/day. After 30 days, cerebellar cortex was evaluated by histomorphometric and immunohistochemical methods.

Results: The results showed that 30 days of exposure to EMF had no significant effect on Purkinje cell size. However, EMF reduced significantly the diameter of astrocytes and increased Glial fibrillary acidic protein (GFAP) expression compared to the controls ($p < 0.05$). Our findings also indicated that crocin treatment could improve the diameter of astrocytes and normalize GFAP expression ($p < 0.05$).

Conclusion: This study concluded that 2100-MHz EMF caused adverse effects on the cerebellum through astrocyte damage and crocin could partially reverse the EMF-related adverse effects.

Please cite this paper as:

Hajinejad M, Narouiepour A, Alipour F, Ebrahimzadeh-bideskan A. Investigation of the neuroprotective effect of crocin against electromagnetic field-induced cerebellar damage in male Balb/c mice. Avicenna J Phytomed, 2024; 14(3): 289-296.

Introduction

Nowadays, with technological advances, the use of mass media such as mobile phones and internet connectivity gadgets (including Wi-Fi modem, computer, etc.) is growing world-wide drastically. Although the potential biohazards of exposure to electromagnetic fields (EMFs) have not been perfectly clear, the usage of these devices has raised concerns regarding exposure. In this regard, the European Union has recommended restrictions on the use of mobile phones and internet access in schools (Obajuluwa *et al.*, 2017) and many studies have been conducted to determine the potential EMF-related adverse bio-effects on various tissue organs. Considering that cell phones are held near the head during the contacts, the brain is exposed to relatively high EMFs compared to other organs and may be a primary target organ for EMFs (Sonmez *et al.*, 2010). Some studies have reported associations between EMFs and impaired spatial learning and memory as well as marked morphological changes in the CA1 sub-region of hippocampus (Li *et al.*, 2012). It has also been found that prenatal exposure to EMFs adversely affects the Purkinje cells development in rats pups (Odacl *et al.*, 2016). From this point of view, EMFs have been indicated to disrupt cerebellar morphology and reduce the number of Purkinje cells in rats (Aslan *et al.*, 2017; Rağbetli *et al.*, 2010).

Crocin, a water-soluble carotenoid compound isolated from *Crocus sativus* L. (saffron), is one of the main saffron bioactive metabolites which is responsible for its color and it is extensively used in modern and traditional medicines. According to previous studies, crocin exhibits antioxidant, anti-tumor, analgesic, anti-inflammatory and anti-anxiety effects (Razavi *et al.*, 2017). Recent evidence has demonstrated the neuro-protective effects of crocin in various neurologic disorders characterized by increased oxidative and inflammatory state. In this way,

administration of crocin might be able to improve memory, behavioral, and histopathological damages in brain tissue. A growing body of evidence indicates that crocin via reducing oxidative stress can be considered a promising candidate for the treatment of oxidative stress-related neurological diseases (Mehri *et al.*, 2015). The aim of this study was to evaluate the neuro-protective effects of crocin on the damage caused by exposure to electromagnetic waves in the cerebellum of male mice.

Materials and Methods

Chemicals

Crocin was purchased from School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran. Primary antibody (Rabbit anti-GFAP antibody; Cat. No. ab7260) and secondary antibody (goat anti-rabbit HRP; Cat. N. ab6721) were purchased from Abcam, USA. Cresyl violet stain solution (Sigma Aldrich, USA). 3, 3'-diaminobenzidine (DAB) was purchased from (Sigma Aldrich, USA), and Hematoxylin stain solution from Merck, Germany.

Animals

Twenty-four male Balb/c mice at 8-weeks of age and weighing 25–35 g were obtained from the animal center of the faculty of medicine, Mashhad University of Medical Science, Mashhad, Iran. All animals were kept in standard condition ($23 \pm 2^\circ\text{C}$ and humidity of $60 \pm 5\%$ with a 12 h light/dark cycle) and experimental procedures were compatible with the Guidelines for Ethical Conduct in the Care and Use of Animals. All experiments were approved by the Ethics Committee of Mashhad University of Medical Sciences, Iran (IR.MUMS.MEDICAL.REC.1399.018).

Crocin preparation

Crocin was purchased in powder from the school of Pharmacy, Mashhad

Crocin against EMF-induced cerebellar damage

University of Medical Sciences, Iran and stored in a dry and dark place. Every day, crocin was freshly prepared at 10 mg/ml concentration by dissolving in distilled water. The volume of injection was adjusted according to the varying weights of mice (50 mg/kg) (Salahshoor et al., 2016)

Electromagnetic field (EMF) delivery

In this study, EMF was generated from a device producing 2100 MHz similar to mobile phones that have 3G feature. For this, a circular box was designed, the device was located in the center of the circle and the animals were placed at equal distances to device and exposed with EMFs for 120 min (Figure 1).



Figure 1. Electromagnetic field-producing device (2100 MHz).

Experimental design

The animals were randomly divided into four groups (n=6):

1. Control group: The animals were injected with distilled water.
2. Electromagnetic field (EMF) group: The animals were exposed to an EMF of 2100 MHz for 120 min daily for 30 consecutive days.
3. EMF+ Crocin group: The animals were exposed to EMF and received crocin (50 mg/kg/day) intraperitoneally (ip) for 30 consecutive days (Vafaei, Motejaded, & Ebrahimzadeh-Bideskan, 2020).
4. Crocin group: The animals received crocin (50 mg/kg/day, ip) for 30 consecutive days.

Tissue procedures

At the end of the experiment, the animals were deeply anesthetized with

ketamine (80 mg/kg) and xylazine (10 mg/kg) and sacrificed by decapitation. The cerebellums were then removed and fixed in 10% formalin solution. After fixation, the specimens were prepared according to the routine histological preparations, embedded in paraffin and then cut into 5- μ m thicknesses coronal sections using Leitz microtome.

Cresyl violet staining

The sections were stained with cresyl violet for morphometric analysis of the cerebellum. Briefly, the sections were dehydrated through descending grades of ethanol. After rinsing with distilled water, the slides were immersed in 0.5% cresyl violet staining solution for 20 min and then washed in distilled water. After dehydration through the ascending grades of ethanol, the sections were immersed two times for 5 min in xylene solution and mounted. Finally, the diameter of the Purkinje cells and astrocytes was measured.

Immunohistochemistry

To evaluate the pattern of glial fibrillary acidic protein (GFAP) in cerebellar cortex, the immunohistochemistry method was performed. Briefly, after dewaxing and gradient rehydration, heat-induced epitope retrieval was done. Then, to quench endogenous peroxidase activity, the sections were incubated in 0.3 % H_2O_2 in methanol for 20 min. Thereafter, blocking solution with 1 % bovine serum albumin (BSA) for 30 min at 37°C was applied to sections. After that, the sections were incubated overnight at 4°C with rabbit polyclonal anti-GFAP antibody (1-2000 diluted). After washing with PBS, the sections were incubated with goat anti-rabbit HRP secondary antibody (1:1000 diluted) at 37°C for 2 hr. The reaction product was visualized using 0.03 % 3, 3'-diaminobenzidine (DAB) in PBS buffer, containing 0.01 % H_2O_2 for 15 min. Finally, hematoxylin staining solution was employed as a counterstain. The cytoplasm

of GFAP -positive cells appeared brown and the nuclei blue.

Morphometric and immunohistochemistry analysis

At first, all sections were photographed using a light microscope (Olympus BX51, Tokyo, Japan) equipped with camera and the taken photos were transferred to a computer. To histomorphometrically evaluate the diameter of Purkinje and astrocyte cells, Image J software was used (Hajje *et al.*, 2014; Rastegar Moghaddam *et al.*, 2022).

Statistical analysis

The statistical analysis was performed with GraphPad Prism version 8.0. The data are shown as the mean±SD. Data from the study was analyzed by one-way Analysis of Variance (ANOVA) followed by Tukey's

test to assess the significance and Kruskal–Wallis followed by Mann-Whitney test. A $p < 0.05$ was considered statistically significant.

Results

Morphometric analysis

The analysis of morphometric parameters showed that there were no significant differences in the diameter of Purkinje cell between the studied groups (Figure 2). However, exposure to EMF significantly reduced the diameter of astrocytes compared with the control group ($p < 0.05$). Our data from the EMF+Crocic groups also showed that the administration of crocin could improve astrocytes diameter compared with the EMF group ($p < 0.05$) (Figures 3 and 4).




Figure 2. Photomicrographs of cerebellar cortex stained with cresyl violet in different studied groups. ML: molecular layer, PCL: Purkinje cell layer, GL: granule cell layer. Purkinje cells (red arrows). No difference was observed between the groups. (Scale bar: 100 μm).




Figure 3. Photomicrographs of cerebellar cortex stained with immunohistochemistry method in different studied groups. GFAP immunostaining was used to compare the astrocytes diameter between the groups. (Scale bar: 20 μm).

Crocini against EMF-induced cerebellar damage




Figure 4. Comparing of the diameter of astrocytes in all the studied groups. *Significant difference between the control and EMF group ($p < 0.05$). #Significant difference between the EMF and EMF+crocini group. Data are presented as mean \pm SD.

Immunohistochemistry analysis

The immunoreactivity of astrocytes was demonstrated using GFAP, a common histological marker for astrocytes. In the control group, GFAP immunoreaction showed a positive brown immunoreactivity in the form of thin and regular brown staining fibers in the astrocytes. In the EMF group, an apparent increase in GFAP immunoreactive astrocytes was observed. Fibers feature in the EMF group was twisted with an irregular course along with an intense deep brown positive immunoreaction for GFAP when compared to the control group ($p < 0.0001$).




Figure 5. Photomicrographs of the GFAP immunostaining sections in cerebellar cortex. Cerebellar cortex of the control group showed a positive GFAP in the form of thin and regular brown fibers. Cerebellar cortex of the animal exposed to EMF showed more diffuse and thick brown fibers of GFAP; cerebellar cortex of the animal treated by crocini with exposure to EMF showed a decrease in GFAP reaction (Scale bar: 20 μ m).




Figure 6. Graph shows GFAP immunoreactivity in different groups. **** $p < 0.0001$ significant difference compared to the control group; * $p < 0.05$ significant difference compared to the EMF group; *** $p < 0.001$ significant difference compared to the crocini group. No significant differences were observed between the control and crocini groups. Data are presented as mean \pm SD.

Comparing with the EMF group, GFAP immunoreactivity in the animals of the EMF+ crocini group, showed an apparent decrease in the amount and intensity of immunoreactivity in the astrocytes ($p < 0.05$). GFAP immunoreactivity in the crocini group was significantly higher than the EMF+crocini group ($p < 0.001$). No significant differences were observed between the control and crocini groups (Figures 5 and 6).

Discussion

Previous studies have indicated widespread debates about the public health

effects of EMF exposure for decades, and a large number of studies have been conducted to assess the potential adverse effects of that exposure (Kivrak *et al.*, 2017; Zymantiene *et al.*, 2020).

In our study, employment of EMF originating from a device similar to mobile phones, disturbed cerebellar cortex architecture. The cerebellum is a region of the brain that plays a major role in motor movement coordination and balance. Furthermore, there is clear evidence that the cerebellum is involved in some complex cognitive processes including emotions, behavior, learning and memory (Ragbetli *et al.*, 2007; Wolf *et al.*, 2009). From this point of view, Purkinje cells as the sole output cells for cerebellar computations play a critical role in cerebellar functioning (Chang *et al.*, 2020). On the other hand, astrocytes as the most abundant glial cells in the CNS have major role in normal and pathological functions of the CNS (Araujo *et al.*, 2019). The worldwide spread of the use of mobile phones has led to concern regarding the possible adverse health effects. These potential health problems, especially for the nervous system, have caused great public concern (Naeem, 2014). Since the cerebellum is the organ that is particularly vulnerable to developmental and environmental insults due to its long developmental schedule, any alteration in the cerebellum seems to lead to motor deficits, dementia, schizophrenia and other psychiatric disorders (Arslan *et al.*, 2022; Othman *et al.*, 2021). Therefore, we aimed to investigate the adverse effect of EMF exposure on male rat cerebellum. The outcomes of the study by Rosli and Teoh (2009) showed that EMFs are capable of causing irreparable damage to adult mice brain which was obviously accompanied by a decrease in the number of Purkinje cells along with the sequel of thinning of the granular layer (Rosli and Teoh, 2009). However, our finding showed that EMF exposure could not significantly reduce the average size of Purkinje cells. On the other

hand, astrocyte activation has been shown to be characterized by increased GFAP expression (Sonmez *et al.*, 2010). Othman *et al.* (2021) showed that GFAP expression in brain tissue after irradiation was not significantly different than those of non-irradiated animals (Othman *et al.*, 2021). Another study reported that reactive astrocytes exhibit no remarkable changes in the volume, but increase the thickness of their main cellular processes (Li *et al.*, 2012). In this study, we also measured the size of astrocytes in the cerebellum. Our findings revealed that EMF reduced the size of astrocyte and administration of crocin could not markedly ameliorate this adverse effect of EMF. Our results also indicated that exposure to EMF induced activation of astrocyte and increased accumulation of GFAP in cellular processes. Interestingly, administration of crocin could markedly improve the detrimental effect of EMF on GFAP expression. Most interestingly, it was reported that astrocytes provide a unique niche with neurogenic capacity capable of promoting proliferation and neuronal fate determination. Furthermore, injury to the CNS activates the astrocytes to be reactive and increase the expression of GFAP. Increase in GFAP levels could be also explained by an increase in the astrocytes to provide more nourishment required for the injured neurons (Ammari *et al.*, 2010; Belal *et al.*, 2020; Pekny *et al.*, 1999). Additionally, GFAP upregulation is considered a major indicator for the astrocytic reactivity that can be commonly detected after neurologic insults. However, the molecular mechanism behind that is poorly understood (Brahmachari *et al.*, 2006; Gomes *et al.*, 1999).

To sum up, our findings showed that exposure to EMF could not significantly affect the change size of Purkinje cells, but it could reduce the size of astrocytes. Furthermore, EMF exposure significantly increased GFAP immunoreactivity. Crocin administration partially could attenuate EMF-induced destructive effects on the

Crocini against EMF-induced cerebellar damage

cerebellum. However, further extensive and rigorous experimental studies and clinical trials are required to adjust the dose and form of crocin treatment.

Acknowledgment

Not applicable.

Conflicts of interest

The authors have declared that there is no conflict of interest.

References

- Ammari M, Gamez C, Lecomte A, Sakly M, Abdelmelek H, De Seze R. 2010. GFAP expression in the rat brain following sub-chronic exposure to a 900 MHz electromagnetic field signal. *Int J Radiat Biol*, 86: 367-375.
- Araujo APB, Carpi-Santos R, Gomes FCA. 2019. The role of astrocytes in the development of the cerebellum. *Cerebellum*, 18: 1017-1035.
- Arslan A, Acer N, Kesici H, Sonmez MF, Ertekin T, Gultekin M, Saracoglu OG. 2022. Stereological Study on the Effect of Carnosine on of Purkinje Cells in the Cerebellum of Rats Exposed to 900 MHz Electromagnetic Field. *Turk Neurosurg*, 32: 618-624.
- Aslan A, İkinci A, Baş O, Sönmez OF, Kaya H, Odacı E. 2017. Long-term exposure to a continuous 900 MHz electromagnetic field disrupts cerebellar morphology in young adult male rats. *Biotech Histochem*, 92: 324-330.
- Belal S, Afifi O, Afeefy A. 2020. Evaluation of mobile phone radiation-induced structural changes of rat brain with emphasis on the possible protective role of pomegranate peel extract. *Int J Radiat Biol*, 18: 753-764.
- Brahmachari S, Fung YK, Pahan K. 2006. Induction of glial fibrillary acidic protein expression in astrocytes by nitric oxide. *J Neurosci*, 26: 4930-4939.
- Chang W, Pedroni A, Hohendorf V, Giacomello S, Hibi M, Köster RW, Ampatzis K. 2020. Functionally distinct Purkinje cell types show temporal precision in encoding locomotion. *Proc Natl Acad Sci*, 117: 17330-17337.
- Gomes FC, Paulin D, Moura Neto V. 1999. Glial fibrillary acidic protein (GFAP): modulation by growth factors and its implication in astrocyte differentiation. *Braz J Med Biol Res*, 32: 619-631.
- Hajje G, Saliba Y, Itani T, Moubarak M, Aftimos G, Farès N. 2014. Hypothyroidism and its rapid correction alter cardiac remodeling. *PLoS One*, 9: e109753.
- Kivrak EG, Altunkaynak BZ, Alkan I, Yurt KK, Kocaman A, Onger ME. 2017. Effects of 900-MHz radiation on the hippocampus and cerebellum of adult rats and attenuation of such effects by folic acid and *Boswellia sacra*. *J Microsc Ultrastruct*, 5: 216-224.
- Li Y, Shi C, Lu G, Xu Q, Liu S. 2012. Effects of electromagnetic radiation on spatial memory and synapses in rat hippocampal CA1. *Neural Regen Res*, 7: 1248-1255.
- Mehri S, Abnous K, Khooei A, Mousavi SH, Shariaty VM, Hosseinzadeh H. 2015. Crocin reduced acrylamide-induced neurotoxicity in Wistar rat through inhibition of oxidative stress. *Iran J Basic Med Sci*, 18: 902-908.
- Naeem Z. 2014. Health risks associated with mobile phones use. *Int J Health Sci (Qassim)*, 8: 5-6.
- Obajuluwa AO, Akinyemi AJ, Afolabi OB, Adekoya K, Sanya JO, Ishola AO. 2017. Exposure to radio-frequency electromagnetic waves alters acetylcholinesterase gene expression, exploratory and motor coordination-linked behaviour in male rats. *Toxicol Rep*, 4: 530-534.
- Odacı E, Hancı H, İkinci A, Sönmez OF, Aslan A, Şahin A, Baş O. 2016. Maternal exposure to a continuous 900-MHz electromagnetic field provokes neuronal loss and pathological changes in cerebellum of 32-day-old female rat offspring. *J Chem Neuroanat*, 75: 105-110.
- Othman H, López-Furelos A, Leiro-Vidal JM, Ammari M, Sakly M, Abdelmelek H, López-Martín E. 2021. Exposure to 2.45 GHz Radiation Triggers Changes in HSP-70, Glucocorticoid Receptors and GFAP Biomarkers in Rat Brain. *Int J Mol Sci*, 22: 5103.
- Pekny M, Johansson CB, Eliasson C, Stakeberg J, Wallén A, Perlmann T, Frisén J. 1999. Abnormal reaction to central nervous system injury in mice lacking glial fibrillary acidic protein and vimentin. *J Cell*

- Biol, 145: 503-514.
- Rağbetli MC, Aydinlioğlu A, Koyun N, Rağbetli C, Bektas S, Ozdemir S. 2010. The effect of mobile phone on the number of Purkinje cells: a stereological study. *Int J Radiat Biol*, 86: 548-554.
- Ragbetli MC, Ozyurt B, Aslan H, Odaci E, Gokcimen A, Sahin B, Kaplan S. 2007. Effect of prenatal exposure to diclofenac sodium on Purkinje cell numbers in rat cerebellum: a stereological study. *Brain Res*, 1174: 130-135.
- Rastegar Moghaddam SH, Hosseini M, Sabzi F, Hojjati Fard F, Marefati N, Beheshti F, Anaigoudari A. 2022. Cardiovascular protective effect of nano selenium in hypothyroid rats: protection against oxidative stress and cardiac fibrosis. *Clin Exp Hypertens*, 44: 268-279.
- Razavi BM, Sadeghi M, Abnous K, Vahdati Hasani F, Hosseinzadeh H. 2017. Study of the role of CREB, BDNF, and VGF neuropeptide in long term antidepressant activity of crocin in the rat cerebellum. *Iran J Pharm Res*, 16: 1452-1462.
- Rosli Y, Teoh PJ. 2009. The effect of low electromagnetic field in the cerebellar layers of mice. *J Biol Sci*, 9: 601-606.
- Salahshoor MR, Khazaei M, Jalili C, Keivan M. 2016. Crocin improves damage induced by nicotine on a number of reproductive parameters in male mice. *Int J Fertil Steril*, 10: 71-78.
- Sonmez OF, Odaci E, Bas O, Kaplan S. 2010. Purkinje cell number decreases in the adult female rat cerebellum following exposure to 900 MHz electromagnetic field. *Brain Res*, 1356: 95-101.
- Vafaei S, Motejaded F, Ebrahimzadeh-Bideskan A. 2020. Protective effect of crocin on electromagnetic field-induced testicular damage and heat shock protein A2 expression in male BALB/c mice. *Iran J Basic Med Sci*, 23: 102-110.
- Wolf U, Rapoport MJ, Schweizer TA. 2009. Evaluating the affective component of the cerebellar cognitive affective syndrome. *J Neuropsychiatry Clin Neurosci*, 21: 245-253.
- Zymantiene J, Juozaitiene V, Zelvyte R, Oberauskas V, Spancerniene U, Sederevicius A, Aniuliene A. 2020. Effect of electromagnetic field exposure on mouse brain morphological and histopathological profiling. *J Vet Res*, 64: 319-324.