

Review Article

Neuroprotective effects of *Coriandrum sativum* and its constituent, linalool: A review

Mahmoud Hosseini¹, Mohammad Hossein Boskabady², Mohammad Reza Khazdair^{3,*}

¹Division of Neurocognitive Sciences, Psychiatry and Behavioral Sciences Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

²Applied Biomedical Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

³Cardiovascular Diseases Research Center, Birjand University of Medical Sciences, Birjand, Iran

Article history:

Received: Feb 13, 2021

Received in revised form:

Mar 08, 2021

Accepted: Mar 10, 2021

Vol. 11, No. 5, Sep-Oct 2021,
436-450.

<https://dx.doi.org/10.22038/AJP.2021.55681.2786>

* Corresponding Author:

Tel: +98-5631626466

Fax: +98-5632234717

khazdairMR@Bums.ac.ir

Keywords:

Coriandrum sativum L.

Neuro-pharmacological effects

Memory impairment

Linalool

Abstract

Objective: Coriander (*Coriandrum sativum* L.) is an annual herb belonging to the Apiaceae (Umbellifera) family that is used as food additives traditionally. This plant is called “*Geshniz*” in Persian and is native to Mediterranean regions but it is currently cultivated in several countries. All parts of coriander are edible and have been traditionally used to treat different disorders, including digestive problems, flatulence, diarrhea, colic and other gastrointestinal diseases.

Materials and Methods: The databases PubMed, Web of Science, Google Scholar and Scopus were considered. The search terms were “*Coriandrum sativum*” or “linalool” and “anti-anxiety”, “sedative”, “antioxidant effect”, “anticonvulsant” and “neuroprotective effect”.

Results: Antioxidant, diuretic, cholesterol lowering, anxiolytic, sedative-hypnotic and anticonvulsant activities were reported for the seeds and leaves of the plant. Furthermore, linalool as the main component of coriander has different neuropharmacological effects, including anti-anxiety, sedative, anticonvulsant and anti-Alzheimer’s disease activities.

Conclusion: Various neuropharmacological effects of *C. sativum* and its component which have antioxidant and anti-inflammatory effects, have been summarized in the current review article.

Please cite this paper as:

Hosseini M, Boskabady MH, Khazdair MR. Neuroprotective effects of *Coriandrum sativum* and its constituent, linalool: A review. Avicenna J Phytomed, 2021; 11(5): 436-450.

Introduction

Coriander (*Coriandrum sativum* L.) (*C. sativum*), is an annual herb originating from Mediterranean areas. The leaves are small branches and sub-branches, the flowers are white and the fruits are almost ovate globular, with multiple longitudinal ridges on the surface (Figure 1) (Yeung

and Bowra, 2011). All segments of this plant are edible and have been traditionally used for treatment of different disorders (Sahib et al., 2013). The fresh leaves can be used for garnishing and are used in many foods like chutneys, salads and soups (Bhat et al., 2014). Green coriander contains 84% water, a low level of

Neuroprotective effects of *Coriandrum sativum*

saturated fats, and cholesterol, but a high level of thiamine, zinc and dietary fibers (Kandlakunta et al., 2008). Coriander seeds are also considered a source of vitamins, lipids and minerals, such as potassium, calcium, phosphorus, magnesium, sodium and zinc (Bhat et al., 2014; Iwatani et al., 2003). The main coriander essential oil (EO) components are linoleic and linolenic acids (Sahib et al., 2013). Nutritional supplementation with coriander seeds is suggested to be able to decrease saturated but increase unsaturated fatty acids (Ertas et al., 2005).

Coriander is commonly used to attenuate digestive problems, flatulence, diarrhea and colic and other gastrointestinal diseases in Iranian traditional medicine (Zargari, 1991). Hepatoprotective properties of *C. sativum* due to the antioxidant potential of its phenolic compounds (Pandey et al., 2011) and cardiovascular protective effect of the plant were demonstrated (Dhanapakiam et al., 2007). The plant also showed antioxidant activities in different organs (Kansal et al., 2011; Wangenstein et al., 2004) including a preventive effect on the changes in the brain histology such as gliosis, lymphocytic infiltration and cellular edema (Vekaria et al., 2012).

It has been reported that treatment with leaves of *C. sativum* (5, 10 and 15% w/w) improved memory scores and prevented

the process of aging in mice dose-dependently, reversed the memory deficits induced by scopolamine and diazepam and significantly reduced brain cholinesterase activity and serum total cholesterol levels (Mani et al., 2011). The preventive effect of the hydro-alcoholic extract of *C. sativum* (50, 100 and 200 mg/kg) on neuronal damages induced by pentylentetrazole (PTZ) in a rat model of seizure, was also demonstrated (Pourzaki et al., 2017). Ethanolic extract of *C. sativum* seeds on learning in second-generation mice was assessed but the plant extract did not improve learning in a short period of time after training; however, it improved learning in long term (Zargar-Nattaj et al., 2011).

There is currently an increasing interest to discover effective drugs based on folk medicinal and medicinal plant. It should be noted that many modern drugs have plants origin, and were developed through observation of therapeutic practices of native people (Balick and Cox, 1996; Gilani, 2005). Therefore, in the present article, neuropharmacological properties of *C. sativum* and its component including their sedative, anti-anxiety, anti-depressive, anticonvulsant and anti-seizures, memory improving, anti-Parkinson's diseases properties and protective effects on neurotoxicity, were reviewed.



Figure 1. Leaf (left), flower (middle) and seeds (right) of coriander (Asgarpanah and Kazemivash, 2012).

Chemical composition of *C. sativum* essential oil

C. sativum is cultivated in different soil, climatic, seasonal and other ecological conditions and therefore, has a broad range of compositions. Different essential oil

(EO) components and different chemical compounds have been extracted from different parts of *C. sativum* which seems to be dependent to the variety of the plant (Diederichsen and Hammer, 2003). EO of premature fruits mainly contains

compounds such as geranyl acetate (46.27%) and linalool (10.96%), while in mature fruits linalool (87.54%), followed by cis-dihydrocarvone (2.36%), were detected as the considerable compounds (Msaada *et al.*, 2007).

The coriander seeds EO from different geographical origins also showed variations in constituents. The major components in Europe type of the plant are linalool (58.0–80.3%), γ -terpinene (0.3–11.2%) and α -pinene (0.2–10.9%) (Orav *et al.*, 2011), while, the main oil components of the plant from Brazil were linalool (77.48%), γ -terpinene (4.64%) and α -pinene (5.5%) (de Figueiredo *et al.*, 2004). Additionally, the seeds oil of the plant grown in Iran contain linalool (40.9–79.9%), neryl acetate (2.3–14.2%) and γ -terpinene (0.1–13.6%) (Nejad Ebrahimi *et al.*, 2010).

The oils extracted from the leaves of the plant have been shown to contain 44 compounds, most of them are aromatic acids, including 2-decenoic (30.82%) and E-11-tetradecenoic acids (13.4%). Capric acid (12.7%) and undecyl alcohol (6.4%). Other compounds including 2-undecenal (3.87%), dodecanoic acid (2.63%), cyclododecane (2.45%), decanal (1.35%) and decamethylene glycol (1.15%) were also reported to be present in the leaves of the plant (Nurzyńska-Wierdak, 2013). Monoterpenoids such as limonene and beta-myrcene are also reported as the constituents of coriander EO (Tashinen and Nykänen, 1975).

Sedative, anti-anxiety and anti-depressive properties of the plant and its component

C. sativum

C. sativum has been recommended for relieving insomnia in Iranian folk medicine (Mir, 1992; Zargari, 1991). Pre-sleep eating of a combination of chopped fresh leaves (30 g) or seeds of the plant with tea, as a single dose, has been suggested to relieve anxiety and insomnia (Mir, 1992). Similar uses have also been

advised in other folk medicines (Duke, 2002). These effects were also confirmed using animal models. Intraperitoneal (i.p.) administration of aqueous extracts (200, 400 and 600 mg/kg) and hydroalcoholic extracts (400 and 600 mg/kg), as well as 600 mg/kg EO of coriander in male albino mice increased sleep duration to 160-220, 130-180 min and 210 min, respectively. These results showed that the extracts and EO of coriander exhibit sedative and hypnotic activities in pentobarbital-induced hypnotic model (Emamghoreishi and Heidari-Hamedani, 2015). In another study, the sleep-prolonging effects of the fractions of aerial parts of the plant including water (WF), ethyl acetate (EAF) and n-butanol (NBF) fractions were also confirmed in mice. The hydro-alcoholic extract (HAE), EAF and NBF significantly prolonged sleep duration but only the NBF decreased sleep latency significantly (Rakhshandeh *et al.*, 2012). Using an animal model of anxiety, the effects of 10, 25, 50 and 100 mg/kg (i.p.) of aqueous extract of the seeds on motor activity and neuromuscular function have been evaluated. In elevated plus-maze, the seeds' aqueous extract at 100 mg/kg (i.p.) showed an anxiolytic effect by increasing the time spent in and the entries to the open arms. Furthermore, administration of the extract (50, 100 and 500 mg/kg) was accompanied with motor activity impairments and neuromuscular dysfunction (Emamghoreishi *et al.*, 2005). In another study, anti-anxiety effects of 100 and 200 mg/kg of hydroalcoholic extract of the seeds were comparable to the effects of 0.5 mg/kg diazepam, however, 50 mg/kg of the extract did not show an anti-anxiety activity (Mahendra and Bisht, 2011). The antianxiety activity of 50, 100 and 200 mg/kg (i.p.) of *C. sativum* leaves aqueous extract has also been evaluated in mice, and was compared with the effects of diazepam (5 mg/kg, i.p.) as a standard drug. Treatment by 200 mg/kg of leaves extract increased the time spent in and the number of entries into the open arms of

elevated plus maze which was assumed to be due to anxiolytic effects of the extract (Pathan et al., 2011). Interestingly, 100 and 200 mg/kg of the fruit extract also had an anxiolytic effect which was similar to the effects of the leave extract. Locomotion activity and frequency of rearing in the groups treated by 200 mg/kg of the extract were also lower than that of the control group. Furthermore, 100 and 200 mg/kg of the plant extract significantly increased social interactions of treated animals (Mahendra and Bisht, 2011). In another study, 10 ml/kg (i.p.) of an extract obtained from the fresh plant attenuated immobility time and was more effective than the extract of dried plant. It was concluded that the extract of green but not dried plant have anti-depressive effect (Kishore and Siddiqui, 2003).

Using tail suspension and forced swimming tests, aqueous extract and fixed oil of the seeds showed anti-depressant activity at the end of 14 consecutive days of treatment. Treatment by aqueous extract 200 and 400 mg/kg, orally (p.o.) and also by 2 and 4 ml/kg (p.o.) of an extract prepared by diethyl ether showed a significant anti-depressant like activity, which was similar to the effects of commonly used anti-depressant drugs fluoxetine 20 mg/kg (p.o.) and imipramine 15 mg/kg (p.o.). Aqueous extract from the seeds also inhibited monoamine oxidase-B (MAO-B) activity, which was suggested as a mechanism of anti-depressant activity (Kharade et al., 2011). Essential oil from coriander seeds and its major component linalool has also been administered intracerebroventricularly (i.c.v.) to examine their effects on locomotor activity in chicks. Similar to diazepam, coriander oil and linalool (0.86, 8.6 and 86 µg/chick, i.c.v. for both agent) significantly decreased escapes, defecations and crossing numbers in open field, while increased the sleeping posture (Gastón et al., 2016).

The chemical composition of *C. sativum* volatile oil included linalool

(69.358%), as the major component, γ -terpinene (7.729%), and α -pinene (6.509%). Inhalation of coriander oil (1 and 3%) significantly increased time spent in the open arms, open-arm entries and the number of crossing (exploratory locomotor activity) in a β -amyloid (1–42) rat model of Alzheimer's disease (Cioanca et al., 2014). *C. sativum* oil also reduced significantly immobility time, but increased swimming time and content of glutathione. Furthermore, amyloid deposits were scarce in rats treated with *C. sativum* oil, which were obvious and abundant in the hippocampus of rats treated with i.c.v. β -amyloid (1–42) (Cioanca et al., 2014).

The hydro-alcoholic extract of *C. sativum* (50, 100 and 200 mg/kg) reduced duration, frequency and amplitude of the burst discharges while prolonged the latency of the seizure attacks. The extract also prevented from production of dark neurons and apoptotic cells in different areas of the hippocampus (Pourzaki et al., 2017) (Table 1).

Linalool

Inhalation of 1 and 3% of linalool increased the sleeping time and induced similar effects as 1 mg/kg diazepam. It also reduced body temperature but was not able to affect the bar descent latency in rota-rod performance test. Linalool 3% also significantly impaired exploratory and motor performances in spontaneous locomotor activity evaluation (de Moura Linck et al., 2009).

Similar to administration of 1 mg/kg (i.p.) diazepam, inhalation of 3% linalool increased the time spent in the light compartment of light/dark test box and increased the latency in the first crossing between the light and dark compartments, but had no effect on the crossing number (Linck et al., 2010). Both diazepam and linalool (1%) improved the social interaction behaviors. Furthermore, both linalool 3% and diazepam postponed the first attack and significantly reduced the number and duration of the attacks in the

aggressive behavior test (Linck et al., 2010).

In elevated plus-maze test, inhalation of 0.65, 2.5 and 5.0% mass fraction (w/w) of linalool oxide and injection of 0.5 mg/kg of diazepam (i.p.) were able to increase the open arm entries and the time spent in open arms. Additionally, both linalool oxide and diazepam prolonged the time spent in the light compartment, as well as the crossing numbers between the light and dark compartments of light-dark box (Souto-Maior et al., 2011). Intraperitoneal administration of linalool (30 mg/kg) showed a significant antidepressant activity by increasing the latency time and self-cleaning time, while decreased immobility time in rats (dos Santos et al., 2018). Linalool (100 mg/kg, i.p.) significantly improved the cognitive performance in Morris water maze (MWM) test in an amyloid-beta (A β 1-40)-induced model of cognitive deficits in mice. Treatment with linalool significantly decreased the prolonged time in dark chamber in mice. Treatment with linalool also reduced the pathological lesions induced by A β to normal range. In addition, linalool (100 mg/kg) significantly increased superoxide dismutase (SOD) and glutathione peroxidase (GPX) activities in the hippocampus and cortex compared to the induced animals, while decreased the MDA level in the cortex close to the sham group (Xu et al., 2017). Oral administration of linalool (25 mg/kg), every 48 hr for 3 months, improved learning and spatial memory and behavior in the elevated plus maze in a mouse model of Alzheimer's disease. Linalool (25 mg/kg) also significantly reduced extracellular β -amyloidosis, astrogliosis and microgliosis and reduced the levels of the pro-inflammatory markers including; p38 mitogen-activated protein kinases (p38 MAPK), cyclooxygenase-2 (COX-2) and interleukin (IL)-1 β in hippocampi and amygdalae (Sabogal-Guáqueta et al., 2016) (Table 1).

Anticonvulsant and anti-seizure effects of *C. sativum* and linalool

C. sativum

Anticonvulsant activity of aqueous and ethanolic extracts of coriander seeds was studied using electroshock and PTZ-induced seizure models. Administration of 0.5 g/kg (i.p.) of aqueous and 3.5 and 5 g/kg (i.p.) of ethanolic extracts shortened tonic seizures period and inhibited maximal electroshock-induced seizures. In addition, both of the extracts postponed the onset of clonic convulsions. The ethanolic extract effect was similar to that of 20 mg/kg of phenobarbital in the PTZ test (Hosseinzadeh and Madanifard, 2000).

The effects of 5 mg/kg (i.p.) of the plant extract, as well as the effect of essential oil, on PTZ (85 mg/kg, i.p.)-induced convulsions were also assessed. Intraperitoneal administration of aqueous extracts, as well as essential oils, obtained from the seeds 30 min before PTZ injection, was effective to postpone myoclonic and clonic seizure onset in mice in a dose-dependent manner (Emam and Heydari, 2008). Similarly, 200, 400, 600 and 800 mg/kg of aqueous and ethanolic extracts, as well as essential oil, of the seeds were administered 30 min before PTZ injection (90 mg/kg, i.p.) to evaluate the anticonvulsant effects. The results showed that 600 and 800 mg/kg of all extracts increased myoclonic and clonic seizures latencies (Emamghoreishi and Heidari-Hamedani, 2010).

The effect of leaves hydroalcoholic extract of coriander on oxidative damages in the brain tissues of PTZ-induced seizure was investigated (Karami et al., 2015). Administration of 100, 500 and 1000 mg/kg (i.p.) of the extract postponed the onset of the minimal clonic (MCS) and generalized tonic-clonic (GTCS) seizures. Furthermore, cortical and hippocampal tissues MDA levels in the animals pretreated by the extract decreased significantly.

Neuroprotective effects of *Coriandrum sativum*

Table 1. Sedative, anti-anxiety and anti-depressive effects of *C. sativum* and linalool

Extract/Component	Type of study	Doses	Results	Ref.
Aqueous fresh leaves	Traditional medicine	-	Anxiety relieving and insomnia treating.	(Mir, 1992)
Aqueous fresh leaves	Traditional medicine	-	Anxiety relieving and insomnia treating.	(Duke, 2002)
Aqueous seed extract	Elevated plus maze in mice	10, 25, 50, 100 mg/kg, i.p.	Increasing the time spent on elevated plus maze open arms motor activity and neuromuscular function impairments.	(Emamghoreishi et al., 2005)
Aqueous seed extract	Pentobarbital-induced hypnosis in mice	100, 200, 400 and 600 mg/kg, i.p.	200, 400 and 600 mg/kg of the extract prolonged duration of sleeping time.	(Emamghoreishi and Heidari-Hamedani, 2015)
Hydroalcoholic seed extract	Pentobarbital-induced hypnosis in mice	100, 200, 400 and 600 mg/kg, i.p.	400 and 600 mg/kg of the extract increased duration of sleeping time	(Emamghoreishi and Heidari-Hamedani, 2015)
Essential oil of seed	Pentobarbital-induced hypnosis in mice	100, 200, 400 and 600 mg/kg, i.p.	Only 600 mg/kg of essential oil of seeds increased duration of sleeping time	(Emamghoreishi and Heidari-Hamedani, 2015)
Aqueous leaves extract	Elevated plus maze in mice	50, 100 and 200 mg/kg, i.p.	200 mg/kg of the extract increased the time spent on and the number of entries into the open arms	(Pathan et al., 2011)
Hydroalcoholic seed extract	Induced anxiety in mice	50, 100 and 200 mg/kg, i.p.	Similar to 0.5 mg/kg diazepam, 100 and 200 mg/kg of the extract attenuated anxiety like behaviors	(Mahendra and Bisht, 2011)
An extract obtained from fresh and dried plant	Induced depression in mice	10 ml/kg, i.p.	The extract from fresh plant had an anti-depression effect	(Kishore and Siddiqui, 2003)
Aqueous extracts and extract obtained by diethyl ether	Induced depression in mice	200 and 400 mg/kg, p.o. 2 and 4 ml/kg, p.o.	Showed anti-depressant like activity comparable to the effects of fluoxetine and imipramine	(Kharade et al., 2011)
Diethyl ether extract from the seeds	Induced depression in mice	6 and 8 ml/kg, p.o.	Showed better bioactivity than aqueous seeds extract	(Kharade et al. 2011)
Coriander oil and linalool	Induced anxiety in chick	0.86, 8.6 and 86 µg/chick, i.c.v.	Distress calls, escapes, defecation and crossing number on open field test were decreased, while sleeping posture was increased	(Gastón et al., 2016)
Inhalation coriander oil	Induced Alzheimer's disease in rat	1% and 3%	Increased time spent in the open arms, open-arm entries and the number of crossing in rat model of Alzheimer's disease. <i>C. sativum</i> oil also reduced immobility time, but increased swimming time and content of glutathione. Furthermore, reduced amyloid deposits in the hippocampus	(Cioanca et al., 2014).
<i>C. sativum</i> hydro-alcoholic extract	Induced seizure in rat	50, 100 and 200 mg/kg	Reduced duration, frequency and amplitude of the burst discharges while prolonged the latency of the seizure attacks. The extract also prevented from production of dark neurons and apoptotic cells in different areas of the hippocampus	(Pourzaki et al., 2017)
Inhaled linalool	Induced anxiety in mice	1% and 3%	Linalool 3% decreased exploring and motor activities. Linalool (1% and 3%) increased sleeping time and reduced body temperature.	(de Moura Linck et al., 2009)

Inhaled linalool	Induced anxiety in mice	1% and 3%	Linalool (3%) increased exploring behavior comparable to the effects of diazepam (1.0 mg/kg, i.p.) Linalool (1%) increased social interactions	(Linck et al., 2010)
Inhaled linalool oxide	Induced anxiety in mice	0.65%, 2.5% and 5.0% (w/w)	Increased open entries and open arm duration and also increased the time spent on light compartment.	(Souto-Maior et al., 2011)
linalool	Induced depression in rat	30 mg/kg, p.o.	Showed antidepressant activity by increased latency time and self-cleaning time, while decreased immobility time.	(dos Santos et al., 2018)
Linalool	Induced anxiety in mice	100 mg/kg, i.p.	Improved the cognitive performance and decreased the prolonged time in dark chamber of model mice. Linalool also reduced the pathological lesions induced by AB to normal range. It also increased SOD and GPX activities in the hippocampus and cortex compared to the induced animals, while decreased the MDA level in the cortex close to the sham group	(Xu et al., 2017)
linalool	Induced Alzheimer's disease in mice	25 mg/kg, p.o.	Improved learning and spatial memory and significantly reduced extracellular β -amyloidosis, astrogliosis and microgliosis as well as reduced the levels of the pro-inflammatory markers including; p38 MAPK, COX2 and IL-1 β in hippocampi and amygdalae.	(Sabogal-Guáqueta et al., 2016)

Pretreatment with 500 mg/kg of the extract also significantly improved total thiol contents in the cortical tissues (Karami et al., 2015). Similarly, the effects of 25 and 100 mg/kg (i.p.) of WF, NBF and EAF fractions of the plant on the brain tissues oxidative damages due to seizures induced by PTZ, were investigated. The GTCS latency in WF and EAF treated rats (100 mg/kg, i.p) was longer than that of PTZ group. Interestingly, WF, NBF, and EAF attenuated hippocampal MDA levels. Additionally, both WF and NBF increased cortical and hippocampal thiol contents (Aanaigoudari et al., 2016), (Table 2).

Linalool

Anticonvulsant effects of linalool on experimental seizure models related to glutamate including N-methyl-D-aspartate (NMDA), quinolinic acid-induced convulsions, and the behavioral and

neurochemical factors related to PTZ-kindling were investigated. Linalool at 0.3 and 1.0 mM was able to modulate glutamate activation expression in the rat cortical membrane cells (*in vitro*). Linalool (350 mg/kg, i.p.) and diazepam also significantly prolonged the delay time to the onset of NMDA (270 mg/kg, i.p.)-induced seizures. Administration of linalool (15, 30 and 45 mM, i.c.v.) significantly inhibit quinolinic acid (9.2 mM, i.c.v.)-induced seizure attack as dose-dependent manner. In addition, linalool (2.2 and 2.5 g/kg, p.o) significantly inhibited and delayed the behavioral expression in experimental seizure models (Elisabetsky et al., 1999). In mice cortical synaptosomes, linalool (1.0 or 3.0 mM) reduced potassium-stimulated glutamate release as well as glutamate uptake (90%) (Brum et al., 2001) (Table 2).

Neuroprotective effects of *Coriandrum sativum*

Table 2. Anticonvulsant and anti-seizure activity of *C. sativum* and linalool

Extract/ Component	Type of study	Doses	Results	Ref.
The seeds aqueous extract		0.5 g/kg, i.p.	Tonic seizures duration was decreased, clonic convulsion onsets were postponed.	(Hosseinzadeh and Madanifard, 2000)
The seeds ethanolic extract		3.5 and 5 g/kg, i.p.	Tonic seizures duration was decreased, clonic convulsion onsets were postponed.	(Hosseinzadeh and Madanifard, 2000)
Aqueous and ethanolic extracts and essential oil	Induced seizure in mice	200, 400, 600 and 800 mg/kg, i.p.	Myoclonic and clonic seizures onsets were postponed.	(Emamghoreishi and Heidari-Hamedani, 2010)
The seeds hydroalcoholic extract and essential oil		5 mg/kg, i.p.	Myoclonic and clonic seizures onsets were postponed.	(Emam and Heydari, 2008)
The leaves hydroalcoholic extract		100, 500 and 1000 mg/kg, i.p.	MCS and GTCS latencies were increased Cortical tissues total thiol contents were improved.	(Karami et al., 2015)
Aerial parts fractions	PTZ-induced seizure in rat	25 and 100 mg/kg, i.p.	GTCS latency was significantly improved Hippocampal MDA concentrations were decreased cortical and hippocampal tissues thiol contents were improved.	(Anaegoudari et al., 2016)
Linalool	<i>In vitro</i>	0.3 mM or 1.0 mM	Modulated glutamate activation expression in the rat cortex membrane cells.	(da-Silva et al., 1990)
Linalool	Quinolinic acid induced seizure in rat	350 mg/kg, i.p.	Linalool delayed of NMDA (270mg/kg, i.p.) induced seizures onsets which was comparable to the effects of diazepam.	(da-Silva et al. 1990)
Linalool	Quinolinic acid induced seizure in mic	1.0 or 3.0 mM	Reduced potassium-stimulated glutamate release as well as glutamate uptake (90%)	(Brum et al., 2001)

Memory improving effects of *C. sativum* and its component *C. sativum*

Effects of daily inhalation of volatile oil (1 and 3%) for 21 days on spatial memory performance were assessed in an A β 1-40-induced Alzheimer's disease model in rats. Volatile oils (both 1 and 3%) improved and reversed deleterious effects of (A β 1-40 i.c.v.) injection and increased percentage of spontaneous alternations and number of arm entries, reduced working and improved errors in radial arm-maze task. Coriander volatile oil also improved hippocampal tissues oxidative stress markers in A β 1-40-treated rats by increasing GPX levels and reduction of SOD, lactate dehydrogenase (LDH) and MDA levels in a dose-dependent manner. The amyloid deposits were scarce in volatile oil-treated animals. Also, cleavages of DNA were absent in the coriander oil groups (Cioanca et al., 2013). Similarly, both 1 and 3% of volatile oil increased the time spent in and the

entering number into the open arms in elevated plus-maze compared to the A β group. The volatile oil also enhanced the swimming time and decreased the immobility time in Alzheimeric rats. The antioxidant parameters such as catalase (CAT) activity decreased and total content of reduced glutathione (GSH) were increased in the hippocampal tissues of the rats (Cioanca et al., 2014).

The protective and therapeutic effects of coriander seed aqueous extract in an aluminum chloride (AlCl₃)-induced animal model of Alzheimer's disease and its effects on the cortical pyramidal cells were investigated in male albino rats. Treatment with aqueous seed coriander extract (0.5 mg/kg, p.o) for one month after stopping AlCl₃ (300 mg/kg, p.o) treatment, restored deleterious effects on the pyramidal cells including, dilatation of blood capillaries and presence of many shrunken pyramidal cells (Enas, 2010).

Anti-amnestic and anti-stress properties of *C. sativum* extract on vanillylmandelic

acid (VMA) and ascorbic acid urinary levels in scopolamine-induced amnesia model were evaluated in rats. Daily administration of 100, 200 and 300 mg/kg of the extract one-hour former to exposure of animals to stress, decreased the stress-induced urinary levels of VMA, while increased excretion of ascorbic acid. Moreover, *C. sativum* extract dose-dependently reversed acquisition, retention and recovery impairments induced by scopolamine (1 mg/kg, i.p.) which was accompanied with lipid peroxidation inhibition in both liver and brain tissues (Koppula and Choi, 2012).

Oral administration of young and aged rats by the leaves extract (5, 10 and 15 % w/w) for 45 days, dose-dependently improved memory. The leaves extract also reversed effectively the memory impairments induced by scopolamine (0.4 mg/kg, i.p.) or diazepam (1 mg/kg, i.p.) administration in plus-maze and Hebb-Williams maze models for testing memory (Mani and Parle, 2009; Mani et al., 2011). In learning and memory impairment due to seizures induced by pilocarpine (30 mg/kg, i.p.), administration of 200 mg/kg (i.p.) of the extract for 7 consecutive days, increased the latencies of seizure onset which was accompanied with a decreased level of latency to find the platform in Morris water maze (Elahdadi-Salmani et al., 2015), (Table 3).

Linalool

Beneficial properties of linalool on triple transgenic Alzheimer's disease model (3xTg-AD) in mice were studied. Both non transgenic and the 3xTg-AD mice received 25 mg/kg of linalool (p.o.) every 48 hr for 3 months, significantly better retention performance and swam a longer distance in the platform location compared to vehicle-treated 3xTg-AD mice. Linalool treatment successfully increased the time spent in and the entries into the open arms compared to vehicle-

treated mice. Linalool treatment also delayed deposits and abundance of A β peptide and significantly reduced the A β (1–40), (1–42) protein and pair helical filaments (PHFs) levels in the hippocampal tissues.

Additionally, immunohistochemistry study showed that glial fibrillary acidic protein (GFAP) decreased in the CA1 area of the hippocampus, the entorhinal cortex (EC) and amygdala in mice. In addition, linalool significantly reduced extracellular of A β concentration, astroglia and microglia as well as the pro-inflammatory marker levels, including mitogen-activated protein kinases p38, nitric oxide (NO) synthase 2, cyclooxygenase-2 and interleukin 1 beta in the AD treated mice (Sabogal-Guáqueta et al. 2016).

The effects of linalool (1%) inhalation on acquisition memory in the step-down inhibitory avoidance test showed the anxiolytic properties, increased social interaction and reduced aggressive behavior as well as improved memory in comparison with the control mice (Linck et al., 2010), (Table 3).

Anti-Parkinson effects

The effects of 100 and 200 mg/kg of the seeds ethanolic extract were evaluated on orofacial dyskinesia induced by tacrine. Administration of 2.5 mg/kg of tacrine (i.p), induced some of orofacial dyskinesia symptoms including tongue protrusions (TP), vacuous chewing movements (VCM) and orofacial bursts (OB) 1 hr after administration. Pre-treatment by 100 and 200 mg/kg (p.o.) of the seed extract for 15 days considerably diminished VCM, TP and OB induced by tacrine, and improved locomotor activity and cognitive functions. In addition, administration of the extract significantly improved SOD, CAT and GSH levels, while attenuated lipid peroxidation (Mohan et al., 2015), (Table 3).

Neuroprotective effects of *Coriandrum sativum*

Table 3. Memory improving and anti-Alzheimer's disease properties of *C. sativum* and linalool

Extract/ Component	Type of study	Doses	Results	Ref.
Coriander volatile oil		1 and 3%	Improved and reversed the deleterious effects of amyloid β .	(Cioanca et al., 2013)
Coriander volatile oil		1 and 3%	Enhanced the swimming time and attenuated the immobility times	(Cioanca et al., 2014)
Aqueous seed extract		0.5 mg/kg, p.o.	Restored deleterious effects on pyramidal cells.	(Enas, 2010)
Aqueous fruit extract		100, 200 and 300 mg/kg	Decreased VMA urinary levels in stress-induced animal model while, increased excretion of ascorbic acid.	(Koppula and Choi, 2012)
Leaves mixed in diet	Induced Alzheimer's disease in rats	5, 10 and 15% (w/w)	Improved memory scores in aged and young rats.	(Mani and Parle, 2009; Mani et al., 2011)
Ethanollic leaves extract		200 mg/kg; i.p.	Increased the latencies of seizure onsets which was accompanied with a decreased latency to find the platform.	(Elahdadi-Salmani et al., 2015)
Ethanollic seeds extract		100 and 200 mg/kg, p.o.	Decreased VCM, TP and OB behaviors induced by tacrine, and improved locomotor activity and cognitive functions.	(Mohan et al., 2015)
Linalool	Induced Alzheimer's disease in mice	25 mg/kg, p.o.	Showed a higher retention performance.	(Sabogal-Guáqueta et al., 2016)
Inhaled linalool		1%	The social interactions were increased.	(Linck et al., 2010)

The effects of *C. sativum* and its components on neurotoxicity *C. sativum*

The beneficial effects of the seeds hydroalcoholic extract on neurodegeneration induced by lead exposure were evaluated. Daily treatment by 250 and 500 mg/kg of the extract for seven days which was started after 4 weeks of lead exposure, increased reactive oxygen species (ROS) production and lipid peroxidation and diminished total protein carbonyl content in the cerebellum, hippocampus, frontal cortex and brain stem (Velaga et al., 2014). Hydroalcoholic extract, WF, EAF and NBF (0.1, 0.2, 0.4, and 0.8 mg/ml) had a protective activity on neurons (PC12 cells) in glucose/serum deprivation-induced cytotoxicity. The extract and the fractions showed no cytotoxicity in standard conditions while EAF or NBF (1.6 mg/ml) decreased cell survival. On the other hand, glucose/serum deprivation induced cytotoxicity was significantly reduced by WF at 0.4, 0.8 and 1.6 mg/ml (Ghorbani et al., 2011).

The leaves methanolic extract (200 mg/kg, p.o., per day) increased endogenous enzyme levels such as SOD, glutathione, catalase and total protein levels, while reduced infarcted volume of the brain and attenuated the brain tissues lipid peroxidation and calcium concentration in the brains of the rats in ischemia conditions induced by bilateral common carotid arteries occlusion. Additionally, the blood vessels congestions and neutrophil infiltration were observed which were accompanied with an increasing level of intercellular spaces due to ischemia in rats. All the damages were improved by the extract. The infarction area in the caudal side of the hippocampus was also reduced after treatment by the extract (Vekaria et al., 2012), (Table 4).

Linalool

The possible neuroprotective effect of linalool was investigated in a glucose/serum deprivation-induced cytotoxicity model. Glucose/serum deprivation reduced cell viability when

were cultured for 8 hr compared to the standard condition culture but treatment with 16 µg/ml of linalool improved cell viability (Alinejad *et al.*, 2013).

The effect of linalool on neuronal damages induced by acrylamide (ACR) was also evaluated. ACR is potentially a water-soluble neurotoxic monomer and is extensively used in different industries. In both human and animals, this monomer damages both the central nervous system (CNS) and peripheral nervous system (PNS). ACR-induced neurotoxicity has been suggested to be mainly due to oxidative stress. Administration of 12.5, 25, 50 and 100 mg/kg (i.p.) of linalool for 11 days significantly reduced severe gait abnormalities induced by ACR. Linalool also increased GSH content, however, it decreased lipid peroxidation in the cortical tissues of the rat brain (Mehri *et al.*, 2015).

Linalool significantly reduced neuronal injury in the cortical tissues due to oxygen-glucose deprivation/reoxygenation (OGD/R), while it was not able to inhibit NMDA-induced excitotoxicity. It also significantly scavenged peroxy radicals and reduced intracellular oxidative stress during OGD/R. Monocyte-chemoattractant protein-1 (MCP-1), as a chemokine which is released during OGD/R and induces microglial migration was also inhibited by linalool (Park *et al.*, 2016a),

Linalool at lower concentrations (1, 2.5, and 5 µM) had no cytotoxic effects on SH-SY5Y cells, and did not significantly induce NO. Incubation of the cells with sodium nitroprusside (SNP, 2.5 mM) reduced cell viability. Linalool also increased neuronal cell viability (to 60% cell viability) and significantly prevented NO over-production. In addition, linalool significantly increased antioxidant levels and protected SH-SY5Y cells against SNP-induced cytotoxicity (Kim *et al.*, 2015). It has been reported that linalool (10 µM) has therapeutic effects in treating ischemia-induced cerebral neuronal injury, by inhibition of oxidative stress and

inflammatory responses in rat cortical cells (Park *et al.*, 2016b). Treatment of rats with linalool (12.5 mg/kg, i.p.) showed neuroprotective effects by reducing the progressive gait abnormalities promoted by ACR, oxidative stress, lipid peroxidation, and increasing the levels of GSH (Mehri *et al.*, 2015), (Table 4).

Beta-myrcene

The effects of β-myrcene (a monoterpene derived from coriander) on oxidative and histological damages induced by global cerebral ischemia/reperfusion (I/R) in tissues of C57BL/J6 mice brain, were investigated. Apoptosis in the tissues and other damages to the brain were revealed following I/R induction, while the antioxidant factors, including GSH, catalase, GPX and SOD were decreased. Treatment of animals by myrcin (MYR) (200 mg/kg, i.p.) for 10 days, significantly decreased thiobarbituric acid reactive substances (TBARS) formation. Furthermore, MYR protected against oxidative stress factors and significantly increased GSH, GPX, and SOD. Additionally, MYR showed a neuroprotective effect and decreased incidence of histopathological damages and apoptosis in the brain tissue (Ciftci *et al.*, 2014), (Table 4).

Conclusion

Various parts of the plant, such as seeds and leaves, possess antioxidant, anxiolytic, antidepressive, learning and memory improving, neuroprotective, sedative-hypnotic, analgesic and anticonvulsant activities. Furthermore, linalool have different neuropharmacological effects, including anti-anxiety, sedative, anticonvulsant and anti-Alzheimer's activities. Most of neuroprotective effects of *C. sativum* and its main component may be due to their antioxidant and anti-inflammatory properties.

Neuroprotective effects of *Coriandrum sativum*

Table 4. The effects of *C. sativum* and its components on neurotoxicity

Extract/ Component	Type of study	Doses	Results	Ref.
The leaves methanolic extract	Ischemic rat model	200 mg/kg, p.o.,	Increased total proteins, superoxide dismutase, catalase and glutathione while, attenuated cerebral infarcted volume, prevented from increasing of lipid peroxidation and calcium levels in the ischemic rat brain.	(Vekaria et al., 2012)
WF, EAF and NBF of <i>C. Sativum</i>	<i>In vitro</i>	0.1, 0.2, 0.4, and 0.8 mg/ml	WF at 0.4, 0.8 and 1.6 mg/ml significantly reduced glucose/serum deprivation-induced cytotoxicity.	(Ghorbani et al., 2011)
Linalool	<i>In vitro</i>	16 µg/ml	Significantly improved cell viability	(Alinejad et al., 2013)
Linalool	Acrylamide (ACR) induced neural damage in rat	12.5, 25, 50 and 100 mg/kg i.p.	Significantly reduced severe gait abnormalities induced by ACR	(Mehri et al., 2015)
Linalool	<i>In vitro</i>	0.01, 0.1, 1 and 10 µM	Significantly attenuated intracellular oxidative damages in OGD/R model. It also scavenged peroxy radicals and also inhibited the monocyte-chemoattractant protein-1.	(Park et al., 2016a)
Linalool	<i>In vitro</i>	1 µM, 2.5 µM, and 5 µM	Increased neuronal cell viability (about 60% cell viability) and significantly prevented from NO over- production. Linalool also improved antioxidant status.	(Kim et al., 2015)
Linalool	<i>In vitro</i>	10 µM	Showed therapeutically effects in treating ischemia-induced cerebral neuronal injury, by inhibition of oxidative stress and inflammatory responses.	(Park et al., 2016b)
Linalool	ACR-induced neural damage in rat	12.5 mg/kg, i.p.	Showed neuroprotective effects by reducing the progressive gait abnormalities promoted by acrylamide, oxidative stress, lipid peroxidation, and increasing the levels of glutathione (GSH)	(Mehri et al., 2015)
Beta-myrcene (MYR)	Induced ischemia in mice	200 mg/kg; i.p.	Decreased in the antioxidant factors including, GSH, catalase, glutathione peroxidase (GPX) and SOD.	(Ciftci et al., 2014)

Acknowledgment

Not applicable.

Conflicts of interest

The authors have declared that there is no conflict of interest.

References

- Alinejad B, Ghorbani A, Sadeghnia HR. 2013. Effects of combinations of curcumin, linalool, rutin, safranal, and thymoquinone on glucose/serum deprivation-induced cell death. *Avicenna j phytomed*, 3: 321-328.
- Anaeigoudari A, Hosseini M, Karami R, Vafae F, Mohammadpour T, Ghorbani A, Sadeghnia HR. 2016. The effects of different fractions of *Coriandrum sativum* on pentylenetetrazole-induced seizures and brain tissues oxidative damage in rats. *Avicenna J Phytomed*, 6: 223-235.
- Asgarpanah J, Kazemivash N. 2012. Phytochemistry, pharmacology and medicinal properties of *Coriandrum sativum* L. *Af J Pharm Pharmacol*, 6: 2340-2345.
- Balick MJ, Cox PA. 1996. Plants, people, and culture: the science of ethnobotany. Scientific American Library.
- Bhat S, Kaushal P, Kaur M, Sharma H. 2014. Coriander (*Coriandrum sativum* L.): Processing, nutritional and functional aspects. *Af J Plant Sci*, 8: 25-33.
- Brum LS, Emanuelli T, Souza D, Elisabetsky E. 2001. Effects of linalool on glutamate

- release and uptake in mouse cortical synaptosomes. *Neurochem Res*, 26: 191-194.
- Ciftci O, Oztanir MN, Cetin A. 2014. Neuroprotective effects of β -myrcene following global cerebral ischemia/reperfusion-mediated oxidative and neuronal damage in a C57BL/6 mouse. *Neurochem res*, 39: 1717-1723.
- Cioanca O, Hritcu L, Mihasan M, Hancianu M. 2013. Cognitive-enhancing and antioxidant activities of inhaled coriander volatile oil in amyloid β (1–42) rat model of Alzheimer's disease. *Physiol behav*, 120: 193-202.
- Cioanca O, Hritcu L, Mihasan M, Trifan A, Hancianu M. 2014. Inhalation of coriander volatile oil increased anxiolytic–antidepressant-like behaviors and decreased oxidative status in beta-amyloid (1–42) rat model of Alzheimer's disease. *Physiol behav*, 131: 68-74.
- da-Silva V, De-Freitas J, Mattos A, Paiva-Gouvea W, Presgrave O, Fingola F, Menezes M, Paumgarten F. 1990. Neurobehavioral study of the effect of beta-myrcene on rodents. *Brazil j med biol res*, 24: 827-831.
- de Figueiredo RO, Marques MOM, Nakagawa J, Ming LC. 2004. Composition of coriander essential oil from Brazil. *Acta Hort*, 629: 135-138.
- de Moura Linck V, da Silva AL, Figueiró M, Piatto AL, Herrmann AP, Birck FD, Caramao EB, Nunes DS, Moreno PRH, Elisabetsky E. 2009. Inhaled linalool-induced sedation in mice. *Phytomedicine*, 16: 303-307.
- Dhanapakiam P, Joseph JM, Ramaswamy V, Moorthi M, Kumar AS. 2007. The cholesterol lowering property of coriander seeds (*Coriandrum sativum*): mechanism of action. *J Env Biol*, 29: 53-56.
- Diederichsen A, Hammer K. 2003. The infraspecific taxa of coriander (*Coriandrum sativum* L.). *Gen Resourc Crop Evol*, 50: 33-63.
- dos Santos ÉRQ, Maia CSF, Junior EAF, Melo AS, Pinheiro BG, Maia JGS. 2018. Linalool-rich essential oils from the Amazon display antidepressant-type effect in rodents. *J ethnopharmacol*, 212: 43-49.
- Duke JA (2002) *Handbook of medicinal herbs*. CRC press.
- Elahdadi-Salmani M, Khorshidi M, Ozbaki J. 2015. Reversal effect of *Coriandrum sativum* leaves extract on learning and memory deficits induced by epilepsy in male rat. *Zahedan J Res Med Sci*, 17: e24475.
- Elisabetsky E, Brum LS, Souza D. 1999. Anticonvulsant properties of linalool in glutamate-related seizure models. *Phytomedicine*, 6: 107-113.
- Emam GM, Heydari HG. 2008. Effect of extract and essential oil of *Coriandrum sativum* seed against pentylentetrazole-induced seizure. *Pharm Sci*, 3:1-10.
- Emamghoreishi M, Heidari-Hamedani G. 2010. Anticonvulsant effect of extract and essential oil of *Coriandrum sativum* seed in conscious mice. *Iran J Pharm Res*, 3 (Supplement 1): 71-71.
- Emamghoreishi M, Heidari-Hamedani G. 2015. Sedative-hypnotic activity of extracts and essential oil of coriander seeds. *Iran J Med Sci*, 31: 22-27.
- Emamghoreishi M, Khasaki M, Aazam MF. 2005. *Coriandrum sativum*: evaluation of its anxiolytic effect in the elevated plus-maze. *J ethnopharmacol*, 96: 365-370.
- Enas AK. 2010. Study of the possible protective and therapeutic influence of Coriander (*Coriandrum sativum* L.) against neurodegenerative disorders and Alzheimer's disease induced by aluminum chloride in cerebral cortex of male Albino rats. *Nat Sci*, 8: 202-213.
- Ertas O, Guler T, Ciftci M, Dalkilic B, Yilmaz O. 2005. The effect of a dietary supplement coriander seeds on the fatty acid composition of breast muscle in Japanese quail. *Revue Méd. Vét*, 156: 514-518
- Gastón MS, Cid MP, Vázquez AM, Decarlina MF, Demmel GI, Rossi LI, Aimar ML, Salvatierra NA. 2016. Sedative effect of central administration of *Coriandrum sativum* essential oil and its major component linalool in neonatal chicks. *Pharm Biol*, 54: 1954-1961.
- Ghorbani A, Rakhshandeh H, Asadpour E, Sadeghnia HR. 2011. Effects of *Coriandrum sativum* extracts on glucose/serum deprivation-induced neuronal cell death. *Avicenna J Phytomed*, 2: 4-9.
- Gilani AH. 2005. Trends in ethnopharmacology. *J ethnopharmacol*, 100: 43-49.
- Hosseinzadeh H, Madanifard M. 2000.

Neuroprotective effects of *Coriandrum sativum*

- Anticonvulsant effects of *Coriandrum sativum* L. seed extracts in mice. *Arch Iran Med*, 3: 182-184.
- Iwatani Y, Arcot J, Shrestha AK. 2003. Determination of folate contents in some Australian vegetables. *J Food Comp Anal*, 16: 37-48.
- Kandlakunta B, Rajendran A, Thingnganing L. 2008. Carotene content of some common (cereals, pulses, vegetables, spices and condiments) and unconventional sources of plant origin. *Food Chem*, 106: 85-89.
- Kansal L, Sharma V, Sharma A, Lodi S, Sharma S. 2011. Protective role of *coriandrum sativum* (coriander) extracts against lead nitrate induced oxidative stress and tissue damage in the liver and kidney in male mice. *Int J Appl Biol Pharm Technol*, 2: 65-83.
- Karami R, Hosseini M, Mohammadpour T, Ghorbani A, Sadeghnia HR, Rakhshandeh H, Vafae F, Esmaeilzadeh M. 2015. Effects of hydroalcoholic extract of *Coriandrum sativum* on oxidative damage in pentylenetetrazole-induced seizures in rats. *Iran J Neurol*, 14: 59-66.
- Kharade S, Gumate D, Patil V, Kokane S, Naikwade N. 2011. Behavioral and biochemical studies of seeds of *Coriandrum sativum* in various stress models of depression. *Int J Curr Res Rev*, 3: 4-11.
- Kim K, Lee H, Min S, Seol G. 2015. Neuroprotective Effect of (-)-Linalool against Sodium Nitroprusside-Induced Cytotoxicity. *Med chem*, 5: 178-182.
- Kishore K, Siddiqui N. 2003. Anti-despair activity of green and dried plant extract of *Coriandrum sativum* in mice. *Indian drugs*, 40: 419-421.
- Koppula S, Choi C. 2012. Anti-stress and anti-amnesic effects of *Coriandrum sativum* Linn (Umbelliferae) extract—an experimental study in rats. *Trop J Pharm Res*, 11: 36-42.
- Linck V, Da Silva A, Figueiró M, Caramão E, Moreno P, Elisabetsky E. 2010. Effects of inhaled Linalool in anxiety, social interaction and aggressive behavior in mice. *Phytomedicine*, 17: 679-683.
- Mahendra P, Bisht S. 2011. Anti-anxiety activity of *Coriandrum sativum* assessed using different experimental anxiety models. *India j pharmacol*, 43: 574-577.
- Mani V, Parle M. 2009. Memory-enhancing activity of *Coriandrum sativum* in rats. *Pharmacologyonline*, 2: 827-839.
- Mani V, Parle M, Ramasamy K, Majeed A, Bakar A. 2011. Reversal of memory deficits by *Coriandrum sativum* leaves in mice. *J Sci Food Agric*, 91: 186-192.
- Mehri S, Meshki MA, Hosseinzadeh H. 2015. Linalool as a neuroprotective agent against acrylamide-induced neurotoxicity in Wistar rats. *Drug chem toxicol*, 38: 162-166.
- Mir H. 1992. *Coriandrum sativum* In: Application of plants in prevention and treatment of illnesses. *Persian*, 1: 257-252.
- Mohan M, Yarlagadda S, Chintala S. 2015. Effect of ethanolic extract of *Coriandrum sativum* L. on tacrine induced orofacial dyskinesia. *India j exp biol*, 53: 292-296.
- Msaada K, Hosni K, Taarit MB, Chahed T, Kchouk ME, Marzouk B. 2007. Changes on essential oil composition of coriander (*Coriandrum sativum* L.) fruits during three stages of maturity. *Food Chem*, 102: 1131-1134.
- Nejad Ebrahimi S, Hadian J, Ranjbar H. 2010. Essential oil compositions of different accessions of *Coriandrum sativum* L. from Iran. *Nat prod res*, 24: 1287-1294.
- Nurzyńska-Wierdak R. 2013. Essential oil composition of the coriander (*Coriandrum sativum* L.) herb depending on the development stage. *Acta Agrobotanica*, 66: 53-60.
- Orav A, Arak E, Raal A. 2011. Essential oil composition of *Coriandrum sativum* L. fruits from different countries. *J Essent Oil Bearing Plant*, 14: 118-123.
- Pandey A, Bigoniya P, Raj V, Patel K. 2011. Pharmacological screening of *Coriandrum sativum* Linn. for hepatoprotective activity. *J Pharm Bioall Sci*, 3: 435-441.
- Park H, Seol GH, Ryu S, Choi I-Y. 2016a. Neuroprotective effects of (-)-linalool against oxygen-glucose deprivation-induced neuronal injury. *Arch pharm res*: 1-10.
- Park H, Seol GH, Ryu S, Choi I-Y. 2016b. Neuroprotective effects of (-)-linalool against oxygen-glucose deprivation-induced neuronal injury. *Arch pharm res*, 39: 555-564.
- Pathan A, Kothawade K, Logade MN. 2011. Anxiolytic and analgesic effect of seeds of *Coriandrum sativum* Linn. *Int J Res Pharm Chem*, 1: 1087-1099.
- Pourzaki M, Homayoun M, Sadeghi S, Seghatoleslam M, Hosseini M, Bideskan

- AE. 2017. Preventive effect of *Coriandrum sativum* on neuronal damages in pentylentetrazole-induced seizure in rats. *Avicenna J Phytomed*, 7: 116-128.
- Rakhshandeh H, Sadeghnia HR, Ghorbani A. 2012. Sleep-prolonging effect of *Coriandrum sativum* hydro-alcoholic extract in mice. *Nat prod res*, 26: 2095-2098.
- Sabogal-Guáqueta AM, Osorio E, Cardona-Gómez GP. 2016. Linalool reverses neuropathological and behavioral impairments in old triple transgenic Alzheimer's mice. *Neuropharmacology*, 102: 111-120.
- Sahib NG, Anwar F, Gilani AH, Hamid AA, Saari N, Alkharfy KM. 2013. Coriander (*Coriandrum sativum* L.): A potential source of high-value components for functional foods and nutraceuticals-A review. *Phytother Res*, 27: 1439-1456.
- Souto-Maior FN, de Carvalho FL, de Moraes LCSL, Netto SM, de Sousa DP, de Almeida RN. 2011. Anxiolytic-like effects of inhaled linalool oxide in experimental mouse anxiety models. *Pharmacol Biochem Behav*, 100: 259-263.
- Tashinen J, Nykänen L. 1975. Volatile constituents obtained by the extraction with alcohol-water mixture and by steam distillation of coriander fruit. *Acta Chem Scand*, 29: 425-429.
- Vekaria RH, Patel MN, Bhalodiya PN, Patel V, Desai TR, Tirgar PR. 2012. Evaluation of neuroprotective effect of *Coriandrum sativum* linn. against ischemic-reperfusion insult in brain. *Int J Phytopharmacol*, 3: 186-193.
- Velaga MK, Yallapragada PR, Williams D, Rajanna S, Bettaiya R. 2014. Hydroalcoholic seed extract of *Coriandrum sativum* (Coriander) alleviates lead-induced oxidative stress in different regions of rat brain. *Biol trace element res*, 159: 351-363.
- Wangensteen H, Samuelsen AB, Malterud KE. 2004. Antioxidant activity in extracts from coriander. *Food Chem*, 88: 293-297.
- Xu P, Wang K, Lu C, Dong L, Gao L, Yan M, Aibai S, Yang Y, Liu X. 2017. Protective effects of linalool against amyloid beta-induced cognitive deficits and damages in mice. *Life Sci*, 174: 21-27.
- Yeung EC, Bowra S. 2011. Embryo and endosperm development in coriander (*Coriandrum sativum*). *Botany*, 89: 263-273.
- Zargar-Nattaj SS, Tayyebi P, Zangoori V, Moghadamnia Y, Roodgari H, Jorsaraei SG, Moghadamnia AA. 2011. The effect of *Coriandrum sativum* seed extract on the learning of newborn mice by electric shock: interaction with caffeine and diazepam. *Psychol Res Behav Manag*, 4: 13-19.
- Zargari A. 1991. Medicinal plants. 5th ed. Tehran University Publications, 2: p. 1-942.